# Exploring Maths at Advanced Level Maths Activities for the HP Prime and HP39gII





# Contents

1.	Studying Advanced Level Maths with an HP Prime or HP39gII Graphing Calculator	3
2.	Basic Operation	4
3.	Graphs and Functions	8
4.	Matrices	14
5.	Sequences and Series	16
6.	Projectiles. Parametric Functions	20
7.	Statistics: Probability Distributions and Inference	22
8.	Calculus	28

#### Introduction

This book is aimed at users of the HP Prime and HP39gII graphing calculators, who are studying for an advanced level exam in mathematics (English A Levels, Scottish Highers, French and International Baccalaureat, German Abitur, Netherlands HAVO and VWO 3/4 level and equivalent in other countries). The aim is to provide a range of activities which will help you become a confident user of the calculator while developing your skills in different mathematical topics. There is no attempt to cover the entire course, but there is a good range of topics covered. The power of this technology is in its capacity to generate lots of mathematical information very quickly, so you can get a good feeling for mathematical ideas. Mathematics needs to be explored and with your HP Prime or HP39gII you can get under the skin of the ideas you need to learn about. I hope that you will try out the activities in this booklet and get into the spirit of exploration that you can then use in all of the topics you need to study.

**NOTICE**: There are significant differences between the HP Prime and the HP39gII. Most important is that the HP Prime has an inbuilt Computer Algebra System (CAS). In this booklet we will not engage with the CAS at all. There will be a separate booklet for activities using CAS. There are also some PRIME only Apps, notably dynamic geometry and spreadsheet which will not be included in this booklet. The remainder of the Apps and the non-CAS operation are broadly similar between HP Prime and HP39gII and this booklet will make use of these areas only.

### About the Author

Chris Olley was a secondary school maths teacher in a range of comprehensive schools in London and East Africa. He currently directs the secondary maths PGCE course at King's College London. He has worked with graphing calculators since they first arrived in the late 1980s and has run sessions nationally and internationally on different approaches to dynamic ICT in maths education, of which graphing calculators are an excellent example.

#### **Further Information and Support**

Please visit www.hpgraphingcalc.org and join in the discussion threads. Share your ideas and your understanding of mathematics with others.

# 1. Studying Advanced Level Maths with an HP Prime or HP39gII Graphing Calculator

The HP Prime and HP39gII are very powerful calculators. They can find solutions to a vast range of mathematical problems in algebra, calculus, probability and statistics, complex numbers, matrices and much more. Press the Suitcase Key on the HP Prime or the Math key on the HP39gII to see the menu of commands and get a feeling for how much this machine will do.

The big message for advanced level maths students, is that you can use this machine in your exams (those for which a calculator is allowed). I would assume you have bought the calculator because you know this. It is really important that you do not think that the calculator will answer the exam questions for you. You actually have to do the steps in the process. However, the main source of difficulty in an advanced level exam is making small errors along the way. With these calculators you can quickly calculate the solution and check that you have done it correctly and be able to move on confidently. Knowing the answer first is often a very helpful way of deciding what steps to take. Also, being able to see different representations of an object, quickly, helps you decide what the solution could be. With these calculators you can draw graphs and see tables of



values of different types of functions, which you can zoom in and out of, make calculations on, even do numeric integration and differentiation.

So, they won't tell you how to solve the problem, but will get you confidently to a solution with a range of better ways of seeing. These are a fantastically powerful tools and all of the evidence from different countries where graphing calculators are used in exams says that if you can use it properly, then it will give you an advantage.

However, it's not just in the exam that the calculator is useful. While you are learning maths, it is vital that you get a good intuitive feeling about how maths works. These calculators are fantastic tools for exploring mathematics. What happens to the graph of a quadratic  $y = ax^2 + bx + c$  when you change the coefficients a, b and c? You would need to draw dozens of graphs to get a good feeling for what goes on. With these calculators you can draw as many as you like, changing the coefficients selectively. Working this way helps you get out of the other big problem with advanced level exams. Memorising methods is fine until you find a question you can't recall the method for.

That normally means most of them, because they never come up just like you were expecting. Far better to understand what is going on and be able to see the mathematics from different starting points, then you can work your way through even when you can't decide which method to use. Maths is the method and you make yourself a mathematician, by exploring. The HP Prime and HP39gII are the tools to make that possible.

This book has a range of examples. Many of them will suggest areas of maths you can explore. But remember, they are just examples. Use the ideas to explore any new area of maths you are learning. By the time of your exam you will be so skilled in using the calculator that it will be able to support you quickly and powerfully in the exam itself.



# 2. Basic Operation

# The Home Screen and the Suitcase (Prime) or Math (HP39gII) Menu

When you switch the calculator on, you are in the Home screen. Here you can do any type of calculation. Not just arithmetic, but using matrices, summing sequences, calculus, complex numbers and so on. Pressing the Suitcase (Prime) or Math (39gII) key shows you the menu of all of the things you can do on the Home screen. Notice that the Math menu has several sub menus, so make sure that the MATH submenu is selected (by touching Math (Prime) or pressing F1 (39gII)). Navigate with touch (Prime) or the cursor. For details of syntax press the Help key (Prime) or note that the bottom of the menu shows you how to enter things when more than one input is needed (39gII).

### **Direct Entry versus Template Entry**

On the HP39gII, to do a numeric differential we find the  $\partial$  command which tells us to enter an expression i.e. the function, then a value, i.e. the value at which we are calculating the differential, i.e. the slope.

Let's calculate the differential of  $x^3$  at x = 7. We know the differential is  $3x^2$ , so at x = 7 the result should be  $3 \times 7^2 = 147$ .

You will have used the X,T, $\theta$ ,N key to enter the X, the  $x^{y}$  key for the power and the = key (shift and decimal point) to set the value.

On the HP Prime we have templates for many mathematical functions. Press the template key to the right of the suitcase. Choose the differential. Use the X,T, $\theta$ ,N and the  $x^{\gamma}$  key to enter the expression then press Enter.

You will get an odd answer here because it will have been evaluated with the whatever value X has.



We wanted X=7, so we must say so. Enter 7 and press the Sto button on the screen and Enter.

Now press on the differential you entered before and Copy (on the screen), then Enter. Now we have evaluated the differential at X=7 (giving the same answer as we got on the HP39gII). The moral here is to enter the values of variable before you use a template.

en al de la desarro de la construcción de la construcción de la construcción de la construcción de la construc	Function	un nu ser ser ser ser ser ser se se ser se
3		
θX		
дX		.03
7 <b>►</b> X		7
3		
θX		
ЭХ		147
Sto 🕨		Copy Show

18:38 6

OK. Now get started. Explore! Find any functions you already know and make them work for you. Then explore ones you've vaguely heard about.

(If you are doing Physics and/or Chemistry as well, then you will be tempted to check the UNITS and PHYS menus as well. These are on the Math menu on the HP39gII. On the HP Prime they are under Units, the shift option on the template key. I suggest you save this till later, but it is good to see that for scientific work, this calculator will make calculations with correct units attached. It also knows all of the key constants scientists need to know).

## Working with Apps

The home screen is where you do calculations. You can always get back to the home screen by pressing the Home button. Pretty much everything else happens through Apps. Press the Apps key and scroll down the list. You will see most of your advanced level topics are covered. All of the Apps work the same way. The Symb, Plot and Num keys show you the three representations of all the maths you can explore with Apps. Symb is for the algebraic view, Plot shows the graph view and Num shows the numeric or table view. Different Apps start in different views according to the maths, for example graphing Apps start in Symb mode, so you can enter a function, but Statistics Apps start in **Num** mode, so you can enter the data. The bar at the top of the screen tells you which App is running at any time. If you start up a new App, all of your data from the last App you were using is saved automatically and you can come back to it any time.

You can change the way each of these representations looks using the Setup option on each key (using the SHIFT key). Explore to see what the possibilities are. Finally, you can change the way the screen is organised using the Views menu. For example, you can split the screen and see the graph and table of values at the same time.

ZOOM TRACE GOTO

Figure 3: The Polar App

There are three main types of Apps:

1. Function Apps. Here you enter the data or the function without restriction.

2. Solver Apps. Here you get fast solutions to specifics types of problems.

3. Explorer Apps. Here you get a

pre-configured set up to make



DEEN MENU

your understanding.

Figure 5: The Quadratic Explorer App

X=0

Eq Graph• Incr 1 Lev 4 Test

HP calculator users are a great enthusiast community and there are additional Apps created and being created for you to download and add to the functionality of your machine.

So, check regularly to see what is available, and of course, contribute yourself.

Check at:

- www.hpgraphingcalc.org which is a general purpose support site
- www.calc-bank.com which has programmes and activities

## **Hints and Tips**

- If you are stuck on a menu press ON to cancel menus. The HP Prime has an Esc key which does the same from a more convenient place.
- This calculator has detailed help built in. Press the Help button (Shift Views on the HP39gII) and it will give you help on whatever you were doing.
- On the HP39gII you can show mathematics correctly, like this ... Press Modes (SHIFT Home), Press F4 and check the Textbook display option. (HP Prime defaults to this mode)
- You can save apps with your settings preserved and attach notes to remind yourself what you were doing. Press Info (SHIFT Apps) to add the notes. Then select the App in the Apps menu. Click F1 for SAVE and give it a new name. You can then share this with anyone else, or indeed online.
- Install the connectivity software onto your computer. This way, the main operating system of the calculator can be updated when changes become available, you can upload and download Apps and you can type notes for your Apps or Programme your calculator much more efficiently with a keyboard!

RAD	Function	
10 ∑X		
X=1		55
STO ►		
0.00		
	Quadratic Explorer	18:53 47
10 \(v)		
X=1		55

# 3. Graphs and Functions

## **Linear Functions**

It is very important that you are able to look at a function and have a good feel for the size, shape and position of the graph of that function. Your HP Prime or HP39gII are the perfect tools for exploring functions.

In the first instance you should do this for functions of the form y = f(x) using the Function App. Later on you may need to do the same for polar functions, parametric functions and sequence functions, depending on the modules you are taking. There are Apps for all of these, but for now, let's focus on the Function App. Press the Apps key, choose Function, press RESET and OK to confirm, then START.

You should be very familiar with (linear) functions of the form f(x) = mx + c. You will know that these are straight line graphs and that they have a gradient of m and they cross the y-axis at (0, c) which is called the y-intercept.

First you should make sure you know the relationship between the gradients of linear functions whose graphs are:

- Parallel
- Perpendicular

Your calculator is ready to receive an input for a function F1(x), so try a simple Linear Function, say F1(x) = 3x + 1

Just type 3, press F3 for the X (or press the X,T, $\theta$ ,N key or press ALPHA X), then +, then 1, then press F6 for OK.

(Notice that the F1 now has a tick next to it, which means that when we choose Plot or Num modes, F1 will show up. We can use F2 for  $\checkmark$  CHK to turn functions on and off)

Try a function for F2 which will be parallel to F1. The cursor has already moved down to F2, so you can type this directly. Now press the Plot key to see if you are correct.

Press the Symb key, make sure the cursor is on the F2 line and try a new function for F2. Keep trying different functions until you are sure you know the relationship between the gradients.

You will need to explore different families of functions:

Polynomial functions f(x) = ax + b  $f(x) = ax^{2} + bx + c$   $f(x) = ax^{3} + bx^{2} + cx + d$ (etc.)

Trigonometric functions f(x) = sin(x) f(x) = cos(x)f(x) = tan(x)

Reciprocal Functions  $f(x) = \frac{1}{x}$ 

Exponential Functions  $f(x) = a^x$ 

	 mouth including the	Function Symbolic View 18:	58 D
V	F1(X)=	3*X+1	
	F2(X)=		
	F3(X)=		
	F4(X)=		
	F5(X)=		
	F6(X)=		
	F7(X)=		



Now do the same thing for Linear functions whose graphs are perpendicular.

I'm sure you knew the relationships already. But this will have helped you get used to entering and changing functions and looking at their graphs.



## **Quadratic Functions**

Next, you should explore Quadratic functions. There are three standard ways of expressing a quadratic function:

- 1. Polynomial:  $f(x) = ax^2 + bx + c$
- 2. Factorised: f(x) = (x + a)(x + b)
- 3. Completed square:  $f(x) = (ax + b)^2 + c$

The first one is most familiar, but the factorised form gives the most information quickly (if the quadratic can be factorised). The completed square form probably gives the most complete picture,

but the algebra involved in changing to this form is trickier. The important thing is that all three give different insights into the nature of the function. Your calculator will not do the algebra for you, so you will have to practice converting between these three forms with pencil and paper.

Press Apps, select Function, then RESET/OK/START and enter a function F1 in factorised form, say F1(x) = (x - 3)(x + 5) press OK, then press Plot.

Press Menu, then Fcn. Select Root.

Now move the cursor so it is closer to the second root (clicking left a fair few times will do it). Now press Menu/Fcn/Root again.

The relationship between the roots and the factorised form is pretty clear. Try a few more examples to make sure.



Now get a feel for the completed square form. Replace your F1 with a function in completed square form, say  $F1(x) = (x + 3)^2 - 4$  (use the  $x^2$  key to enter the 'squared'). Now look at the position of the minimum point with Menu/Fcn/Extremum

Also, check the positions of the roots as before.



You will need to do some exploration before you can be sure of the relationship between the roots, the extremum and the values of a, b and c in  $f(x) = (ax + b)^2 + c$ , so keep going until you are quite sure. Then test your theory on new functions in this form.

Now you should be ready to take on the task of exploring the effect of changing *a*, *b* and *c* in the form  $f(x) = ax^2 + bx + c$ . Work on each one independently. The effect of changing the c is quite straightforward. The effect of changing the *a* is a bit counter-intuitive. However, just changing the *b* on its own is guite hard to describe. So, spend some time exploring until you are guite sure you have a good theory that you can describe accurately and simply.

#### **Polynomials**

Now you should explore other functions. Firstly, get a feel for cubic functions. Remember that different algebraic forms give different insights into the nature of the function. It is quite hard to imagine the graph of a cubic in the form  $f(x) = ax^3 + bx^2 + cx + d$  but much easier in the form f(x) = (x + a)(x + b)(x + c), so factorising first, if possible, is always a good thing.



Try some quartic and quintic functions as well, to get a feeling for the differences between different polynomial functions.

### **Other Functions**

You should explore trigonometric, reciprocal and exponential functions in the same way.

When you have a good feel for the general shape and position of these functions, you can explore how the graphs of functions can be manipulated generally.

The best way into this is to have a good play with the graph explorer apps: Quadratic Explorer and Trig Explorer.

Press Apps and select Quadratic Explorer. (Best to Reset then Start)



Use the cursor to move the graphs up, down, left and right and spread the graph in and out with the + and - keys. Look at the effect this has on the function.

Work through the 4 levels of difficulty (LEV1 to LEV4) and TEST yourself.

Now do the same with the Trig Explorer applet.



## Practice until these conclusions are clear:

A vertical shift of *a* in the:

+ve direction f(x) + a

-ve direction f(x) - a

A horizontal shift of a in the:

+ve direction f(x - a)

-ve direction f(x + a)

A vertical stretch of a: +ve (increase) af(x)-ve (decrease)  $\frac{1}{a}f(x)$ A horizontal stretch of : +ve (increase)  $f\left(\frac{1}{a}x\right)$ -ve (decrease) f(ax) Now you should explore every different function type. Start with a reciprocal function  $f(x) = \frac{1}{x}$  and ensure that you can translate and stretch it horizontally and vertically. Repeat with an exponential function  $f(x) = a^x$  and use HOME/Sto/ALPHA/A to change the value of A.





# **Piecewise functions**

We can graph piecewise functions i.e. ones that are defined differently over different part s of the domain.

For example, we can graph a function for which f(x) = 2x when x < 1 and f(x) = 2 when  $x \ge 1$ 

We write this function as  $f(x) = \begin{cases} 2x & x < 1 \\ 2 & x > 1 \end{cases}$ 

Press Apps and launch the Function applet.

(Reset/OK/Start)

For F1(X) you should

(HP39gII) enter 2\*X then press SHIFT Vars (to get the Chars menu) and find the back slash / character. Click OK.

Now enter the domain in brackets (X<2) getting the < from the Chars menu. Click OK to see the function and the domain shown. Now enter the second part of the function as F2(X)=2

You can find a  $\geq$  symbol but it is easier to find >= which means the same.

RAD	Function Syr	nbolic View		RAD Func	tion Syn	nbolic View		RAD	Fund	ction Syl	mbolic View		
F1(X)=				✓F1(X)= 2*X				<b>√</b> F1(X	)= 2*X			[	
F2(X)=				(X<1)	Ī				X<1	ĩ			
F3(X)=				$F_2(x) = $				✓F2(X)	)= 2	,		[	
F4(X)=				F3(X) =				11200		1			
F5(X) =			-	1000					(^	IJ.			÷
2X/(X<1)				Enter function			_						
	X	CANC	L OK	EDIT ↓ ✔ CHK	Х	SHO'	W EVAL	EDIT	√СНК	Х	SH	OW EVAL	

 $\square$  and enter 2X then the domain X<2 us ing the menu with (HP Prime) use the template menu shift and the 6 key.

·····		Functi	on Syn	nbolic	View	11 11 11	20:20	Function Symbolic View 20:18	Function Symbolic View 20:19
F1(X)=							√ F1(X)= <sup>2</sup> *X X<1	√ F1(X)= <sup>2</sup> *X X<1	
📕 F2	2(X)=						_	F2(X)=	$\sqrt{1}$ F2(X)= $\frac{2}{X \ge 1}$
<b>F</b>	븝		미	<u>90</u> 90	0,0  0,0	問	_	<b>F3(X)=</b>	<b>F3(X)</b> =
F F	$\sqrt{\Box}$	₽□	lim□ n→n <sup>0</sup>	∫90	[00]	[8]		F4(X)=	F4(X)=
F	Inl	Π.□	τη	log []	-°′″			<b>F5(X)=</b>	F5(X)=
F								F6(X)=	F6(X)=
F7	7(X)=							F7(X)=	F7(X)=
Edit	√ √		Х		Sh	ow	Eval		Edit √ X Show Eval

Finally press Plot to see the effect. You should explore setting up different piecewise functions, including more than two parts.



## 4. Matrices

Mainly you will be using matrices for Linear Algebra. The Matrix button (SHIFT + 4) is the editor. Here you can create Matrices in a convenient way. You can also type matrices directly onto the Home screen. When you have created a matrix, you should store it as a variable like M1, M2 etc.

Let's do a task. Solve the linear system:

$$2x_1 + 3x_2 - 5x_3 = 13$$
  

$$x_1 - 3x_2 + 8x_3 = -13$$
  

$$2x_1 - 2x_2 + 4x_3 = -6$$



Use the Matrix editor to enter the matrix of coefficients.

This matrix is already called M1 by the editor. Just click Edit to get started and use the cursor keys and OK to enter each value. The Editor quickly works out the size of the matrix. When you are finished, press Home.

To solve the problem we pre-multiply the solution matrix, by the inverse of the coefficients matrix. So create a second matrix M2 with the right hand sides of the equations. This is a column matrix.

Use the matrix editor to enter M2.



Now we are ready to find the solutions ...

On the Home screen type  $M1^{-1} \times M2$ 

(use the  $x^{y}$  key and type -1)

(HP 39gII) If the Textbook display is turned off, then we can use the Show key (F5) to see the matrix correctly.

M1<sup>-1</sup> \*M2 [1 2 -1] Sto► [1 2 -1] OK

So, we have solutions:  $x_1 = 1, x_2 = 2, x_3 = -1$ 



Press the Suitcase key (HP Prime) or the Math key (HP39gII) to find the Matrix section to see the range of operations we can apply to our matrix.

[2 1 2	9- Loop Matrix Polynomia Probability	Math Fu Il	INCTIONS SVL TRACE TRN 1		·5 3 4
	lobject^-1				
Ν	1AT• UNITS	PHYS	CATLG	CANCL	OK

20:31 Matrices M2 Math 1 Numbers 1 Transpose <sup>2</sup>Arithmetic > 2 Determinant 3Trigonometry > 3RREF I Cross Product 4 Hyperbolic 4 Create 5 Probability 5 Basic 2Dot Product 6 List > 3L<sup>2</sup> Norm 6Advanced 7 Matrix 7 Factorize 4L<sup>1</sup> Norm Special <sup>8</sup>Vector 5Max Norm Math App Catlg OK

Entering the matrices can be a bit time consuming, but when they are entered you have an enormous amount that you can do with the matrices. So, now explore the other matrix functions available on the Math menu.

You can find the determinant.

You can find inverse matrices (notice that M2 has no inverse, so will generate a syntax error if you try this).

(HP39gII) Here we have used the up cursor to highlight the matrix and SHOW to see the matrix correctly.

HP Prime shows the matrix in textbook format by default.

braith doraith alth a th doraith doraith doraith	Function	20:40 <1
D		
DEI(M1)		24.00
-1.00		.17083 .38
M1		.50 .7588
		.17 .4238
Sto 🕨		

 RAD
 Function

 DET(M1)
 24

 DEG
 Function

 M1<sup>-1</sup>
 [-1.5,2.5,-1],[-4.25,5,-1.75],[4.5,-5.5,2]]

 STO ►
 [-1.5 2.5 -1]

 (-1.5 2.5 -1],[-4.25,5,-1.75],[4.5,-5.5,2]]

 STO ►

 (-1.5 2.5 -1]

 (-1.5 2.5 -1)

 (-1.5 2.5 -1)

 (-1.5 2.5 -1)

 (-1.5 2.5 -1)

 (-1.5 2.5 -1)

 (-1.5 2.5 -1)

 (-1.5 2.5 -1)

 (-1.5 2.5 -1)

 (-1.5 2.5 -1)

 (-1.5 2.5 -1)

 (-1.5 2.5 -1)

 (-1.5 2.5 -1)

 (-1.5 2.5 -1)

 (-1.5 2.5 -1)

 (-1.5 2.5 -1)

 (-1.5 2.5 -1)

 (-1.5 2.5 -1)

 (-1.5 2.5 -1)

 (-1.5 2.5 -1)

 (-1.5 2.5 -1)

 (-1.5 2.5 -1)

 (-1.5 2.5 -1)

 (-1.5 2.5 -1)

 (-1.5 2.5 -1)

 (-1.5 2.5 -1)

 (-1.5 2.5 -1)

 (-1.5 2.5 -1)

 (-1.5 2.5 -1)

 (-1.5 2.5 -1)

 (-1.5 2.5 -1)

 (-1.5 2.5 -1)

 (-1.5 2

You can find Eigen Values and Vectors and do row and column operations. You should experiment to get a feel for all of the available functions and operations.

# 5. Sequences and Series

You will be very familiar with different continuous functions. You will have explored the relationship between the function, its graph and tables of values. With your HP39gII you can get a feel for the variation in functions which are not continuous, by exploring sequences and series.

Look at a simple sequence: 3, 5, 7, 9, 11, 13, ...

The first term is 3 and the rule is that each number is 2 more than the previous number (we have  $u_1 = 3$  and  $u_n = u_{n-1} + 2$ ).

(You may know this as an arithmetic progression with a = 1 and d = 2).

Let's look at this with the sequence app.

Press Apps, select Sequence then Reset/OK/Start

U1 is a sequence. We can enter its first term U1(1), its second term U1(2) and a rule for the *n*th term. You only need to enter what is needed. The first and second terms are not necessarily needed. In our case, we have a first term and a rule.

Enter the first term U1(1)=3

Then navigate down to U1(N). Notice that options are added for the (N-1) which we will need.

Use this to enter  $u_n = u_{n-1} + 2$  as U1(N)=U1(N-1)+2

Press the Num key to see the sequence.

I pressed F4 to turn off the BIG font so you can see more of the sequence.

Use the down arrow to see even more.

Notice that using the up arrow will not allow us to go back further than the first term, because we entered a value for U1(1) saying that this is where the sequence starts.

	inna fi seinna fi sinna fi s	Sequence S	ymbolic View	08:51
	U1(1)=			
	U1(2)=			
	U1(N)=			
	U2(1)=			
	U2(2)=			
	U2(N)=			
	U3(1)=			
_				
	dit	$\checkmark$	Sh	iow Eval
		Sequence S	ymbolic View	08:56
	U1(1)=	3		
	U1(2)=			
√	U1(N)=	U1(N-1)+2		
	U2(1)=			
	U2(2)=			
	U2(N)=			
	U3(1)=			
	-die	/	L Ch	our Fuel
	N	 U1		
1		3		
2 3		5 7		
4		9		
5 6		13		
7		15		
9		19		
10		21 23		
1		2.5		

Size

Defn Column

Zoom

Now press Plot to see what the sequence looks like graphically.



Press Shift Plot to see the plot setup options.

I have changed the plot type to Cobweb and increased the end of the Y range (YRNG) to 50 to see more of the sequence.

We have specified the sequence with a term-to-term rule. Now we can compare with a position to term rule, i.e. a rule for the *n*th term.

The first term is three and the sequence is increasing by two for each extra position so the position-to-term rule must be u(n) = 2(n-1) + 3. Notice it must be n-1because the first term has zero 2s added on.

Press Symb and enter this rule as the second sequence U2. Enter the rule as U2(N)=2(N-1)+3

(On the HP39gII notice that when you do this U2(1) and U2(2) are greyed, because with a position-to-term rule, any value can be calculated, so the initial terms are irrelevant).

Press Num and compare the sequences.

-		Sequence	e Symbolic	View		09:39
	U1(2)=	5				
$\checkmark$	U1(N)=	U1(N-1)+2	2			-
	U2(1)=					
	U2(2)=					
√	U2(N)=	2*(N-1)+3				
	U3(1)=					
	113(2)=					
E	dit	√ [		Sł	iow	Eval
E	dit N	√ U1	U2	Sł	iow [	Eval
1 2	dit N	√ U1 3	U2	[ Sŀ	iow [	Eval
1 2 3	N	✓ U1 3 5 7	U2 3 5 7	Sł	iow	Eval
1 2 3 4 5	N	✓ U1 3 5 7 9 11	3 5 7 9 11	Sł	iow	Eval
1 2 3 4 5 6	idit N	✓ U1 3 5 7 9 11 13	U2 3 5 7 9 11 13	Sł	iow	Eval
1 2 3 4 5 6 7 8	N	U1 3 5 7 9 11 13 15 17	U2 3 5 7 9 11 13 15 17	Sł		Eval
E 1 2 3 4 5 6 7 8 9	N	U1 3 5 7 9 11 13 15 17 19	U2 3 5 7 9 11 13 15 17 19 21	Sł		Eval

This provides an excellent platform to explore different sequences and practice finding position-to-term rules to match term-to-term rules.

Now that we know the position-to-term rule, we can find values for the corresponding series.

Zoom

Size Defn Column

To find 3+5+7+9+11, we are summing terms in the sequence from n = 1 to n = 5

So, we find  $\sum_{1}^{5} 2(n-1) + 3$ 

(HP39gII) Press Math, navigate to Loop and choose  $\Sigma$ 

Note that on the menu, it tells you how to enter the values. Click OK, then enter the expression 2(n - 1) + 3 using the ALPHA key to enter the N. The *var* is N and the limits are 1 then 5. Make sure to put the commas in between each entry.

Now press ENTER to evaluate.

(This is how it appears with textbook display switched on).

HP Prime) Use the template key

Check that you agree with the result! (3+5+7+9+11 = 35)

You can easily change the values by using the cursor to select the entry, then pressing Copy and editing. You should explore the effect of making changes to the limits and to the rule.

You can easily explore more complicated sequences. An

excellent example would be Fibonacci sequences.

1, 1, 2, 3, 5, 8, 13, ...

Here, we know the first and second terms and a rule that each term is the sum of the preceding two.

Relaunch the sequence applet and reset it, ready to enter the new sequence.

We have U1(1)=1, U1(2)=1 and U1(N)=U1(N-1)+U1(N-2)

Check that it has worked by looking at the table.

(Press Num)

8-2 Math Functions	
Hyperbolic IITERATE	
Integer <b>S</b>	
List	
Loop	
Σ( expr, var, ivalue, fvalue)	
MAT• UNITS PHYS CATLG CANCL	OK
RAD Sequence	
$\Sigma(2(N-1)+3.N.1.5)$	
STO F	
RAD Sequence	
5	
5 3*(N 4)+ 3	
2*(IN=1)+3	
Ň=1	35
	- 35
STO F	
	10:04-
Sequence	10:04



Size

Defn Column

Zoom



A square is divided into quarters and one quarter is shaded. A quarter of the opposite unshaded quarter is shaded. A quarter of this quarter is shaded and so on as shown. What fraction of the original square is shaded?

If the original square has unit area, then the shaded part is  $\frac{1}{4} + \frac{1}{16} + \frac{1}{64} + \cdots$ 

That is, adding up the terms of the sequence with  $u_n = \left(\frac{1}{4}\right)^n$ 

So, see what happens to this value as n increases.

Enter  $\sum_{1}^{1} \left(\frac{1}{4}\right)^{n}$ 

Press  $a \leftrightarrow b/c$  to see the outcome as a fraction.

Use the up arrow to COPY and EDIT the sum. Keep increasing the last term to see how the sum is changing. What value does it tend towards as the number of terms increases?

Try to explain how you know that the shaded area must be this fraction of the whole.



Sequence

10:12

# **6: Projectiles. Parametric Functions**

We can set up graphs with variable coefficients. In this way we can explore the effect of changing the coefficients. A good example of this is to graph a general projectile.

We know the horizontal equation of motion is x = utcos(A)

The vertical equation is  $y = utsin(A) - 0.5gt^2$ 

Where u is the initial velocity, A the angle of projection, g acceleration due to gravity and t the time taken.

So, we have two parametric equations with parameter t

First, we enter the equations: (note that T is the parameter so use the X,T, $\theta$ ,N key).

Press Apps, find Parametric, press Reset then OK, then Start

For X1(T) type ALPHA U  $\times$  T  $\times$  COS(ALPHA A) then OK

For Y1(T) type ALPHA U  $\times$  T  $\times$  SIN(ALPHA A) - 0.5  $\times$  ALPHA G  $\times$  T2 then  $\,$  OK

If you press Plot now, you will see nothing, since the value of	
all of the variables defaults to zero. So, now we assign values	
to the variables. First, we will take $g = 9.8$	

Press HOME, type 9.8 press STO ALPHA G ENTER

Now choose some sensible values for the other variables, (say):

30 STO ALPHA A ENTER

#### 5 STO ALPHA U ENTER

Press (HP Prime) Shift Home Settings or (HP39gII) SHIFT Modes and set the Angle Measure to degrees.

Now press Plot again.

	والمحاو المحاو المح	Parametric Symbolic View	10:16
V	X1(T)=	U*T*COS(A)	
	Y1(T)=	U*T*SIN(A)5*G*T²	
	X2(T)=		
	Y2(T)=		
	X3(T)=		
	Y3(T)=		
	X4(T)=		
	dit	/ T Show	Eval

Laire	× ×		511011	L VOI
		Paramet	ric	10:17



We can go back to the original zoom level using ZOOM Decimal.

Having set this up you can now explore changing the variables. How much difference does it make if we double the initial angle?

Press Home and 60 STO ALPHA A ENTER, then Plot.

We can solve sloping ground problems. For example, suppose the projectile is launched from a point on a slope with gradient 0.2. We can model this with a function f(x) = 0.2x. However, we are in parametric mode, so we use the parameter T for X as x(t) = t and then y(t) = 0.2tso we enter X2(T)=T and Y2(T)=0.2T.

Press Plot to look at the motion.

Now use zoom box to find out when the projectile lands.

First use zoom box to zoom in close on the landing point. Then TRACE to find the value of T. You can increase the precision by changing the T steps using SHIFT Plot. To 2 d.p. the projectile lands after 7.82 seconds.



X: 7.81880877743 Y: 1.56146788991 Menu

Finally you can even see what happens going back in time! Press SHIFT plot to Setup the plot. The T range defaults to starting at time T=0, which is logical. But change the TRNG to start in the past e.g. T=-2 ...

# 7. Statistics: Probability Distributions and Inference

For statistical work, you should use the three statistics applets:

- 1 variable statistics (for box plots, histograms etc. with central tendency and spread)
- 2 variable statistics (for scatter plots, correlation and regression)
- Inference (for inferential statistics and confidence intervals)

Also, press Math and look at the following sub menus:

- Distribution (binomial, normal, poisson, etc.)
- Probability (permutations, combinations, random numbers, etc.)

#### **One Variable Data**

Get started by entering some data for 2 single variable lists.

Apps/Statistics 1Var/Reset/OK/Start

For example, the ages of visitors to a swimming pool on Friday are as follows:

{12, 13, 12, 12, 12, 12, 3, 38, 12, 13, 12, 13, 12, 12, 12, 13} enter this under D1

... and on Saturday are as follows:

{3, 56, 23, 12, 14, 15, 7, 5, 35, 28, 17, 2, 6, 15, 21} enter this under D2

Press Symb and change Plot 1 and Plot 2 from Histogram to BoxWhisker (notice the wide range of other plot types ... experiment to see what they are like)

Also, enter D2 for the list associated with plot H2.

Press Plot (then Menu/Zoom/Autoscale) to compare the two distributions.

Press Num and Stats to see the summary statistics. (HP39gII) Press down to see mean and standard deviation(s).

Click OK to exit the STATS view.

	Sta	atistics 1Var N	lumeric Viev	V 10:29
	D1	D2	D3	D4
7	3	7		
8	38	5		
9	12	35		
10	13	28		
11	12	17		
12	13	2		
13	12	6		
14	12	15		
15	12	21		
16	13			

Edit Ins	Sort	Size Make	Stats
Stati	stics 1Var S	Symbolic View	10:30
√ H1: D1		Freq	
Plot1: Box\	Whisker		×
√ H2: D2		Freq	
Plot2: Box\	Whisker		Ŧ
H3:			
Plot3: Histo	ogram		Ŧ
H4:			



	_										<u> </u>						_			<u> </u>			_			
																										1

Zoom	race•		Jefn Menu
Х	H1	H2	
n	16	15	
Min	3	2	
Q1	12	6	
Med	12	15	
Q3	13	23	
Max	38	56	
ΣΧ	213	259	
ΣX <sup>2</sup>	3569	7317	
x	13.3125	1.7266667E	1
sX	6.992555565	1.4255158E	1
σX	6.770512813	1.3771791e	1
16	I		
		Size Co	olumn OK

For grouped data use one list e.g. D1 for the values and a second list for the frequencies e.g. D2.

For example	here is a	distribution	showing the	sizes of	mens'	shoes
FUI example,	liele is a	uistribution	showing the	SIZES UI	mens	shoes.

• •			•	<u> </u>											
Size	5	6	7	8	9	10	11	12	13						
Frequency	7	13	21	35	39	28	21	8	1						

	Stati	istics 1Var N	umeric Viev	v 10:36
	D1	D2	D3	D4
1	5	7		
2	6	13		
3	7	21		
4	8	35		
5	9	39		
6	10	28		
7	11	21		
8	12	8		
9	13	1		
10				
				I

Edit Ins Sort Size Make Stats



Statistics 1V	ar Symbolic View	10:37
<mark>√ H1:</mark> D1	D2	
Plot1: Histogram		v
H2:		
Plot2: Histogram		Ψ.
H3:		
Plot3: Histogram		*
H4:		
Choose 🗸		

Х	H1		
n	173		
Min	5		
Q1	8		
Med	9		
Q3	10		
Max	13		
ΣΧ	1511		
ΣX <sup>2</sup>	13733		
x	8.734104046		
sX	1.764917789		
σX	1.759809478		
173			
		Sizo Col	
		Size Col	

#### Two Variable Data

Start the 2 variable statistics applet: Apps/Statistics 2Var/RESET/OK/START

Enter some paired data, e.g. data comparing shoe size with handspan:

Shoe Size	6	9	11	9	10	12	9	7	5	12	9	7	8	9	4	8
Hand Span (cm)	18	23	25	22	26	27	24	23	19	28	21	22	20	24	19	20

Enter Shoe size as C1 and Handspan as C2

	Statistics 2Var Numeric View 10:40						
	C1	C2	C3	C4			
8	7	23					
9	5	19					
10	12	28					
11	9	21					
12	7	22					
13	8	20					
14	9	24					
15	4	19					
16	8	20					
17							

Press Symb and notice that the defualt is to plot C1 against C2 which is what we want.

Edit	Ins Sort	Size	Make	Stats
ann an straith straith stra	Statistics 2Va	r Symbolio	: View	10:41
√ S1:C1		C2		
Type1: Li	near			Ψ.
Fit1:	M*X+B			
S2:				
Type2: Li	near			*
Fit2:	M*X+B			
S3:				
Enter Ind Edit	ependent colu	imn Fit•	Show	Eval
		/		
		_/		
, I		/		
÷.	. :/			
	•∕+			
Į,	/			
X				
S1:1 X:6	_	Y:18		Menu
<u>X</u>	S1			
r	8.717070E-1			
sCOV	5.9375			
σCOV ΣΧΥ	5.56640625 3135			
16				
Stats•	ХҮ	Size	Column	OK

Press Plot/Menu/Zoom/Autoscale to see a scatter plot with the linear regression line, or on the HP Prime simply drag the points into view.

Pressing Fit on the Symb screen will turn the line on and off.

Press Num and Stats to see the summary statistics with the correlation coefficient and covarience.

Notably, the correlation coefficient r = 0.87 (t 2 d.p.) shows a reasonably strong relationship.

Press OK to leave the stats screen.

Press Symb again and notice that the details for the regression line are now shown.

The line is f(x) = 1.14x + 12.9 (to 3 s.f.) and our x values (i.e. the first list) showed shoe size. This suggests a model of the relationship between show size and handspan as:

H = 1.14S + 12.9 where H = handpsan and S = shoesize.

If we want to see if we would get a better fit with a different model, then we can change the model with the Type options. Navigate to the Type1 entry and choose Quadratic.

Press Num and Stats to calculate the statistics (many of which are undefined in this new model) then press OK and press Symb again to see the coefficients for the new model.

This gives us a new model of  $H = 0.94S^2 - 0.4S + 18.8$  where H =handpsan and S = shoesize.

Press Plot to see the new model.

Statistics 2Va	r Symbolic View	10:45
√ S1: <mark>C1</mark>	C2	
Type1: Linear		*
Fit1: 1.1427425822*	×X+12.92060946	27
S2:		
Type2: Linear		×
Fit2: M*X+B		
S3:		
Enter independent colu Edit	mn Fit• Show	Eval
Statistics 2Vai	r Symbolic View	10.04
Type1: Quadratic	C2	*
Fit1: A*X2+B*X+C		
52:		
Type2: Linear		v
Fit2: M*X+B		
53:		
Enter independent colu	mn	-
Edit √ C	Fit• Show	Eval
Statistics 2Va	r Symbolic View	10:55 4°
√ S1:C1	C2	
Type1: Quadratic		*
Fit1: .09405687637	2 99∗X −.4002578	34887*X
S2:		
Type2: Linear		v
Fit2: M*X+B		
ça. Enter independent colu	Imp	
Edit V C	Fit• Show	Eval
	· · · · / · · · · · · ·	
	/	
	/	
	/	

Y:18

Menu

S1:1 X:6

#### **Confidence Intervals**

We have some data about the shoe sizes of thirty female maths students. We want to use this data to answer the question, "What is the mean shoe size for UK female maths students?"

Here we are using a sample to 'infer' something about the whole population. This is called inferential statistics. The issue will be how confident we can be in the outcomes. We can say that with a given level of probability, the mean will lie within a certain interval. At a 95% level of confidence, we can calculate the range within which the mean will lie, with a probability of 0.95.

To do this, we need to know the mean and standard deviation and the number of data values of the sample.

First, we enter the sample data into the 1 variable statistics applet. Press Apps, choose Statistics 1Var then Reset/OK/Start then enter the data into list D1

The data is: {2, 7, 6, 4, 5, 3, 4, 5, 6, 7, 4, 3, 5, 4, 4, 3, 5, 6, 6, 8, 6, 7, 8, 4, 3, 9, 5, 7, 6, 4}

Now start the inference applet:

Press Apps, choose Inference then Reset/OK/Start

Chhose to change the Method to Confidence Interval and the Type to T-Int  $1\mu$ 

Press Num to get the statistics we entered earlier.

Press Import/ IMPRT. Check that we will import list D1 from the Statistics 1Var applet (which it will be expecting) and click OK

You can now see the sample mean  $\overline{x}$ , the sample standard deviation s and the number of sample data points n. You need to ch ange the confidence interval C to 0.95

	Statistics 1Var Numeric View 11:00						
	D1	D2	D3	D4			
22	7						
23	8						
24	4						
25	3						
26	9						
27	5						
28	7						
29	6						
30	4						
31							

dit Ins	Sort	Size	Make	Stats
Inf	ference Sy	mbolic	View	11:03
Method:	Confiden	ce inter	val	*
Type:	T–Int: 1 μ			*

In	nport Sample Statistics 11:	04
x:	5.2	
n:	30	
s:	1.73005879908	
App:	Statistics 1Var	Ŧ
Column:	D1	Ŧ

#### Choose app from which to import data

Choose a distribution statistic Choose

Choc	se		Cancel	ОК
	nference	e Numeric	View	11:08
х: 5.2				
s: 1.730058	879908			
n: 30				
C:: 0.95				
Edit		Impor	t	Calc

Press F6 for CALC to make the calculations.

This tells us that the probability is 0.95 that the shoe sizes of the whole population will fall between 4.55 and 5.85 (to 2 d.p.)



Press Plot to see this graphically.

As usual, experiment! See what impact changing the confidence interval has. Look at the other tests available. Research them. See what difference they make.

# 8. Calculus

It is important to remember that the HP39gII is a numeric calculator. The HP Prime has both numeric and symbolic modes. This chapter lets you explore how calculus works using the numeric mode.

#### Differentiation

Let's explore the differential of  $f(x) = x^2$ . The differential is the function which gives values of the slope of the original function at all values of x.

Open and reset the Function applet:

Apps/Function/Reset/OK/Start

Enter the function  $f(x) = x^2$ 

Press plot to see the graph.

Make sure that Trace is on (the little blob is showing next to Trace) and move the cursor to the furthest right hand/positive end of the visible graph. Now press Fcn and choose Slope.



Before you move the cursor, guess what the slope values will be as you move the cursor to the left hand/negative end of the visible graph. When you have formed a good idea, move the cursor and see.  $x = \frac{1}{2}$ 

Look to see the relationship between the value of the slope and the value of x. (It is easiest to see at key points like the one shown where the slope is 4 and you can see the value of x is 2).

You can see the relationship more clearly by splitting the screen. Press View/Views. Choose the second option (Split Screen: Plot table).

Press Fcn and choose slope, then choose another value of x in the table and find the slope.

Jot down the pairs of values if x and the corresponding slopes that you have found and decide what function shows the relationship between x and the slope. Go back to the Symb page and enter this function for F2(X). Press Num and check that the values for F2(X) are the same as the ones you jotted down.



Repeat this process with a range of functions. The sine function is very interesting. Make slight changes to the functions and see what difference it makes.

(e.g. change  $f(x) = x^2$  to  $f(x) = x^2 + 1$ )

Get a good feel for the differential functions generated by different types of functions.

## Integration

You can apply the same process to get a feel for the integral function. This time we are looking at the area under the graph.

Launch and reset the function applet.

Enter the function  $f(x) = x^2$  and Plot it.

Press Menu then Fcn and select Signed area. The cursor will already be at the origin, if not, move it there. Click OK. Now move the cursor to x = 1 and click OK.

Now repeat the process starting from x = 0 to x = 2 ... then from x = 0 to x = 3



Writing the values as fractions makes things clearer:

x	0	1	2	3
Area from 0 to <i>x</i>	0	$\frac{1}{3}$	83	$\frac{27}{3}$

Decide what you think the area function (the integral) is, Press Symb and enter it as F2(X). Press Num and check that this generates the same values.

Press Plot and TRACE over F2(X) to check.



Again, do this with different types of functions. Make small changes to existing functions. The aim is to get a feeling for how the area function varies. To be able to work out exactly what the function is from this small amount of data would be very difficult for most functions. But the key point is to experiment and get that all important feeling for how calculus works.

# Notes

For further information and educational support contact:

The Maths Zone

10 Staplehurst Road

London

SE13 5NB

T: (+44) (0)20 8318 6380

F: (+44) (0)20 8318 6610

info@themathszone.co.uk

www.themathszone.co.uk

www.hpgraphingcalc.org

