



HP35

MATH PAC

The program material contained herein is supplied without representation or warranty of any kind. Hewlett-Packard Company therefore assumes no responsibility and shall have no liability, consequential or otherwise, of any kind arising from the use of this program material or any part thereof.

Your Math Pac

The Math Pac is a representative collection of key sequence routines for solving problems with your HP-35 Pocket Calculator.

We suggest that you first read the introductory material explaining the Standard Key Sequence Format. Then find the routine you want, and use it. Numerical examples are provided to enable you to try out routines before using them. In most cases, the given answers are rounded. An understanding of the HP-35 Operating Manual is also required if, in addition, you wish to track the changes in the memory register and stack registers on a step-by-step basis.

To make for easy reference, the body of the book is arranged in alphabetic sequence of key topics—those you are likely to think of when you want a routine. Thus, Complex Number Operations are presented in the “C” section, Progressions are presented in the “P” section, etc. Statistics (in the “S” section) includes the Mean and Standard Deviation. Functions that are thus grouped and presented under higher level headings are also given cross reference entries in the alphabetic sequence. Thus, if you happen to search for Mean in the “M” section, you will find there a reference to the page in the “S” section where mean is presented under “Statistics”. The Contents section at the front is arranged logically (instead of by page order) to show all the routines available under each subject. The back cover contains an index.

Mr. Lee Skinner of Albuquerque, New Mexico (a prolific HP-35 user) contributed all but a few of the routines in this book. Two key sequence forms are included inside the back cover of this manual. You may wish to duplicate these forms and record your own key step programs.

2 Standard Key Sequence Format

Standard Key Sequence Format

Shown below is the key sequence routine for computing the roots of the Quadratic Equation:

$$AX^2 + BX + C = 0$$

Using A, B, and C, which you supply as data, the routine produces an intermediate result, D. If $D < 0$, one root is the complex conjugate of the other; the real and imaginary parts of one complex root develop on lines 7 and 8. Except for the opposite sign of the imaginary part, the other complex root is identical. If $D \geq 0$, the roots are real and develop on lines 5 and 6.

LINE	DATA	OPERATIONS	DISPLAY	REMARKS
1	A	\uparrow + STO		
2	B	RCL \div CHS \uparrow \uparrow		
3		x		
4	C	\uparrow + RCL \div -	D	If $D < 0$, go to 7
5		\sqrt{x} STO +	Root 1	
6		$x \leftrightarrow y$ RCL -	Root 2	Stop
7		CHS \sqrt{x} $x \leftrightarrow y$	Real part	
8		$x \leftrightarrow y$	Imag. part	

To execute the sequence, start with line 1 and read from left to right, making the appropriate keystrokes as you proceed. Interpret the respective columns as follows:

Data: Information to be supplied by you, the user. In the sample case, lines 1, 2, and 4 prompt the reader to enter coefficients A, B, and C.

Operations: The keys to be pressed after you enter any requested data item for the line. \uparrow is the symbol used to denote the $\text{ENTER } \uparrow$ key of the HP-35. All other key designations are identical to the HP-35 keys. Ignore any blank positions in the operations column.

Display: *Intermediate or final results which you should*, depending upon the case, *jot down*. In the sample case, D is developed so that the reader can decide which line (5 or 7) to execute next.

Remarks: Conditional and unconditional jumps to specified lines or other information for the reader. In the sample case, the reader is prompted to continue with line 7 (ignoring lines 5 and 6) if D is negative. If the condition fails, execution continues on the next line. In the sample case, the reader proceeds to line 5, if D is zero or positive.

Thus, lines are read in sequential order except where the remarks column directs otherwise (as in line 4 of the sample case). To assist the reader in distinguishing lines to be repeated, a sequence of lines making up an iterative process is outlined with a bold border. The following sequence for computing Chi-Square (lines 2, 3, and 4) illustrate this convention.

Formula:

$$\chi^2 = \sum_{i=1}^n \frac{(O_i - E_i)^2}{E_i}$$

where: O_i = observed frequency
 E_i = expected frequency

LINE	DATA	OPERATIONS	DISPLAY	REMARKS
1		CLX ↑ [] [] []		
2	O_i	↑ [] [] [] []		Perform lines 2–4 for $i = 1, \dots, n$
3	E_i	STO – ↑ × RCL		
4		÷ + [] [] []		

Contents

<i>Your Math Pac</i>	1
<i>Standard Key Sequence Format</i>	2
MACHINE RELATED OPERATIONS	
Register Operations	122
Clear stack (retain storage)	122
Delete x (lower stack)	122
Delete y (lower stack above X)	122
Reverse the stack	123
Fetch t (roll up stack)	123
Fetch t to Y	123
Fetch z to X	124
Copy x into Z and T	124
Copy y into Z	124
Copy y into Z and T	125
Copy y and x into Z and T	125
Copy y and x into T and Z	125
Swap x and s	126
Swap y and z	126
Swap z and t	126
Swap x and t	127
Swap x and z	127
Swap y and t	127
Rounding Numbers	131
Conversions	55
Centigrade to Fahrenheit	55
Fahrenheit to centigrade	55
Feet, inches, to centimeters	56
Centimeters to feet, inches	56
Exponentiation	70
Multiple power operations	70
e^x for large positive x	71
x^y for large x and/or y	72
Converging $u^u u^{u^u} \dots$	73
Diverging $u^u u^{u^u} \dots$	73
Limits of x^y	74
Double Precision Product	59

Simple Monadic Functions of x

Square of x	137
Antilog of x	16
Factorial of x	75

Simple Dyadic Functions of x and y

Powers	112
Roots	130
Logarithms	106
Permutations	108
Combinations	23
Remainder function: y modulo x	129

NUMBER THEORY AND ALGEBRA**Numbers**

Fibonacci numbers	79
Harmonic numbers	83
Bernoulli numbers	21
Euler numbers	69

Factoring Integers and Determining Primes 78**Highest Common Factor** 84**Least Common Multiple** 102**Fractions** 80

Improper fractions	80
Reduction to lowest terms	80
Addition and subtraction	81
Multiplication and division	82

Progressions 113

Arithmetic progressions	113
Geometric progressions	114
Harmonic progressions	114

Quadratic Equations 119**Simultaneous Linear Equations** 132**Complex Number Operations** 24

Complex add	24
Complex subtract	24
Complex multiply	24
Complex divide	26
Complex reciprocal	27

6 Contents

Complex absolute value	28
Complex square	28
Complex square root	28
Complex natural logarithm	29
Complex exponential	30
t^{a+ib}	30
Integral powers	31
Integral roots	32
Complex number to complex power	33
Complex root of complex number	34
Logarithm of complex number to complex base	34

Vector Algebra

Vector dot product	161
Vector cross product	160

Base Conversion	17
Decimal integer to another base	17
Integer, base b to decimal	19
Integer with exponent in base b to decimal	19
Fractional decimal number to another base	20
Fractional number in another base to decimal	20

GEOMETRY AND TRIGONOMETRY

Coordinate Conversion	57
Polar to rectangular	57
Rectangular to polar	57
Angle Conversions	11
Degrees, minutes, seconds to decimal degrees	11
Decimal degrees to degrees, minutes, seconds	11
Bearing to azimuth	12
Azimuth to bearing	12
Radians to degrees	13
Degrees to radians	14
Mils to degrees	14
Degrees to mils	14
Grads to degrees	15
Degrees to grads	15
Trigonometric Functions	154
Secondary arc sine	154
Secondary arc cosine	154
Secondary arc tangent	155
Cotangent	155

Cosecant	156
Secant	156
Versine	156
Coversine	157
Haversine	157
Arc cotangent	157
Arc cosecant	158
Arc secant	158
Hyperbolic and Inverse Hyperbolic Functions	85
Gudermannian function	85
Hyperbolic sine	85
Hyperbolic cosine	86
Hyperbolic tantent	86
Hyperbolic cotangent	86
Hyperbolic cosecant	87
Hyperbolic secant	87
Inverse Gudermannian	87
Inverse hyperbolic sine	88
Inverse hyperbolic tangent	88
Inverse hyperbolic secant	88
Inverse hyperbolic cosine	89
Inverse hyperbolic cotangent	89
Inverse hyperbolic cosecant	89
Complex Trigonometric and Complex Hyperbolic Functions	36
Complex sine	36
Complex cosine	36
Complex tangent	37
Complex cotangent	38
Complex cosecant	38
Complex secant	39
Complex arc sine	40
Complex arc cosine	41
Complex arc tangent	42
Complex arc cotangent	43
Complex arc cosecant	44
Complex arc secant	45
Complex hyperbolic sine	46
Complex hyperbolic cosine	46
Complex hyperbolic tangent	47
Complex hyperbolic cotangent	48
Complex hyperbolic cosecant	48
Complex hyperbolic secant	49

8 Contents

Complex inverse hyperbolic sine	50
Complex inverse hyperbolic cosine	50
Complex inverse hyperbolic tangent	51
Complex inverse hyperbolic cotangent	52
Complex inverse hyperbolic cosecant	53
Complex inverse hyperbolic secant	54
Triangles	147
Given a, b, C; find A, B, c	148
Given a, b, c; find A, B, C	148
Given a, A, C; find B, b, c	149
Given a, B, C; find A, b, c	150
Given B, b, c; find a, A, C	150
Given a, b, c; find area	151
Given a, B, C; find area	153
Given a, b, C; find area	152
STATISTICS	138
Means	
Arithmetic mean	138
Geometric mean	138
Harmonic mean	139
Average and Count Simultaneously	140
Sum Two Columns Simultaneously	141
Mean, Standard Deviation, and Sums	
Ungrouped data	141
Grouped data	142
Cross product sums	144
Normal or Gaussian Probability Function	82
T-Statistic (Paired)	158
Chi-Square	22
Random Number Generation	120
Linear Regression and Correlation Coefficient	144
NUMERICAL METHODS	
Interpolation (Linear)	100
Polynomial Evaluation	111

Infinite Power Series Summation	90
Factorial and Gamma Functions	75
Stirling's Approximation	75
Polynomial approximation	76
Plotting Curves	109
Equations, Iterative Solutions	63
Integration, Numerical	92
FINANCE	
Interest and Discount (Simple)	94
Interest	94
Loan payments	94
Discount	95
Interest (Compound)	96
Annuities	15
Sinking Fund and Amortization	134
Sinking fund method	134
Amortization method	135
Loan Repayments	102
Interest Rates	98
Percent	107
MISCELLANY	
Weekday from Date Computation	161
Speedometer/Odometer Adjustments	136
Resistance Equivalent	130
Register Use Optimization	128
APPENDIX	163
Index	back cover

Amortization

See page 135

Angle Conversions**Degrees, minutes, and seconds to decimal degrees***Example:*

$$46^{\circ} 17' 32.6'' = 46.29^{\circ}$$

LINE	DATA	OPERATIONS	DISPLAY	REMARKS
1	Seconds	[↑] [6] [0] [STO] [÷]		
2	Minutes	[+] [RCL] [÷] [] []		
3	Degrees	[+] [] [] [] []		

Decimal degrees to degrees, minutes and seconds*Note:* x = decimal degrees.*Example:*

$$23.32916667^{\circ} = 23^{\circ} 19' 45''$$

LINE	DATA	OPERATIONS	DISPLAY	REMARKS
1	X	[↑] [] [] [] []		Let D = integer part of X
2	D	[] [] [] [] []	Degrees	
3		[-] [6] [0] [STO] [x]	A	Let M = integer part of A
4	M	[] [] [] [] []	Minutes	
5		[-] [RCL] [x] [] []	B	Let S = nearest integer to B
6	S	[] [] [] [] []	Seconds	

12 Angle Conversions

Bearing to azimuth

Note:

X = bearing in decimal degrees

Example:

S 42.6° E = 137.4°

LINE	DATA	OPERATIONS	DISPLAY	REMARKS
1	X	↑		If S X° E, go to 2;
				If S X° W, go to 3;
				If N X° W, go to 4;
			Azimuth	If N X° E, stop
2		1 8 0 x ⁻² y -	Azimuth	Stop
3		1 8 0 +	Azimuth	Stop
4		CHS ↑ 3 6 0		
5		+	Azimuth	

Azimuth to bearing

Method:

If $0 < X < 90^\circ$, convert to $N^\circ \cdot' \cdot'' E$;
if $90^\circ < X < 180^\circ$, convert to $S^\circ \cdot' \cdot'' E$;
if $180^\circ < X < 270^\circ$, convert to $S^\circ \cdot' \cdot'' W$;
if $270^\circ < X < 360^\circ$, convert to $N^\circ \cdot' \cdot'' W$.

Note:

X = azimuth in decimal degrees

Example:

$226^\circ 23' = 226.3833333^\circ = S 46^\circ 23' W$

LINE	DATA	OPERATIONS	DISPLAY	REMARKS
1	X	↑	D	If $0 < X < 90$, go to 5;
		↑		If $90 < X < 180$, go to 2;
		↑		If $180 < X < 270$, go to 3;
		↑		If $270 < X < 360$, go to 4
2		1 8 0 $x \leftrightarrow y$ -	D	Go to 5
		↑		
3		1 8 0 -	D	Go to 5
		↑		
4		3 6 0 - CHS	D	
5	D ¹	↑	Degrees	Note: D ¹ is the integer part of D.
6		- 6 0 STO x	A	Let M = integer part of A
7	M	↑	Minutes	
8		- RCL x	B	Let S = nearest integer to B
9	S	↑	Seconds	

Radians to degrees

Examples:

- 1 radian = 57.29577951°
- $\frac{3}{4}\pi$ radians = 135°

LINE	DATA	OPERATIONS	DISPLAY	REMARKS
1		1 8 0 π \div		
2		↑ ↑ ↑		
3		CLX		
4	Radians	x	Degrees	Stop. For new case, go to 3.

14 Angle Conversions

Degrees to radians

Examples

1. $1^\circ = .0174532925$ radians
2. $266^\circ = 4.64$ radians

LINE	DATA	OPERATIONS	DISPLAY	REMARKS
1		π 1 8 0 \div		
2		\uparrow \uparrow \uparrow \square \square		
3		CLX \square \square \square		
4	Degrees	x \square \square \square	Radians	Stop. For new case, go to 3

Mils to degrees

Example:

$$1600 \text{ mils} = 90^\circ$$

LINE	DATA	OPERATIONS	DISPLAY	REMARKS
1	Mils	\uparrow 9 x 1 6		
2		0 \div \square \square	Degrees	

Degrees to mils

Example:

$$90^\circ = 1600 \text{ mils}$$

LINE	DATA	OPERATIONS	DISPLAY	REMARKS
1	Degrees	\uparrow 1 6 0 x		
2		9 \div \square \square	Mils	

Grads to degrees

Example:

$$300 \text{ grads} = 270^\circ$$

LINE	DATA	OPERATIONS	DISPLAY	REMARKS
1	Grads	<input type="button" value="↑"/> <input type="button" value="•"/> <input type="button" value="9"/> <input type="button" value="x"/> <input type="button" value=""/>	Degrees	

Degrees to grads

Example:

$$360^\circ = 400 \text{ grads}$$

LINE	DATA	OPERATIONS	DISPLAY	REMARKS
1	Degrees	<input type="button" value="↑"/> <input type="button" value="•"/> <input type="button" value="9"/> <input type="button" value="÷"/> <input type="button" value=""/>	Grads	

Angles of Triangle

See pages 147–151

Annuities

Present Value

The present value (PV) of an ordinary annuity is the total discounted value of a series of payments

$$PV = PMT \left[\frac{1 - (1 + i)^{-n}}{i} \right]$$

where PMT is the payment
 i is the periodic interest rate
 and n is the number of periods.

Example:

What is the present value of \$200 received at the end of each year for 10 years if the money earns 3% annually? (PMT = 200, i = .03, n = 10)

Answer: \$1706.04

16 Anti-Log₁₀ of X

LINE	DATA	OPERATIONS	DISPLAY	REMARKS
1	i	↑ STO 1 +		
2	n	x ^{-y} x ^y 1/x 1 x ^{-y}		
3		- RCL ÷		
4	PMT	x	PV	

Anti-Log₁₀ of X

Example:

$$\text{Antilog}_{10} 2 = 100.$$

LINE	DATA	OPERATIONS	DISPLAY	REMARKS
1	X	↑ 1 0 x ^y		

Arc Cosecant

See page 158

Arc Cosine, Secondary

See page 154

Arc Cotangent

See page 157

Arc Secant

See page 158

Arc Sine, Secondary

See page 154

Arc Tangent, Secondary

See page 155

Area of Triangle

See pages 151–153

Arithmetic Mean

See page 138

Arithmetic Progressions

See pages 113, 115-116

Average

See pages 138, 140–142

Azimuth to Bearing

See page 12

Base Conversion

Base conversion algorithms are given for positive values only. To convert a negative number, (1) change the sign, (2) convert the number, and (3) change the sign of result.

(A)_b means, “A is a number in base b.”

Decimal integer to integer in any base

$$(I)_{10} \rightarrow (J)_b$$

In the following key sequence, f + 1 is the number of digits in (J)_b. d_i (i = 1, f + 1) represents the ith digit in (J)_b, counting from left to right, i.e.

$$(J)_b = (d_1 d_2 \dots d_{f+1})_b$$

Notes: 1. If d_i = 0, line 7 can be replaced by the **[x]** key.

2. For large numbers, f is the exponent for base exponentiation.

$$(J)_b = (d_1 \cdot d_2 \dots d_{f+1})_b \times b^f$$

LINE	DATA	OPERATIONS	DISPLAY	REMARKS
1	b	↑ ↑		
2	I	STO In x ² y In ÷	D	Let f be the largest integer ≤ D
3	f	x ² y CLX +	f	
4		x ² y ↑ ↑ RCL R↓		
5		R↓ x ^v ÷	E ₁	Let d _i = integer part of E ₁
				(i = 1, ..., f)
6	d ₁	- x	E ₂	
7	d _i	- x	E _{i+1}	Perform line 7 for i = 2, ..., f
8				d _{f+1} = nearest integer to E _{f+1}

18 Base Conversion

Example 1:

Convert $(1206)_{10}$ to hexadecimal (base 16).

(The Hexadecimal digits are 0, 1, 2, 3, 4, 5, 6, 7, 8, 9, A, B, C, D, E, F).

Answer:

$$(1206)_{10} = (4B6)_{16} \quad (f = 2)$$

Example 2:

Convert 513 to octal (base 8).

Answer:

$$(513)_{10} = (1001)_8 \quad (f = 3)$$

Example 3:

Convert 6.023×10^{23} to octal.

Answer:

$$6.023 \times 10^{23} = (1.7743)_8 \times 8^{26}$$

Note:

If we consider 6.023×10^{23} to be a scientific measurement good only to 4 significant digits, it is meaningless for the octal representation to contain more than 5 significant digits. Therefore, we stop before the loop is completed.

LINE	DATA	OPERATIONS	DISPLAY	REMARKS
1	8	↑ ↑		
2	6.023	EEX		
3	23	STO In x \bar{x} y In \div	26.33160947	
4	26	x \bar{x} y CLX +	26	f = 26
5		x \bar{x} y ↑ ↑ RCL R↓		
6		R↓ x \bar{y} \div	1.992843519	d ₁ = 1
7	1	- x	7.942748152	d ₂ = 7
8	7	- x	7.541985216	d ₃ = 7
9	7	- x	4.335881728	d ₄ = 4
10	4	- x	2.687053824	d ₅ = 3 (rounded), Stop

Integer without exponent in base b to decimal integer

$$(d_1 d_2 \dots d_{n-1} d_n)_b \rightarrow (I)_{10}$$

Examples:

1. $(730020461)_8 = (123740465)_{10}$
2. $(7D0F)_{16} = (32015)_{10}$
(A = 10, B = 11, C = 12, D = 13, E = 14, F = 15 in the hexadecimal system)

Note: If $d_i = 0$, line 3 can be replaced by the **X** key.

LINE	DATA	OPERATIONS	DISPLAY	REMARKS
1	b	↑ ↑ ↑		
2	d_i	x		
3	d_i	+ x		Perform line 3 for $i = 2, \dots, n-1$
4	d_n	+	$(I)_{10}$	

Integer with exponent in base b to decimal integer

$$d_1 \cdot d_2 \cdots d_n \times b^{\text{Exp}} \rightarrow (I)_{10}$$

Examples:

1. $(3.0002)_8 \times 8^{11} = 2.577399803 \times 10^{10}$ ($b = 8$, $\text{Exp} = 11$, $n = 5$)
2. $(D2EE4)_{16} \times 16^{28} = (D.2EE4)_{16} \times 16^{32}$
 $= 4.485999088 \times 10^{39}$

Note: If $d_i = 0$, line 7 can be replaced by the **X** key.

LINE	DATA	OPERATIONS	DISPLAY	REMARKS
1	b	↑ ↑		
2	Exp	↑		
3	n	- 1 + x^y x^y		
4		STO		
5	b	↑ ↑ ↑		
6	d_i	X		
7	d_i	+ x		Perform line 7 for $i = 2, \dots, n-1$
8	d_n	+ RCL x	$(I)_{10}$	

20 Base Conversion

Fractional decimal number to base b

$$(x)_{10} \rightarrow (y)_b$$

Method:

In the following algorithm, c is the number of significant digits in x.

d_i ($i = 1, \dots, c$) represents the i^{th} digit in $(y)_b$, counting from left to right.

$$(y)_b = (d_1 \cdot d_2 \dots d_c)_b \times b^{-H}$$

Examples:

1. $(0.2937)_{10} = (2.263)_8 \times 8^{-1} = (.2263)_8 \quad (c = 4)$
2. $(3.688)_{10} \times 10^{-54} = (5.A6E)_{16} \times 16^{-45}$
 $(c = 4, \text{ in hexadecimal system } A = 10, E = 14)$

Note:

If $d_i = 0$, line 8 can be replaced by the $\boxed{\times}$ key.

LINE	DATA	OPERATIONS	DISPLAY	REMARKS
1	b	\uparrow \uparrow \square \square \square		
2	x	STO $1/x$ In $x \leftrightarrow y$ In		
3		\div \square \square \square	D	Let H be the smallest integer $\geq D$
4	H	$x \leftrightarrow y$ CLX + \square \square		
5		$x \leftrightarrow y$ \uparrow \uparrow RCL R↓		
6		R↓ x^y x \square \square	E ₁	$d_i = \text{integer part of } E_i$
		\square \square \square \square		$(i = 1, \dots, c-1)$
7	d ₁	- x \square \square	E ₂	
8	d _i	- x \square \square	E _{i+1}	Perform line 8 for $i = 2, \dots, c-1$
9		\square \square \square \square		$d_c = \text{nearest integer to } E_c$

Fractional number in base b to decimal number

$$(\cdot d_1 d_2 \dots d_n)_b \rightarrow (A)_{10}$$

or

$$d_1 \cdot d_2 \dots d_n \times b^{-\text{Exp}} \rightarrow (A)_{10}$$

Examples:

1. $(.0E728)_{16} = (.0564)_{10} \quad (\text{In hexadecimal system, } E = 14)$
2. $(7.200067)_8 \times 8^{-29} = 4.6853 \times 10^{-26} \quad (\text{Exp} = 29)$

Note: Exponents are in base 10.

Note: If $d_i = 0$, line 3 can be replaced by the \times key.

LINE	DATA	OPERATIONS	DISPLAY	REMARKS
1	b	$1/x$ \uparrow \uparrow \uparrow		
2	d_n	x		
3	d_i	+ x		Perform line 3 for $i = n-1, \dots, 1$
4			(A) ₁₀	If there is no exponent, stop; Otherwise go to 5
5	Exp	\uparrow 1 -		
6	b	x^y $1/x$ x	(A) ₁₀	

Bearing to Azimuth

See page 12

Bernoulli Numbers

The Bernoulli numbers $B_1, B_2, B_3 \dots$ are defined by

$$B_n = \frac{(2n)!}{\pi^{2n} 2^{2n-1}} \left[1 + \frac{1}{2^{2n}} + \frac{1}{3^{2n}} + \frac{1}{4^{2n}} + \dots \right].$$

Specifically,

$$\frac{1}{6}, \frac{1}{30}, \frac{1}{42}, \frac{1}{30}, \frac{5}{66}, \frac{691}{2730}, \frac{7}{6}, \dots$$

Example:

The 8th Bernoulli number = 7.092

(i takes the values 1, 2, ... and
j takes the values 15, 14, ..., 2 in the example.)

LINE	DATA	OPERATIONS	DISPLAY	REMARKS
1	n	\uparrow 2 x \uparrow \uparrow		
2		\uparrow 1		
3		STO		Perform lines 3–5 for $i = 1, 2, \dots$
4	i	\uparrow 2 x 1 +		until A_i does not change
5		x^y $1/x$ RCL +	A_i	
6		R↓	D	If $D \leq 2$, go to 8
7	j	x		Perform line 7 for $j = D-1, D-2, \dots, 2$
8		2 x RCL x R↓		
9		R↓ 2 x^y 1 -		
10		x^{2-y} π x^y x \div		

22 Chi-Square Evaluation

Centigrade to Fahrenheit

See page 55

Centimeters to Feet, Inches

See page 56

Chi-Square Evaluation

Formula:

$$\chi^2 = \sum_{i=1}^n \frac{(O_i - E_i)^2}{E_i}$$

where O_i = observed frequency

E_i = expected frequency

Example:

O_i	8	50	47	56	5	14
E_i	9.6	46.75	51.85	54.4	8.25	9.15

Answer:

$$\chi^2 = 4.844$$

LINE	DATA	OPERATIONS	DISPLAY	REMARKS
1		CLX ↑ [] [] []		
2	O_i	↑ [] [] [] []		Perform lines 2–4 for $i = 1, \dots, n$
3	E_i	STO – ↑ x RCL		
4		÷ + [] [] []		

Combinations

y objects taken x at a time (binomial coefficient)

Formula:

$$\binom{y}{x} = {}_y C_x = C_x^y = C(y, x) = \frac{y!}{x!(y-x)!} = \frac{y \cdot (y-1) \cdot \dots \cdot (y-x+1)}{1 \cdot 2 \cdot \dots \cdot x}$$

Notes:

1. This algorithm is for $y \geq x \geq 2$.
2. ${}_y C_0 = {}_y C_y = 1$; ${}_y C_1 = {}_y C_{y-1} = y$
3. ${}_y C_x = {}_y C_{y-x}$; so if $y - x < x$, compute ${}_y C_{y-x}$ instead of ${}_y C_x$.

Example:

$${}_7 C_5 = 21. \text{ (Compute } {}_7 C_2 \text{ to get the answer)}$$

LINE	DATA	OPERATIONS	DISPLAY	REMARKS
1	y	\uparrow <input type="text"/> <input type="text"/> <input type="text"/> <input type="text"/> <input type="text"/>		
2	i	x <input type="text"/> <input type="text"/> <input type="text"/> <input type="text"/> <input type="text"/> <input type="text"/> <input type="text"/> <input type="text"/> <input type="text"/> <input type="text"/>		Perform line 2 for $i = y - x + 1$, ..., $y - 1$
3	x	<input type="text"/> <input type="text"/> <input type="text"/> <input type="text"/> <input type="text"/> <input type="text"/> <input type="text"/> <input type="text"/> <input type="text"/> <input type="text"/>		If $x = 2$, go to 6.
4		\uparrow <input type="text"/> <input type="text"/> <input type="text"/> <input type="text"/> <input type="text"/>		
5	j	x <input type="text"/> <input type="text"/> <input type="text"/> <input type="text"/> <input type="text"/> <input type="text"/> <input type="text"/> <input type="text"/> <input type="text"/> <input type="text"/>		Perform line 5 for $j = x - 1$, $x - 2, \dots, 2$
6		\div <input type="text"/> <input type="text"/> <input type="text"/> <input type="text"/> <input type="text"/>		

Complex Number Operations

Complex add

Formula:

$$(a_1 + ib_1) + (a_2 + ib_2) = (a_1 + a_2) + i(b_1 + b_2)$$

$$= u + iv$$

Example:

$$(3 + 4i) + (7.4 - 5.6i) = 10.4 - 1.6i$$

LINE	DATA	OPERATIONS	DISPLAY	REMARKS
1	a_1	<input type="button" value="↑"/> <input type="button" value=" "/> <input type="button" value=" "/> <input type="button" value=" "/> <input type="button" value=" "/>		
2	a_2	<input type="button" value="+"/> <input type="button" value=" "/> <input type="button" value=" "/> <input type="button" value=" "/> <input type="button" value=" "/>	u	
3	b_1	<input type="button" value="↑"/> <input type="button" value=" "/> <input type="button" value=" "/> <input type="button" value=" "/> <input type="button" value=" "/>		
4	b_2	<input type="button" value="+"/> <input type="button" value=" "/> <input type="button" value=" "/> <input type="button" value=" "/> <input type="button" value=" "/>	v	

Complex subtract

Formula:

$$(a_1 + ib_1) - (a_2 + ib_2) = (a_1 - a_2) + i(b_1 - b_2)$$

$$= u + iv$$

Example:

$$(3 + 4i) - (7.4 - 5.6i) = -4.4 + 9.6i$$

LINE	DATA	OPERATIONS	DISPLAY	REMARKS
1	a_1	<input type="button" value="↑"/> <input type="button" value=" "/> <input type="button" value=" "/> <input type="button" value=" "/> <input type="button" value=" "/>		
2	a_2	<input type="button" value="-"/> <input type="button" value=" "/> <input type="button" value=" "/> <input type="button" value=" "/> <input type="button" value=" "/>	u	
3	b_1	<input type="button" value="↑"/> <input type="button" value=" "/> <input type="button" value=" "/> <input type="button" value=" "/> <input type="button" value=" "/>		
4	b_2	<input type="button" value="-"/> <input type="button" value=" "/> <input type="button" value=" "/> <input type="button" value=" "/> <input type="button" value=" "/>	v	

Complex multiply

(Use for short data input)

Formula:

$$(a_1 + ib_1)(a_2 + ib_2) = (a_1a_2 - b_1b_2) + i(a_1b_2 + a_2b_1)$$

$$= u + iv$$

Example:

$$(3 + 4i)(7 - 2i) = 29 + 22i$$

LINE	DATA	OPERATIONS	DISPLAY	REMARKS
1	b_1	\uparrow 		
2	a_1	\uparrow \uparrow 		
3	a_2	x 		
4	b_2	$R\downarrow$ $R\downarrow$ $R\downarrow$ STO x		
5		$-$ 	u	
6	a_2	RCL x $R\downarrow$ $R\downarrow$ $R\downarrow$		
7	b_2	x $+$ 	v	

Complex multiply, alternate method

(Data are entered only once, use for long data input)

Method:

If a_1, a_2, b_1, b_2 are all non-zero, and if $\theta = \theta_1 + \theta_2 \neq 90^\circ$ or -90° , then

$$(a_1 + ib_1)(a_2 + ib_2) = \frac{b_1 \cdot b_2}{\sin \theta_1 \cdot \sin \theta_2} (\cos \theta + i \sin \theta) = u + iv.$$

Note: $a_1 + ib_1 = r_1 e^{i\theta_1}$,

$$a_2 + ib_2 = r_2 e^{i\theta_2}$$

Example:

$$(3 + 4i)(5 - 12i) = 63 - 16i$$

LINE	DATA	OPERATIONS	DISPLAY	REMARKS
1	a_1	\uparrow 		
2	b_1	STO $x \div y$ \div arc tan		
3	a_2	\uparrow 		
4	b_2	\uparrow $R\downarrow$ $x \div y$ \div arc		
5		tan RCL $R\downarrow$ $R\downarrow$ $R\downarrow$		
6		x STO $R\downarrow$ \uparrow $R\downarrow$		
7		$+$ 	D	If D = 90 or -90, stop. Use other method.
8		RCL $R\downarrow$ STO $R\downarrow$ sin		
9		$x \div y$ sin x \div RCL		
10		cos x \uparrow \uparrow RCL		
11		tan x $x \div y$ 	u	
12		$x \div y$ 	v	

26 Complex Number Operations

Complex divide

(Use for short data input)

Formula:

$$(a_1 + ib_1) \div (a_2 + ib_2) = \frac{(a_1 a_2 + b_1 b_2) + i(a_2 b_1 - a_1 b_2)}{a_2^2 + b_2^2}$$

$$= u + iv$$

Example:

$$\frac{3 + 4i}{7 - 2i} = .245 + .64i$$

LINE	DATA	OPERATIONS	DISPLAY	REMARKS
1	b_1	↑		
2	a_1	↑ ↑		
3	a_2	x		
4	b_2	R↓ R↓ R↓ STO x		
5		+		
6	a_2	RCL x R↓ R↓ R↓		
7	b_2	x - STO		
8	a_2	↑ x		
9	b_2	↑ x + RCL x \neq y		
10		STO ÷ x \neq y RCL ÷	u	
11		x \neq y	v	

Complex divide, alternate method

(Data are entered only once; use for long data input)

Formula:

If a_1, a_2, b_1, b_2 are all non-zero, and if $\theta = \theta_1 - \theta_2 \neq 90^\circ$ or -90° , then

$$(a_1 + ib_1) \div (a_2 + ib_2) = \left(\frac{b_1}{b_2} \right) \left/ \left(\frac{\sin \theta_1}{\sin \theta_2} \right) \right. (\cos \theta + i \sin \theta) = u + iv$$

Note: $a_1 + ib_1 = r_1 e^{i\theta_1}$,

$a_2 + ib_2 = r_2 e^{i\theta_2}$.

Example:

$$(63 - 16i) \div (5 - 12i) = 3 + 4i$$

LINE	DATA	OPERATIONS	DISPLAY	REMARKS
1	a ₁	↑ [] [] [] []		
2	b ₁	STO x ² y ÷ arc tan		
3	a ₂	↑ [] [] [] []		
4	b ₂	↑ R↓ x ² y ÷ arc		
5		tan RCL R↓ R↓ R↓		
6		÷ STO R↓ ↑ R↓		
7		- [] [] [] []	D	If D = 90 or -90, stop.
		[] [] [] [] []		Use other method.
8		RCL R↓ STO R↓ sin		
9		x ² y sin ÷ ÷ RCL		
10		cos x ↑ ↑ RCL		
11		tan x x ² y [] []	u	
12		x ² y [] [] [] []	v	

Complex reciprocal

Formula:

$$\frac{1}{a + ib} = \frac{a - ib}{a^2 + b^2} = u + iv$$

Example:

$$\frac{1}{2 + 3i} = .15 - .23i$$

LINE	DATA	OPERATIONS	DISPLAY	REMARKS
1	b	↑ [] [] [] []		
2	a	↑ ↑ x STO CLX		
3		+ R↓ x RCL +		
4		STO ÷ CHS x ² y RCL		
5		÷ [] [] [] []	u	
6		x ² y [] [] [] []	v	

28 Complex Number Operations

Complex absolute value

Formula:

$$|a + bi| = \sqrt{a^2 + b^2}$$

Example:

$$|3 + 4i| = 5$$

LINE	DATA	OPERATIONS	DISPLAY	REMARKS
1	a	<input type="button" value="↑"/> <input type="button" value="x"/> <input type="button" value=" "/> <input type="button" value=" "/> <input type="button" value=" "/>		
2	b	<input type="button" value="↑"/> <input type="button" value="x"/> <input type="button" value="+"/> <input type="button" value="√x"/> <input type="button" value=" "/>		

Complex square

Formula:

$$(a + bi)^2 = (a^2 - b^2) + i(2ab) = u + iv$$

Example:

$$(7 - 2i)^2 = 45 - 28i$$

LINE	DATA	OPERATIONS	DISPLAY	REMARKS
1	a	<input type="button" value="↑"/> <input type="button" value="↑"/> <input type="button" value="x"/> <input type="button" value=" "/> <input type="button" value=" "/>		
2	b	<input type="button" value="STO"/> <input type="button" value="↑"/> <input type="button" value="x"/> <input type="button" value="-"/> <input type="button" value=" "/>	u	
3		<input type="button" value="x<sup>2</sup>y"/> <input type="button" value="RCL"/> <input type="button" value="x"/> <input type="button" value="2"/> <input type="button" value="x"/>	v	

Complex square root

Formula:

$$\sqrt{a + bi} = \pm \left[\sqrt{\frac{a + \sqrt{a^2 + b^2}}{2}} + i \frac{b}{\sqrt{\frac{a + \sqrt{a^2 + b^2}}{2}}} \right] = \pm(u + iv)$$

Example:

$$\sqrt{7 + 6i} = \pm(2.85 + 1.05i)$$

LINE	DATA	OPERATIONS	DISPLAY	REMARKS
1	b	\uparrow $\boxed{}$ $\boxed{}$ $\boxed{}$ $\boxed{}$		
2	a	STO $x \leftrightarrow y$ $\boxed{}$ $\boxed{}$ $\boxed{}$		If $b = 0$ and $a < 0$, go to 7; if $b = 0$ and $a \geq 0$, go to 9.
		$\boxed{}$ $\boxed{}$ $\boxed{}$ $\boxed{}$ $\boxed{}$		
3		\uparrow \uparrow x RCL \uparrow		
4		x $+$ \sqrt{x} RCL $+$		
5		2 \div \sqrt{x} STO $\boxed{}$	u	
6		2 RCL x \div $\boxed{}$	v	Stop
		$\boxed{}$ $\boxed{}$ $\boxed{}$ $\boxed{}$ $\boxed{}$		
7		RCL CHS \sqrt{x} $x \leftrightarrow y$ $\boxed{}$	u	
8		$x \leftrightarrow y$ $\boxed{}$ $\boxed{}$ $\boxed{}$ $\boxed{}$	v	Stop
		$\boxed{}$ $\boxed{}$ $\boxed{}$ $\boxed{}$ $\boxed{}$		
9		RCL \sqrt{x} $\boxed{}$ $\boxed{}$ $\boxed{}$	u	
10		$x \leftrightarrow y$ $\boxed{}$ $\boxed{}$ $\boxed{}$ $\boxed{}$	v	

Complex natural logarithm (base e)

Formula:

$$\ln(a + ib) = \ln(\sqrt{a^2 + b^2}) + i \left(\arctan \frac{b}{a} \right) \frac{\pi}{180}$$

$$= u + iv$$

Example:

$$\ln i = 1.57i$$

LINE	DATA	OPERATIONS	DISPLAY	REMARKS
1	b	\uparrow $\boxed{}$ $\boxed{}$ $\boxed{}$ $\boxed{}$		
2	a	$\boxed{}$ $\boxed{}$ $\boxed{}$ $\boxed{}$ $\boxed{}$		If $a \neq 0$, go to 4
		$\boxed{}$ $\boxed{}$ $\boxed{}$ $\boxed{}$ $\boxed{}$		
3		CLX EEX CHS 9 9		
4		STO $x \leftrightarrow y$ \uparrow \uparrow x		
5		RCL \uparrow x $+$ \sqrt{x}		
6		in $\boxed{}$ $\boxed{}$ $\boxed{}$ $\boxed{}$	u	
7		$x \leftrightarrow y$ RCL \div arc tan $\boxed{}$		If $a \geq 0$, go to 9
		$\boxed{}$ $\boxed{}$ $\boxed{}$ $\boxed{}$ $\boxed{}$		
8		1 8 0 + $\boxed{}$		
9		π x 1 8 0		
10		\div $\boxed{}$ $\boxed{}$ $\boxed{}$ $\boxed{}$	v	

30 Complex Number Operations

Complex exponential

Formula:

$$e^{(a+ib)} = e^a (\cos \theta + i \sin \theta) = u + iv$$

where

$$\theta = \frac{180b}{\pi}$$

Example:

$$e^{1.570796327i} = i$$

LINE	DATA	OPERATIONS	DISPLAY	REMARKS
1	b	\uparrow $\boxed{}$ $\boxed{}$ $\boxed{}$ $\boxed{}$		
2	a	e^x $\boxed{}$ STO $x \leftrightarrow y$ 1 8		
3		0 x π \div \uparrow		
4		\cos RCL x $\boxed{}$ $\boxed{}$	u	
5		$x \leftrightarrow y$ \sin RCL x $\boxed{}$	v	

Complex exponential (t^{a+ib})

Formula:

$$t^{a+ib} = e^{(a+ib)\ln t} = u + iv$$

Restriction:

$$(t > 0)$$

Example:

$$2^{3+4i} = -7.46 + 2.89i$$

LINE	DATA	OPERATIONS	DISPLAY	REMARKS
1	b	\uparrow $\boxed{}$ $\boxed{}$ $\boxed{}$ $\boxed{}$		
2	a	$x \leftrightarrow y$ $\boxed{}$ $\boxed{}$ $\boxed{}$		
3	t	\ln $\boxed{}$ STO x $x \leftrightarrow y$ RCL		
4		x e^x $\boxed{}$ STO $x \leftrightarrow y$ 1		
5		8 0 x π \div		
6		\uparrow \cos RCL x $\boxed{}$	u	
7		$x \leftrightarrow y$ \sin RCL x $\boxed{}$	v	

Complex number to integral power

Formula:

$$(a + ib)^n = r^n (\cos n\theta + i \sin n\theta) = u + iv$$

where: $r = \sqrt{a^2 + b^2}$,

$$\theta = \text{arc tan } \frac{b}{a}, \text{ and}$$

n is a positive integer.

Example:

$$(3 + 4.5i)^5 = 926.44 - 4533.47i$$

LINE	DATA	OPERATIONS	DISPLAY	REMARKS
1	b	↑		
2	a			If $a \neq 0$, go to 4
3		CLX EEX CHS 9 9		
4		STO x ² y ↑ ↑ x		
5		RCL ↑ x + √x		
6		x ² y RCL ÷ arc tan		If $a \geq 0$, go to 8
7		1 8 0 +		
8		STO		
9	n	↑ ↑ R↓ R↓ R↓		
10		x ^y RCL x ² y STO CLX		
11		+ x ↑ cos RCL		
12		x	u	
13		x ² y sin RCL x	v	

32 Complex Number Operations

Integral roots of complex number

Formula:

$$(a + ib)^{\frac{1}{n}} = r^{\frac{1}{n}} \left(\cos \frac{\theta + 360k}{n} + i \sin \frac{\theta + 360k}{n} \right)$$

$$= u_k + iv_k$$

where: n is a positive integer, and

$$k = 0, 1, \dots, n - 1.$$

Example:

$5 + 3i$ has three cube roots:

$$u_0 + iv_0 = 1.77 + .32i$$

$$u_1 + iv_1 = -1.16 + 1.37i$$

$$u_2 + iv_2 = -.61 - 1.69i$$

LINE	DATA	OPERATIONS	DISPLAY	REMARKS
1	b	\uparrow $\boxed{}$ $\boxed{}$ $\boxed{}$ $\boxed{}$		
2	a	$\boxed{}$ $\boxed{}$ $\boxed{}$ $\boxed{}$ $\boxed{}$		If $a \neq 0$, go to 4
		$\boxed{}$ $\boxed{}$ $\boxed{}$ $\boxed{}$ $\boxed{}$		
3		CLX EEX CHS 9 9		
4		STO $x \leftrightarrow y$ \uparrow \uparrow x		
5		RCL \uparrow x + \sqrt{x}		
6		$x \leftrightarrow y$ RCL \div arc tan		If $a \geq 0$, go to 8
		$\boxed{}$ $\boxed{}$ $\boxed{}$ $\boxed{}$ $\boxed{}$		
7		1 8 0 +		
8		$x \leftrightarrow y$		
9	n	\uparrow $1/x$ R↓ R↓ $x \leftrightarrow y$		
10		R↓ x STO CLX +		
11		\div \uparrow \uparrow cos RCL		
12		x	u ₀	
13		$x \leftrightarrow y$ sin RCL x	v ₀	
14		R↓ R↓		Perform lines 14–18 for k = 1,
		$\boxed{}$ $\boxed{}$ $\boxed{}$ $\boxed{}$ $\boxed{}$	2, ..., n-1	
15		3 6 0 \uparrow		
16	n	\div + \uparrow \uparrow cos		
17		RCL x	u _k	
18		$x \leftrightarrow y$ sin RCL x	v _k	

Complex number to a complex power

Formula:

If $a_1 + ib_1 \neq 0$,

$$(a_1 + ib_1)^{(a_2 + ib_2)} = e^{(a_2 + ib_2)\ln(a_1 + ib_1)} = u + iv$$

Example:

$$(1 + i)^{(2-i)} = 1.49 + 4.13i$$

LINE	DATA	OPERATIONS	DISPLAY	REMARKS
1	b_1	\uparrow [] [] [] []		
2	a_1	[] [] [] [] []		If $a_1 \neq 0$, go to 4
		[] [] [] [] []		
3		CLX EEX CHS 9 9		
4		STO $x \leftrightarrow y$ \uparrow \uparrow x		
5		RCL \uparrow x + \sqrt{x}		
6		In $x \leftrightarrow y$ RCL \div arc		
7		tan [] [] [] []		If $a_1 \geq 0$, go to 9
		[] [] [] [] []		
8		1 8 0 + []		
9		π x 1 8 0		
10		\div $x \leftrightarrow y$ \uparrow \uparrow []		
11	a_2	x [] [] [] []		
12	b_2	R↓ R↓ R↓ STO x		
13		- [] [] [] []		
14	a_2	RCL x R↓ R↓ R↓		
15	b_2	x + $x \leftrightarrow y$ e ^x STO		
16		$x \leftrightarrow y$ 1 8 0 x		
17		π \div \uparrow cos RCL		
18		x [] [] [] []	u	
19		$x \leftrightarrow y$ sin RCL x []	v	

34 Complex Number Operations

Complex root of a complex number

Formula:

If $a_1 + ib_1 \neq 0$

$$(a_1 + ib_1)^{\frac{1}{a_2 + ib_2}} = e^{[\ln(a_1 + ib_1)] / (a_2 + ib_2)}$$

$$= u + iv$$

Example:

Find the $(2 - i)^{\text{th}}$ root of $1.49 + 4.126i$.

Answer: $1 + i$

LINE	DATA	OPERATIONS	DISPLAY	REMARKS
1	b_1	\uparrow $\boxed{}$ $\boxed{}$ $\boxed{}$ $\boxed{}$		
2	a_1	$\boxed{}$ $\boxed{}$ $\boxed{}$ $\boxed{}$ $\boxed{}$		If $a_1 \neq 0$, go to 4
3		$\boxed{}$ $\boxed{}$ $\boxed{}$ $\boxed{}$ $\boxed{}$		
4		$\boxed{}$ $\boxed{}$ $\boxed{}$ $\boxed{}$ $\boxed{}$		
5		$\boxed{}$ $\boxed{}$ $\boxed{}$ $\boxed{}$ $\boxed{}$		
6		$\boxed{}$ $\boxed{}$ $\boxed{}$ $\boxed{}$ $\boxed{}$		
7		$\boxed{}$ $\boxed{}$ $\boxed{}$ $\boxed{}$ $\boxed{}$		If $a_1 \geq 0$, go to 9
8		$\boxed{}$ $\boxed{}$ $\boxed{}$ $\boxed{}$ $\boxed{}$		
9		$\boxed{}$ $\boxed{}$ $\boxed{}$ $\boxed{}$ $\boxed{}$		
10		$\boxed{}$ $\boxed{}$ $\boxed{}$ $\boxed{}$ $\boxed{}$		
11	a_2	x $\boxed{}$ $\boxed{}$ $\boxed{}$ $\boxed{}$		
12	b_2	$R\downarrow$ $R\downarrow$ $R\downarrow$ STO x		
13		$+ \quad \boxed{}$ $\boxed{}$ $\boxed{}$ $\boxed{}$		
14	a_2	RCL x $R\downarrow$ $R\downarrow$ $R\downarrow$		
15	b_2	x $-$ STO $\boxed{}$ $\boxed{}$		
16	a_2	\uparrow x $\boxed{}$ $\boxed{}$ $\boxed{}$		
17	b_2	\uparrow x $+$ RCL xz^y		
18		STO \div xz^y RCL \div		
19		e^x STO xz^y 1 8		
20		0 x π \div \uparrow		
21		\cos RCL x $\boxed{}$ $\boxed{}$	u	
22		xz^y \sin RCL x $\boxed{}$	v	

Logarithm of a complex number to a complex base

Formula:

$$\log_{(a_1 + ib_1)} (a_2 + ib_2) = \frac{\ln (a_2 + ib_2)}{\ln (a_1 + ib_1)} = u + iv$$

$$a_1 + ib_1 \neq 0$$

Example: $\log_{(1+i)} (1.49 + 4.126i) = 2 - i$

$$(a_3 = .34657359, b_3 = .7853981633)$$

LINE	DATA	OPERATIONS	DISPLAY	REMARKS
1	b_1	\uparrow [] [] [] []		
2	a_1	[] [] [] [] []		If $a_1 \neq 0$, go to 4
		[] [] [] [] []		
3		CLX EEX CHS 9 9		
4		STO x \leftrightarrow y \uparrow \uparrow x		
5		RCL \uparrow x + \sqrt{x}		
6		In [] [] [] []	a_3	
7		x \leftrightarrow y RCL \div arc tan		If $a_1 \geq 0$, go to 9
		[] [] [] []		
8		1 8 0 + []		
9		π x [] []		
10		1 8 0 \div []	b_3	
11		CLR [] [] []		
12	b_2	\uparrow [] [] []		
13	a_2	[] [] [] []		If $a_2 \neq 0$, go to 15
		[] [] [] []		
14		CLX EEX CHS 9 9		
15		STO x \leftrightarrow y \uparrow \uparrow x		
16		RCL \uparrow x + \sqrt{x}		
17		In x \leftrightarrow y RCL \div arc		
18		tan [] [] [] []		If $a_2 \geq 0$, go to 20
		[] [] [] []		
19		1 8 0 + []		
20		π x 1 8 0		
21		\div x \leftrightarrow y \uparrow \uparrow []		
22	a_3	x [] [] []		
23	b_3	R \downarrow R \downarrow R \downarrow STO x		
24		+ [] [] []		
25	a_3	RCL x R \downarrow R \downarrow R \downarrow		
26	b_3	x - STO []		
27	a_3	\uparrow x [] []		
28	b_3	\uparrow x + RCL x \leftrightarrow y		
29		STO \div x \leftrightarrow y RCL \div []	u	
30		x \leftrightarrow y [] [] []	v	

Complex Trigonometric and Hyperbolic Functions

In this section all angles in the equations are in radians.

Complex sine

Formula:

$$\begin{aligned}\sin(a + ib) &= \sin a \cosh b + i \cos a \sinh b \\ &= u + iv\end{aligned}$$

Example:

$$\sin(2 + 3i) = 9.154 - 4.1689i$$

LINE	DATA	OPERATIONS	DISPLAY	REMARKS
1	b	\uparrow 		
2	a	\uparrow 1 8 0 x		
3		π \div STO sin $x \leftrightarrow y$		
4		\uparrow \uparrow R↓ R↓ e^x		
5		\uparrow $1/x$ + 2 \div		
6		x RCL $x \leftrightarrow y$ STO	u	
7		CLX + cos $x \leftrightarrow y$ e^x		
8		\uparrow $1/x$ - 2 \div		
9		x 	v	

Complex cosine

Formula:

$$\begin{aligned}\cos(a + ib) &= \cos a \cosh b - i \sin a \sinh b \\ &= u + iv\end{aligned}$$

Example:

$$\cos(2 + ei) = -4.189 - 9.109i$$

LINE	DATA	OPERATIONS	DISPLAY	REMARKS
1	b	\uparrow \square \square \square \square		
2	a	\uparrow 1 8 0 x		
3		π \div STO cos $x \Rightarrow y$		
4		\uparrow \uparrow R↓ R↓ e^x		
5		\uparrow $1/x$ + 2 \div		
6		x RCL $x \Rightarrow y$ STO \square	u	
7		CLX + sin $x \Rightarrow y$ e^x		
8		\uparrow $1/x$ - 2 \div		
9		x CHS \square \square \square	v	

Complex tangent

Formula:

$$\tan(a + ib) = \frac{\sin 2a + i \sinh 2b}{\cos 2a + \cosh 2b}$$

$$= u + iv$$

Example:

$$\tan(2 + 3i) = -.00376 + 1.003i$$

LINE	DATA	OPERATIONS	DISPLAY	REMARKS
1	b	\uparrow \square \square \square \square		
2	a	\uparrow 2 x STO \uparrow		
3		1 8 0 x π		
4		\div cos $x \Rightarrow y$ 2 x		
5		\uparrow \uparrow R↓ R↓ e^x		
6		\uparrow $1/x$ + 2 \div		
7		+ RCL $x \Rightarrow y$ STO CLX		
8		+ $x \Rightarrow y$ e^x \uparrow $1/x$		
9		- 2 \div RCL \div		
10		$x \Rightarrow y$ 1 8 0 x		
11		π \div sin RCL \div	u	
12		$x \Rightarrow y$ \square \square \square \square	v	

38 Complex Trigonometric and Hyperbolic Functions

Complex cotangent

Formula:

$$\cot(a + ib) = \frac{\sin 2a - i \sinh 2b}{\cosh 2a - \cos 2b}$$

$$= u + iv$$

Example:

$$\cot(2 + 3i) = -.0037 - .9968i$$

LINE	DATA	OPERATIONS	DISPLAY	REMARKS
1	b	↑		
2	a	↑ 2 x STO ↑		
3		1 8 0 x π		
4		÷ cos x ^y 2 x		
5		↑ ↑ R↓ R↓ e ^x		
6		↑ 1/x + 2 ÷		
7		x ^y - RCL x ^y STO		
8		CLX + x ^y e ^x ↑		
9		1/x - 2 ÷ RCL		
10		÷ CHS x ^y 1 8		
11		0 x π ÷ sin		
12		RCL ÷	u	
13		x ^y	v	

Complex cosecant

Formula:

$$\csc(a + ib) = \frac{1}{\sin(a + ib)}$$

$$= u + iv$$

Example:

$$\csc(2 + 3i) = .09 + .0412i$$

LINE	DATA	OPERATIONS					DISPLAY	REMARKS
1	b	↑						
2	a	↑	1	8	0	x		
3		π	÷	STO	sin	x ^z y		
4		↑	↑	R↓	R↓	e ^x		
5		↑	1/x	+	2	÷		
6		x	RCL	x ^z y	STO	CLX		
7		+	cos	x ^z y	e ^x	↑		
8		1/x	-	2	÷	x		
9		RCL	↑	x	x ^z v	↑		
10		↑	x	RCL	R↓	x ^z y		
11		R↓	+	STO	÷		u	
12		x ^z y	RCL	÷	CHS		v	

Complex secant

Formula:

$$\sec(a + bi) = \frac{1}{\cos(a + bi)}$$

$$= u + vi$$

Example:

$$\sec(2 + 3i) = -0.04 + .09i$$

LINE	DATA	OPERATIONS					DISPLAY	REMARKS
1	b	↑						
2	a	↑	1	8	0	x		
3		π	÷	STO	cos	x ^z y		
4		↑	↑	R↓	R↓	e ^x		
5		↑	1/x	+	2	÷		
6		x	RCL	x ^z y	STO	CLX		
7		+	sin	x ^z y	e ^x	↑		
8		1/x	-	2	÷	x		
9		CHS	RCL	↑	x	x ^z y		
10		↑	↑	x	RCL	R↓		
11		x ^z y	R↓	+	STO	÷	u	
12		x ^z y	RCL	÷	CHS		v	

40 Complex Trigonometric and Hyperbolic Functions

Complex arc sine

Formula:

$$\begin{aligned}\arcsin(a + bi) &= \arcsin \beta + i \operatorname{sgn}(b) \ln(\alpha + \sqrt{\alpha^2 - 1}) \\ &= u + iv\end{aligned}$$

$$\text{where: } \alpha = \frac{1}{2} \sqrt{(a+1)^2 + b^2} + \frac{1}{2} \sqrt{(a-1)^2 + b^2}$$

$$\beta = \frac{1}{2} \sqrt{(a+1)^2 + b^2} - \frac{1}{2} \sqrt{(a-1)^2 + b^2}$$

$$\operatorname{sgn}(b) = \begin{cases} 1 & \text{if } b \geq 0 \\ -1 & \text{if } b < 0 \end{cases}.$$

Example:

$$\arcsin(5 + 8i) = .556 + 2.9387i$$

Note: Inverse trigonometric and inverse hyperbolic functions are multiple-valued functions, but only one answer will be given for each of them.

LINE	DATA	OPERATIONS	DISPLAY	REMARKS
1	a	↑		
2	b	↑ x STO x ² y ↑		
3		↑ 1 + ↑ x		
4		x ² y 1 - ↑ x		
5		RCL + √x 2 ÷		
6		STO R↓ + √x 2		
7		÷ ↑ ↑ RCL +		
8		x ² y RCL - arc sin		
9		π x 1 8 0		
10		÷	u	
11		R↓ ↑ ↑ x 1		
12		- √x + ln		If b ≥ 0, go to 14
13		CHS		
14			v	

Complex arc cosine

Formula:

$$\begin{aligned}\text{arc cos}(a + ib) &= \text{arc cos } \beta - i \operatorname{sgn}(b) \ln(\alpha + \sqrt{\alpha^2 - 1}) \\ &= u + iv\end{aligned}$$

$$\text{where: } \alpha = \frac{1}{2} \sqrt{(a+1)^2 + b^2} + \frac{1}{2} \sqrt{(a-1)^2 + b^2}$$

$$\beta = \frac{1}{2} \sqrt{(a+1)^2 + b^2} - \frac{1}{2} \sqrt{(a-1)^2 + b^2}$$

$$\operatorname{sgn}(b) = \begin{cases} 1 & \text{if } b \geq 0 \\ -1 & \text{if } b < 0 \end{cases}$$

Example:

$$\text{arc cos}(5 + 8i) = 1.0147 - 2.9387i$$

LINE	DATA	OPERATIONS	DISPLAY	REMARKS
1	a	↑ [] [] [] []		
2	b	↑ x STO x \leftrightarrow y ↑		
3		↑ 1 + ↑ x		
4		x \leftrightarrow y 1 - ↑ x		
5		RCL + \sqrt{x} 2 \div		
6		STO R↓ + \sqrt{x} 2		
7		\div ↑ ↑ RCL +		
8		x \leftrightarrow y RCL - arc cos		
9		π x 1 8 0		
10		\div [] [] [] []	u	
11		R↓ ↑ ↑ x 1		
12		- \sqrt{x} + ln []		If b < 0, go to 14
13		[] [] [] []		
14		CHS [] [] [] []	v	

42 Complex Trigonometric and Hyperbolic Functions

Complex arc tangent

Formula:

$$\begin{aligned}\text{arc tan}(a + bi) &= \frac{1}{2} \left[\pi - \text{arc tan} \frac{1+b}{a} - \text{arc tan} \frac{1-b}{a} \right] + \frac{i}{4} \ln \left[\frac{(1+b)^2 + a^2}{(1-b)^2 + a^2} \right] \\ &= u + iv\end{aligned}$$

where: $(a + bi)^2 \neq -1$.

Example:

$$\text{arc tan}(5 + 8i) = 1.5142 + .0898i$$

LINE	DATA	OPERATIONS	DISPLAY	REMARKS
1	b	↑ ↑ 1 +		
2	a	STO ÷ arc tan x↔y		
3		↑ 1 x↔y - ↑		
4		↑ RCL ÷ arc tan		
5		x↔y R↓ + x↔y CLX		
6		1 8 0 x↔y -		
7		2 ÷ π x 1		
8		8 0 ÷	u	
9		R↓ 2 x↔y - ↑		
10		x RCL ↑ x STO		
11		+ x↔y ↑ x RCL		
12		+ ÷ ln 4 ÷	v	

Complex arc cotangent

Formula:

$$\text{arc cot } (a + ib) = \frac{\pi}{2} - \text{arc tan } (a + ib) = u + iv$$

Example:

$$\text{arc cot } (5 + 8i) = .0566 - .0898i$$

LINE	DATA	OPERATIONS	DISPLAY	REMARKS
1	b	↑ ↑ 1 +		
2	a	STO ÷ arc tan x↔y		
3		↑ 1 x↔y - ↑		
4		↑ RCL ÷ arc tan		
5		x↔y R↓ + x↔y CLX		
6		+ 2 ÷ π x		
7		1 8 0 ÷	u	
8		R↓ 2 x↔y - ↑		
9		x RCL ↑ x STO		
10		+ x↔y ↑ x RCL		
11		+ ÷ ln 4 ÷		
12	CHS		v	

44 Complex Trigonometric and Hyperbolic Functions

Complex arc cosecant

Formula:

$$\text{arc csc } (a + ib) = \text{arc sin} \left(\frac{1}{a + ib} \right)$$

$$= u + iv$$

Example:

$$\text{arc csc } (5 + 8i) = .05598 - .0899i$$

$$(D < 0)$$

LINE	DATA	OPERATIONS	DISPLAY	REMARKS
1	b	↑ [] [] [] []		
2	a	↑ ↑ x STO CLX		
3		+ R↓ x RCL +		
4		STO ÷ CHS x ² y RCL		
5		÷ x ² y [] []	D	
6		↑ x STO x ² y ↑		
7		↑ 1 + ↑ x		
8		x ² y 1 - ↑ x		
9		RCL + √x 2 ÷		
10		STO R↓ + √x 2		
11		÷ ↑ ↑ RCL +		
12		x ² y RCL - arc sin		
13		π x 1 8 0		
14		÷ [] []	u	
15		R↓ ↑ ↑ x 1		
16		- √x + ln []	If D > 0, go to 18	
17		[] [] [] []		
18		CHS [] [] []	v	

Complex arc secant*Formula:*

$$\text{arc sec } (a + ib) = \text{arc cos} \left(\frac{1}{a + ib} \right)$$

$$= u + iv$$

Example:

$$\text{arc sec } (5 + 8i) = 1.5148 + .0899i$$

$$(D < 0)$$

LINE	DATA	OPERATIONS	DISPLAY	REMARKS
1	b	↑ [] [] [] []		
2	a	↑ ↑ x STO CLX		
3		+ R↓ x RCL +		
4		STO ÷ CHS x ² y RCL		
5		÷ x ² y [] []	D	
6		↑ x STO x ² y ↑		
7		↑ 1 + ↑ x		
8		x ² y 1 - ↑ x		
9		RCL + √x 2 ÷		
10		STO R↓ + √x 2		
11		÷ ↑ ↑ RCL +		
12		x ² y RCL - arc cos		
13		π x 1 8 0		
14		÷ [] [] []	u	
15		R↓ ↑ ↑ x 1		
16		- √x + ln []	If D < 0, go to 18	
17		[] [] [] []		
18		[] [] [] []	v	

46 Complex Trigonometric and Hyperbolic Functions

Complex hyperbolic sine

Formula:

$$\begin{aligned}\sinh(a + ib) &= -i \sin i(a + ib) \\ &= u + iv\end{aligned}$$

Example:

$$\sinh(3 - 2i) = -4.1689 - 9.154i$$

LINE	DATA	OPERATIONS	DISPLAY	REMARKS
1	a	↑		
2	b	CHS ↑ 1 8 0		
3		x π ÷ STO sin		
4		x ² y ↑ ↑ R↓ R↓		
5		e ^x ↑ 1/x + 2		
6		÷ x RCL x ² y STO		
7		CLX + cos x ² y e ^x		
8		↑ 1/x - 2 ÷		
9		x RCL CHS x ² y	u	
10		x ² y	v	

Complex hyperbolic cosine

Formula:

$$\begin{aligned}\cosh(a + ib) &= \cos i(a + ib) \\ &= u + iv\end{aligned}$$

Example:

$$\cosh(1 + 2i) = -.6421 + 1.0686i$$

LINE	DATA	OPERATIONS	DISPLAY	REMARKS
1	a	\uparrow [] [] [] []		
2	b	CHS \uparrow 1 8 0		
3		x π \div STO cos		
4		x^2y \uparrow \uparrow R↓ R↓		
5		e^x \uparrow $1/x$ + 2		
6		\div \times RCL x^2y STO	u	
7		CLX + sin x^2y e^x		
8		\uparrow $1/x$ - 2 \div		
9		x CHS [] [] []	v	

Complex hyperbolic tangent

Formula:

$$\begin{aligned}\tanh(a + ib) &= -i \tan i(a + ib) \\ &= u + iv\end{aligned}$$

Example:

$$\tanh(1 + 2i) = 1.1667 - .243i$$

LINE	DATA	OPERATIONS	DISPLAY	REMARKS
1	a	\uparrow [] [] [] []		
2	b	CHS \uparrow 2 \times STO		
3		\uparrow 1 8 0 \times		
4		π \div cos x^2y 2		
5		x \uparrow \uparrow R↓ R↓		
6		e^x \uparrow $1/x$ + 2		
7		\div + RCL x^2y STO		
8		CLX + x^2y e^x \uparrow		
9		$1/x$ - 2 \div RCL		
10		\div x^2y 1 8 0		
11		x π \div sin RCL		
12		\div CHS x^2y [] []	u	
13		x^2y [] [] []	v	

48 Complex Trigonometric and Hyperbolic Functions

Complex hyperbolic cotangent

Formula:

$$\coth(a + ib) = i \cot i(a + ib)$$

$$= u + iv$$

Example:

$$\coth(1 + 2i) = .8213 + .17138i$$

LINE	DATA	OPERATIONS	DISPLAY	REMARKS
1	a	↑ [] [] [] []		
2	b	CHS ↑ 2 x STO		
3		↑ 1 8 0 x		
4		π ÷ cos x↔y 2		
5		x ↑ ↑ R↓ R↓		
6		e^x ↑ 1/x + 2		
7		÷ x↔y - RCL x↔y		
8		STO CLX + x↔y e^x		
9		↑ 1/x - 2 ÷		
10		RCL ÷ CHS x↔y 1		
11		8 0 x π ÷		
12		sin RCL ÷ x↔y CHS	u	
13		x↔y [] [] []	v	

Complex hyperbolic cosecant

Formula:

$$\operatorname{csch}(a + ib) = i \csc i(a + ib)$$

$$= u + iv$$

Example:

$$\operatorname{csch}(1 + 2i) = -.2215 - .63549i$$

LINE	DATA	OPERATIONS					DISPLAY	REMARKS
1	a	↑						
2	b	CHS	↑	1	8	0		
3		x	π	÷	STO	sin		
4		x ² y	↑	↑	R↓	R↓		
5		e ^x	↑	1/x	+	2		
6		÷	x	RCL	x ² y	STO		
7		CLX	+	cos	x ² y	e ^x		
8		↑	1/x	-	2	÷		
9		x	RCL	↑	x	x ² y		
10		↑	↑	x	RCL	R↓		
11		x ² y	R↓	+	STO	÷		
12		x ² y	RCL	÷			u	
13		x ² y					v	

Complex hyperbolic secant

Formula:

$$\operatorname{sech}(a + bi) = \sec i(a + bi)$$

$$= u + iv$$

Example:

$$\operatorname{sech}(1 + 2i) = -.4131 - .6875i$$

LINE	DATA	OPERATIONS					DISPLAY	REMARKS
1	a	↑						
2	b	CHS	↑	1	8	0		
3		x	π	÷	STO	cos		
4		x ² y	↑	↑	R↓	R↓		
5		e ^x	↑	1/x	+	2		
6		÷	x	RCL	x ² y	STO		
7		CLX	+	sin	x ² y	e ^x		
8		↑	1/x	-	2	÷		
9		x	CHS	RCL	↑	x		
10		x ² y	↑	↑	x	RCL		
11		R↓	x ² y	R↓	+	STO		
12		÷					u	
13		x ² y	RCL	÷	CHS		v	

50 Complex Trigonometric and Hyperbolic Functions

Complex inverse hyperbolic sine

Formula:

$$\begin{aligned}\sinh^{-1}(a + ib) &= -i \operatorname{arc sin} i(a + ib) \\ &= u + iv\end{aligned}$$

Example:

$$\sinh^{-1}(8 - 5i) = 2.9387 - .556i$$

LINE	DATA	OPERATIONS	DISPLAY	REMARKS
1	a	\uparrow \square \square \square \square		
2	b	CHS $x \leftrightarrow y$ \uparrow \times STO		
3		$x \leftrightarrow y$ \uparrow \uparrow 1 +		
4		\uparrow \times $x \leftrightarrow y$ 1 -		
5		\uparrow \times RCL + \sqrt{x}		
6		2 \div STO R↓ +		
7		\sqrt{x} 2 \div \uparrow \uparrow		
8		RCL + $x \leftrightarrow y$ RCL -		
9		arc sin π \times 1		
10		8 0 \div CHS STO		
11		R↓ \uparrow \uparrow \times 1		
12		- \sqrt{x} + ln \square		If $a \geq 0$, go to 14
13		CHS \square \square \square		
14		\square \square \square \square	u	
15		RCL \square \square \square	v	

Complex inverse hyperbolic cosine

Formula:

$$\begin{aligned}\cosh^{-1}(a + ib) &= i \operatorname{arc cos} (a + ib) \\ &= u + iv\end{aligned}$$

Example:

$$\cosh^{-1}(5 + 8i) = 2.9387 + 1.0147i$$

LINE	DATA	OPERATIONS	DISPLAY	REMARKS
1	a	\uparrow \uparrow \uparrow \uparrow \uparrow		
2	b	\uparrow \times STO $x \leftrightarrow y$ \uparrow		
3		\uparrow 1 $+$ \uparrow x		
4		$x \leftrightarrow y$ 1 $-$ \uparrow x		
5		RCL $+$ \sqrt{x} 2 \div		
6		STO R↓ $+$ \sqrt{x} 2		
7		\div \uparrow \uparrow RCL $+$		
8		$x \leftrightarrow y$ RCL $-$ arc cos		
9		π x 1 8 0		
10		\div STO R↓ \uparrow \uparrow		
11		x 1 $-$ \sqrt{x} $+$		
12		In		If $b \geq 0$, go to 14
13		CHS		
14			u	
15		RCL	v	

Complex inverse hyperbolic tangent

Formula:

$$\tanh^{-1}(a + ib) = -i \operatorname{arc tan} i(a + ib)$$

$$= u + iv$$

Example:

$$\tanh^{-1}(8 - 5i) = .0898 - 1.5142i$$

LINE	DATA	OPERATIONS	DISPLAY	REMARKS
1	a	\uparrow \uparrow 1 $+$		
2	b	CHS STO \div arc tan		
3		$x \leftrightarrow y$ \uparrow 1 $x \leftrightarrow y$ $-$		
4		\uparrow \uparrow RCL \div arc		
5		tan $x \leftrightarrow y$ R↓ $+$ $x \leftrightarrow y$		
6		CLX 1 8 0 $x \leftrightarrow y$		
7		$-$ 2 \div π x		
8		1 8 0 \div CHS	v	
9		R↓ 2 $x \leftrightarrow y$ $-$ \uparrow		
10		x RCL \uparrow x STO		
11		$+$ $x \leftrightarrow y$ \uparrow x RCL		
12		$+$ \div ln 4 \div	u	

52 Complex Trigonometric and Hyperbolic Functions

Complex inverse hyperbolic cotangent

Formula:

$$\coth^{-1} (a + ib) = i \operatorname{arc cot} i(a + ib)$$

$$= u + iv$$

Example:

$$\coth^{-1} (8 - 5i) = .0898 + .0566i$$

LINE	DATA	OPERATIONS	DISPLAY	REMARKS
1	a	↑ ↑ 1 +		
2	b	CHS STO ÷ arc tan		
3		x ⁻² y ↑ 1 x ⁻² y -		
4		↑ ↑ RCL ÷ arc		
5		tan x ⁻² y R↓ + x ⁻² y		
6		CLX + 2 ÷ π		
7		x 1 8 0 ÷	v	
8		R↓ 2 x ⁻² y - ↑		
9		x RCL ↑ x STO		
10		+ x ⁻² y ↑ x RCL		
11		+ ÷ ln 4 ÷	u	

Complex inverse hyperbolic cosecant*Formula:*

$$\begin{aligned}\operatorname{csch}^{-1}(a+ib) &= i \operatorname{arc csc} i(a+ib) \\ &= u + iv\end{aligned}$$

Example:

$$\begin{aligned}\operatorname{csch}^{-1}(8 - 5i) &= .0899 + .05598i \\ (D < 0) &\end{aligned}$$

LINE	DATA	OPERATIONS					DISPLAY	REMARKS
1	a	↑						
2	b	CHS	↑	↑	x	STO		
3		CLX	+	R↓	x	RCL		
4		+	STO	÷	CHS	x↔y		
5		RCL	÷	x↔y			D	
6		↑	x	STO	x↔y	↑		
7		↑	1	+	↑	x		
8		x↔y	1	-	↑	x		
9		RCL	+	√x	2	÷		
10		STO	R↓	+	√x	2		
11		÷	↑	↑	RCL	+		
12		x↔y	RCL	-	arc	sin		
13		π	x	1	8	0		
14		÷	STO	R↓	↑	↑		
15		x	1	-	√x	+		
16		In					If D < 0, go to 18.	
17		CHS						
18							u	
19		RCL					v	

54 Complex Trigonometric and Hyperbolic Functions

Complex inverse hyperbolic secant

Formula:

$$\begin{aligned}\operatorname{sech}^{-1}(a + ib) &= i \operatorname{arc sec}(a + ib) \\ &= u + iv\end{aligned}$$

Example:

$$\begin{aligned}\operatorname{sech}^{-1}(5 + 8i) &= -0.0899 + 1.5148i \\ (D < 0)\end{aligned}$$

LINE	DATA	OPERATIONS	DISPLAY	REMARKS
1	b	↑ [] [] [] []		
2	a	↑ ↑ x STO CLX		
3		+ R↓ x RCL +		
4		STO ÷ CHS x ² y RCL		
5		÷ x ² y [] [] []	D	
6		↑ x STO x ² y ↑		
7		↑ 1 + ↑ x		
8		x ² y 1 - ↑ x		
9		RCL + √x 2 ÷		
10		STO R↓ + √x 2		
11		÷ ↑ ↑ RCL +		
12		x ² y RCL - arc cos		
13		π x 1 8 0		
14		÷ STO R↓ ↑ ↑		
15		x 1 - √x +		
16		In [] [] []	If D ≥ 0, go to 18.	
		[] [] []		
17		CHS [] [] []		
18		[] [] [] []	u	
19		RCL [] [] []	v	

Compound Interest

See page 96

Conversions

Repetitive use of formulas with two constants.

The technique used here is to store both constants—one in the S register, and the other in the T register. After initially storing the two constants, the formulas can be used repeatedly without limit.

Centigrade to Fahrenheit ($a^{\circ}\text{C} \rightarrow b^{\circ}\text{F}$)

Formula:

$$b = \frac{9}{5} a + 32$$

Examples:

a	-30	0	28	100	539
b	-22.	32.	82.4	212.	1002.2

LINE	DATA	OPERATIONS					DISPLAY	REMARKS
1		3	2	STO	9	↑		
2		5	÷	↑	↑	↑		
3		CLX						
4	a	x	RCL	+			b	Stop. For new case, go to 3

Fahrenheit to centigrade ($b^{\circ}\text{F} \rightarrow a^{\circ}\text{C}$)

Formula:

$$a = \frac{5}{9} (b - 32)$$

Examples:

b	-460	-40	0	32	212
a	-273.33	-40.	-17.78	0.	100.

LINE	DATA	OPERATIONS					DISPLAY	REMARKS
1		3	2	STO	5	↑		
2		9	÷	↑	↑	↑		
3		CLX						
4	b	RCL	-	x				Stop. For new case, go to 3

56 Conversions

Feet and inches to centimeters

Formulas:

$$1 \text{ foot} = 12 \text{ inches}$$

$$1 \text{ inch} = 2.54 \text{ cm}$$

Examples:

$$4' 8'' = 142.24 \text{ cm}$$

$$5' 5'' = 165.1 \text{ cm}$$

$$6' 3'' = 190.5 \text{ cm}$$

LINE	DATA	OPERATIONS	DISPLAY	REMARKS
1		2 . 5 4 STO		
2		1 2 ↑ ↑ ↑		
3		CLX		
4	Feet	x		
5	Inches	+ RCL x	Centimeters	Stop. For new case, go to 3

Centimeters to feet and inches

Examples:

$$164 \text{ cm} = 5' 4.6''$$

$$180 \text{ cm} = 5' 10.9''$$

LINE	DATA	OPERATIONS	DISPLAY	REMARKS
1		1 2 STO 2 .		
2		5 4 1/x ↑ ↑		
3		↑		
4		CLX		
5	Centimeters	x RCL ÷	D	Let f = integer part of D
6	f		Feet	
7		- RCL x	Inches	Stop. For new case, go to 4

Coordinate Conversions


Polar to rectangular

Formulas:

$$x = R \cos \theta$$

$$y = R \sin \theta$$

(θ is in degrees)



Example:

If $\theta = 40^\circ$, and $R = 5.1$, then

$x = 3.907$, and

$y = 3.278$.


LINE	DATA	OPERATIONS	DISPLAY	REMARKS
1	θ	\uparrow [] [] [] []		
2	R	STO $x \leftrightarrow y$ \uparrow cos RCL		
3		x [] [] []	x	
4		$x \leftrightarrow y$ sin RCL x []	y	

Rectangular to polar

Formulas:

$$R = \sqrt{x^2 + y^2},$$

$$\theta = \arctan\left(\frac{y}{x}\right)$$



Example:

If $x = 5.7$, and $y = 4.2$, then

$R = 7.08$, and

$\theta = 36.38$.

58 Coordinate Conversions

LINE	DATA	OPERATIONS	DISPLAY	REMARKS
1	y	↑ [] [] [] []		
2	x	[] [] [] []		If $x \neq 0$, go to 4
		[] [] [] []		
3		CLX EEX CHS 9 9		
4		STO x \leftrightarrow y ↑ ↑ x		
5		RCL ↑ x + \sqrt{x}	R	
6		x \leftrightarrow y RCL \div arc tan		If $x \geq 0$, go to 8
		[] [] [] []		
7		1 8 0 +		
8		[] [] [] []	θ	

Correlation Coefficient

See page 144

Cosecant

See page 156

Cotangent

See page 155

Coversine

See page 157

Cross Product

See page 160

Cross Product Sum

See page 144

Degrees, Minutes, Seconds to Decimal Degrees

See page 11

Decimal Degrees to Degrees, Minutes, Seconds

See page 11

Degrees to Grads

See page 15

Degrees to Mils

See page 14

Degrees to Radians

See page 14

Discount

See page 95

Dot Product

See page 161

Double Precision Product*Terms used:*

A and **B** are the two numbers to be multiplied (each having up to 10 significant digits).

A_{L5} and **B_{L5}** are the left most 5 significant digits, of A and B.

A_{R5} and **B_{R5}** are the remaining significant digits, (zero filled at the right if the number of significant digits is less than 10).

A_I and **B_I** are integers with the same significant digits as A and B, respectively.

A_n and **B_n** are the number of significant digits each in A and B.

X_{R5} is the rightmost portion (5 digits) of the display, regardless of how many digits are displayed.

X_{L1} is the leftmost digit of the display.

X_{Exp} is the value of the displayed exponent (=0, if none visible).

X_{LD} is the number of displayed digits to the left of the decimal point (= -1 if the number is of the form .od ---).

X_{ND} is the total number of displayed digits.

60 Double Precision Product

LINE	DATA	OPERATIONS	DISPLAY	REMARKS
1	A	\uparrow \uparrow \uparrow \uparrow \uparrow		
2	B	x STO \uparrow \uparrow \uparrow		
3	A_{L5}	\uparrow \uparrow \uparrow \uparrow \uparrow		
4	B_{R5}	x \uparrow \uparrow \uparrow \uparrow		
5	B_{L5}	\uparrow \uparrow \uparrow \uparrow \uparrow		
6	A_{R5}	x \uparrow \uparrow CLX +		
7		$R\downarrow$ + \uparrow \uparrow \uparrow	D1	If D1 < 10^{10} , go to 11
		\uparrow \uparrow \uparrow \uparrow \uparrow		
8		CLX 1 \uparrow EEX 1		
9		0 - + + 1		
10		- \uparrow \uparrow \uparrow \uparrow		
11	X_{R5}	0 0 0 0 0		
12		\uparrow \uparrow \uparrow \uparrow \uparrow		
13	A_{R5}	\uparrow \uparrow \uparrow \uparrow \uparrow		
14	B_{R5}	x \uparrow \uparrow CLX +		
15		$R\downarrow$ + \uparrow \uparrow \uparrow	D2	If D2 < 10^{10} , go to 19
		\uparrow \uparrow \uparrow \uparrow \uparrow		
16		CLX 1 \uparrow EEX 1		
17		0 - + + 1		
18		- \uparrow \uparrow \uparrow \uparrow		
19	A_n	\uparrow \uparrow \uparrow \uparrow \uparrow		
20	B_n	+ \uparrow \uparrow \uparrow \uparrow		
21	A_1	\uparrow \uparrow \uparrow \uparrow \uparrow	A_1	Read note 1
22		CLX \uparrow \uparrow \uparrow \uparrow		
23	999999997	\uparrow \uparrow \uparrow \uparrow \uparrow		Go to 26
		\uparrow \uparrow \uparrow \uparrow \uparrow		
24		CLX \uparrow \uparrow \uparrow \uparrow		
25	9999999949	\uparrow \uparrow \uparrow \uparrow \uparrow		
26		log \uparrow \uparrow \uparrow \uparrow		
27	B_1	\uparrow \uparrow \uparrow \uparrow \uparrow	B_1	Read note 1
		\uparrow \uparrow \uparrow \uparrow \uparrow		
28		CLX \uparrow \uparrow \uparrow \uparrow		
29	999999997	\uparrow \uparrow \uparrow \uparrow \uparrow		Go to 32
		\uparrow \uparrow \uparrow \uparrow \uparrow		
30		CLX \uparrow \uparrow \uparrow \uparrow		
31	9999999949	\uparrow \uparrow \uparrow \uparrow \uparrow		

Continued

LINE	DATA	OPERATIONS	DISPLAY	REMARKS
32		log +	D3	If D3 is an integer, go to 35; otherwise, let f be the integer part of D3.
33		CLX		
34	f			
35		↑ 1 + -	D4	If D4 = 0, go to 37
36		R↓		If $X_{ND} = 10$, go to 39; otherwise, go to 41.
37		R↓		If $x_{ND} < 10$, go to 41; if $X_{L1} > 5$, go to 42
38				Go to 40
39	X_{L1}	EEX 9 -		
40		1 0 x		If $X_{ND} < 10$, go to 41, if $X_{L1} > 5$, go to 42
41		RCL		Go to 47
42		RCL		Let E = X_{EXP} , D = X_{LD}
43	E	↑ 1 0 -		
44	D	+	P	
45		CLX EEX		
46	P	-		
47				Read note 2
48		$x \leftrightarrow y$		Read note 3

Note 1: If A_I (B_I) contains 9 digits of which the first 8 are all 9's and the last digit is 8 or 9, go to 22 (28).

If A_I (B_I) contains 10 digits of which the first 8 are all 9's and the last 2 are 50 or more, go to 24 (30).

Otherwise, go to 26 (32).

Note 2: Read the sign, the leftmost 10 significant digits of the product, the decimal point, and the exponent. If the number starts with .0, the tenth significant digit is hidden. Multiply by 10 to read it. If less than 10 digits are displayed, the balance of the rightmost blank positions are significant zeros (providing the Y register is non-zero).

Note 3: Read the remaining significant digits. If less than 10 digits are displayed, fill in enough preceding zeros on the left to make 10 digits.

62 Double Precision Product

Example:

If $A = 349.27186$, $B = -2.431985517 \times 10^{12}$,

then $A_{L5} = 34927$, $B_{L5} = 24319$

$A_{R5} = 18600$, $B_{R5} = 85517$

$A_I = 34927186$, $B_I = 2431985517$

$A_n = 8$, $B_n = 10$

and $A \times B = -8.4942410501565162 \times 10^{14}$.

(Note that $A_{L5}, B_{L5}, A_{R5}, B_{R5}, A_I, B_I, A_n, B_n$ are all positive integers that contain neither signs nor decimal points.)

LINE	DATA	OPERATIONS	DISPLAY	REMARKS
1	349.27186	↑ CHS		
2	2.431985517	EEX 1 2 × STO		
3	34927	↑		
4	85517	x		
5	24319	↑		
6	18600	x ↑ ↑ CLX +		
7		R↓ +	3439185659.	← D1 < 10 ¹⁰
8	8565900000	↑		
9	18600	↑		
10	85517	x ↑ ↑ CLX +		
11		R↓ +	1.01565162	10 ← D2 ≥ 10 ¹⁰
12		CLX 1 ↑ EEX 1		
13	0	- + + 1		
14	-		156516200.	
15	8	↑ 1 0 +		
16	34927186	log		
17	2431985517	log +	16.92912458	← D3
18		CLX 1 6 ↑ 1		
19		+ -	1.	← D4
20		R↓	156516200.	X _{ND} = 9
21		1 0 x	1565162000.	X _{ND} = 10, X _{L1} = 1
22		RCL	-8.49424105 14	Note: only 9 digits are displayed.
23		x ² y	1565162000.	

Equation Solving (Iterative Techniques)

We will deal here with equations of the form

$$x = f(x)$$

for cases where it is difficult to separate all x 's to one side of the equal sign. The iterative approach is illustrated through the solution of selected equations.

Example 1. Find x such that $x = e^{-x}$

Method:

Choose $X_a = 5$ as an approximation for the solution. Then after 44 iterations, the answer is $X = .5671432904$.

Note: The algorithm will converge to $.5671432904$ after about 50 iterations for any value of X_a .

LINE	DATA	OPERATIONS	DISPLAY	REMARKS
1	5	CHS e ^x [] [] [] []	6.737946999 -03	
2		CHS e ^x [] [] [] []		Perform line 2 forty-three times
3		[] [] [] [] []	.5671432904	

Example 2. $4 = X - \frac{1}{X}$

Method:

Rewrite the equation as

$$X = \frac{1}{X} + 4$$

choose an approximate solution for X (e.g., $X_a = 4$).

Answer:

4.236067978

LINE	DATA	OPERATIONS	DISPLAY	REMARKS
1	4	1/x 4 + [] []	4.25	
2		1/x 4 + [] []		Perform line 2 seven times
3		[] [] [] []	4.236067978	

64 Equation Solving (Iterative Techniques)

Example 3. $X^X = 1000$

Method:

Rewrite the equation in the form

$$x = \ln 1000 / \ln x$$

Pick an approximation x_a for x (e.g., $x_a = 4$). If we use the following sequence, convergence is from both sides and will take a long time.

LINE	DATA	OPERATIONS	DISPLAY	REMARKS
1	1000	\ln \uparrow \uparrow \uparrow		
2	4	\ln \div	4.982892143	
3		\ln \div	4.301189432	
4		\ln \div	4.7349339	
5		\ln \div	4.442378437	
6		\ln \div	4.632377942	
7		\ln \div	4.505830645	
8		\ln \div	4.588735608	
9		\ln \div	4.533824354	
10		\ln \div	4.569933525	etc.

To hasten the convergence, we modify the loop and use the average of the last two approximations as the new approximation.

Answer:

4.555535705

LINE	DATA	OPERATIONS	DISPLAY	REMARKS
1	1000	\ln \uparrow \uparrow \uparrow		
2	4	STO \ln \div RCL $+$		
3		2 \div STO	4.491446072	
4		\ln \div RCL $+$		
5		2 \div STO	4.544974651	
6		\ln \div RCL $+$		
7		2 \div STO	4.553746975	
8		\ln \div RCL $+$		
9		2 \div STO	4.555231426	
24		\ln \div RCL $+$		
25		2 \div STO	4.555535705	
26		\ln \div RCL $+$		
27		2 \div STO	4.555535705	Display converges

7 iterations

Example 4. Largest x^{x^x}

What is the largest value of x such that x^{x^x} does not overflow in the HP-35 (i.e. $x^{x^x} < 9.999999999 \times 10^{99}$)?

Method:

Since $9.999999999 \times 10^{99}$ is only slightly less than a *googol* (10^{100}), let us call this constant G. Then,

$$x^{x^x} = G$$

$$x^x \ln x = \ln G$$

$$x^x = \ln G / \ln x$$

$$x \ln x = \ln(\ln G / \ln x)$$

$$x = [\ln(\ln G / \ln x)] / \ln x$$

Use 3 as an initial approximation for the solution.

The loop in the following routine alternates between the two answers, 3.830482865 and 3.830482864. Upon substitution of both values in x^{x^x} , 3.830482865 gives an overflow while 3.830482864 does not. Thus, the answer is 3.830482864.

LINE	DATA	OPERATIONS	DISPLAY	REMARKS
1		9 e ^x e ^x ln ↑		
2		↑ ↑ [] [] []	230.2585093	← ln G
3		3 ln ÷ ln 3		
4		ln ÷ STO [] []	4.865369574	
5		ln ÷ ln RCL ln		
6		÷ RCL + [] []		
7		2 ÷ STO [] []	4.006633409	
8		ln ÷ ln RCL ln		
9		÷ RCL + [] []		
10		2 ÷ STO [] []	3.84465425	
29		[] [] [] [] []		
30		[] [] [] [] []		
31		[] [] [] [] []	3.830482865	
32		[] [] [] [] []		
33		[] [] [] [] []		
34		2 ÷ STO [] []	3.830482864	etc.

} 6 iterations

66 Equation Solving (Iterative Techniques)

Example 5. $x^2 + 4 \sin x = 0$

(x is in radians).

Method:

Rewrite the equation as

$$x = \pm \sqrt{-4 \sin x}.$$

Since x is in radians, we convert to degrees before using it with the trigonometric function keys.

Upon plotting this curve, we see that there is a root near -2, so take $x_a = -2$ as an approximation of the solution, and substitute in

$$x = -\sqrt{-4 \sin x}.$$

Note: This algorithm converges from both sides, to hasten convergence, the following routine averages two approximations to get the next approximation.

It is only necessary to do the loop 5 times to get 4 digits of accuracy.

LINE	DATA	OPERATIONS	DISPLAY	REMARKS
1	2	CHS STO <input type="text"/> <input type="text"/> <input type="text"/>		
2		<input type="text"/> 1 8 0 <input type="text"/> x		Perform lines 2–5
3		<input type="text"/> ÷ sin CHS 4		until A_i does not change.
4		x \sqrt{x} RCL – 2		
5		÷ CHS STO <input type="text"/> <input type="text"/> <input type="text"/>	A_i	
6		<input type="text"/> <input type="text"/> <input type="text"/> <input type="text"/> <input type="text"/>	-1.933753764	After 17 iterations

A method for faster convergence

Some equations for $f(x) = 0$ converge very slowly by the above methods. The following method gives faster convergence.

Formula:

$$x_{i+1} = x_i - \frac{E_i (x_i - x_{i-1})}{E_i - E_{i-1}}$$

where $i = 1, 2, 3, \dots$

x_i = current trial value

x_{i+1} = next trial value

x_{i-1} = previous trial value

E_i = current error = $f(x_i)$

E_{i-1} = last error = $f(x_{i-1})$

L = lower bound for the solution

U = upper bound for the solution.

The keystrokes evaluating the equation $f(x)$ may not modify the S register, because the trial value is stored there.

Example 6. Solve $x^3 = 3^x$ where $1 < x < e$

Method:

Rewrite the equation in the form

$$x^3 - 3^x = 0.$$

Replace $f(x)$ in the routine (shown after example 7) by the key sequence:

The sequence consists of three up arrow keys followed by the digit 3, then the x^y key, then another x^y key, then the digit 3, then another x^y key, and finally a minus sign key.

Let $L = 1$

$U = e.$

Answer:

2.478052685 ($E_0 = -2.000000001$, $E_1 = .27254617$, $E_2 = .05608461$, $E_3 = -.03342142$, $E_4 = 1.19176 \times 10^{-3}$, $E_5 = 2.254 \times 10^{-5}$, $E_6 = E_7$)

68 Equation Solving (Iterative Techniques)

Example 7. Find a root of a polynomial

$$f(x) = x^4 - 4x^3 + 8x^2 + 20x - 65.$$

Method:

Replace $f(x)$ in the program by

$\uparrow \uparrow \uparrow 4 - \times 8 + \times 20 + \times 65 -$

Note that $f(2) = -9 < 0$, $f(3) = 40 > 0$, so there is a root between 2 and 3.

Let $L = 2$

$U = 3$

Answer:

2.236067977 (or $\sqrt{5}$) ($x_0 = 2$, $x_1 = 3$, $x_2 = 2.183673469$,
 $x_3 = 2.224244398$, $x_4 = 2.236236914$, $x_5 = 2.236067428$,
 $x_6 = x_7 = 2.236067977$)

LINE	DATA	OPERATIONS	DISPLAY	REMARKS
1	L	STO $\uparrow \uparrow \uparrow$		
2		f(x) $\uparrow \uparrow \uparrow$	E_0	Replace f(x) by proper keystrokes
3	U	$\uparrow \uparrow \uparrow$		
4		f(x) $\uparrow \uparrow \uparrow$	E_i	Perform lines 4–8 for $i = 1$,
5		RCL R↓ R↓ STO $x^{-2}y$		2, ..., until either the last
6		CLX + $x^{-2}y$ – x		two trial values are the same
7		RCL $x^{-2}y$ R↓ R↓ R↓		or the last two errors are
8	E_{i-1}	– ÷ – $\uparrow \uparrow$	X_{i+1}	the same.

Euler Numbers

Compute the n^{th} Euler number

Formula

$$E_n = \frac{2^{2n+2} (2n)!}{\pi^{2n+1}} \left[1 - \frac{1}{3^{2n+1}} + \frac{1}{5^{2n+1}} - \frac{1}{7^{2n+1}} + \dots \right]$$

Note: The Euler numbers are 1, 5, 61, 1385, 50521, ...

Example:

The 5th Euler number = 50521.

LINE	DATA	OPERATIONS	DISPLAY	REMARKS
1	n	↑ 2 x ↑ ↑		
2		↑	D	If D ≤ 2 , go to 4
3	i	x		Perform line 3 for i = D-1,
				D-2, ..., 2
4		x ² y 2 + 2 x ^y		
5		x x ² y 1 + π		
6		x ^y ÷ EEX 9 +		
7		EEX 9 -		

Exponentiation

Multiple successive power operations

As written, these terms must be executed from right to left. For example,

$$e^x^{1.5}$$

means

$$e^{(x^{1.5})}, \text{ not } (e^x)^{1.5}$$

$$a^{b^c}$$

means

$$a^{(b^c)}, \text{ not } (a^b)^c.$$

Example:

$$\text{Evaluate } t^{t^{t^{\dots}}} \text{ where } t = e^{\frac{1}{e}}.$$

Answer:

$$1.98957$$

LINE	DATA	OPERATIONS	DISPLAY	REMARKS
1		1 e^x $1/x$ e^x \uparrow		
2		STO x^y RCL x^y RCL		
3		x^y [] [] [] []		

Example:

$$\text{Find the limit of } t^{t^{t^{\dots}}} \text{ where } t = \sqrt{3}.$$

Answer:

$$\infty$$

LINE	DATA	OPERATIONS	DISPLAY	REMARKS
1		3 \sqrt{x} \uparrow STO []		
2		RCL x^y [] []		Perform line 2 six times
3		[] [] [] [] []	9.999999999 99	

Example:

Find L, the limit of t^t where $t = \sqrt{2}$.

Answer:

2 (rounded)

LINE	DATA	OPERATIONS	DISPLAY	REMARKS
1		2 \sqrt{x} \uparrow STO		
2		RCL x^y	L_i	Perform line 2 until $L_i = L_{i+1}$
3			1.999999995	Unchanging L after 54 iterations

e^x for large positive x ($x > 230$)

Formula:

$$\text{Suppose } e^x = a \times 10^b$$

$$\text{where } a < 10$$

Find a, b for a given x.

Example:

$$e^{300} = 1.942426525 \times 10^{130}$$

LINE	DATA	OPERATIONS	DISPLAY	REMARKS
1	x	\uparrow 1 0 ln STO		
2		\div \uparrow \uparrow \uparrow .		
3		5 - EEX 9 +		
4		EEX 9 - \uparrow R↓		
5		- RCL x e^x	a	
6		R↓ R↓	b	

72 Exponentiation

x^y for large x and/or y ($y \ln x > 230$)

Formula:

Suppose

$$x^y = a \times 10^b$$

where

$$a < 10,$$

find a, b for a given x, y.

Example:

$$75^{100} = 3.207202635 \times 10^{187}$$

LINE	DATA	OPERATIONS	DISPLAY	REMARKS
1	y	\uparrow [] [] [] []		
2	x	[ln] [x] [1] [0] [ln]		
3		[STO] [\div] [\uparrow] [\uparrow] [\uparrow]		
4		[.] [5] [-] [EEX] [9]		
5		[+] [EEX] [9] [-] [\uparrow]		
6		[R↓] [-] [RCL] [x] [e ^x]	a	
7		[R↓] [R↓] [] [] []	b	

Example:

The largest number that can be written with three digits and no other symbols is 9^9 . How big is this number?

Answer:

$$9^9 = 3.981071706 \times 10^{369693099}$$

(Due to machine accuracy limitations, this is only an approximate answer.)

LINE	DATA	OPERATIONS	DISPLAY	REMARKS
1		[9] [\uparrow] [\uparrow] [\uparrow] [x]		
2		[x] [x] [x] [x] [x]		
3		[x] [x] [9] [ln] [x]		
4		[1] [0] [ln] [STO] [\div]		
5		[\uparrow] [\uparrow] [\uparrow] [.] [5]		
6		[-] [EEX] [9] [+] [EEX]		
7		[9] [-] [\uparrow] [R↓] [-]		
8		[RCL] [x] [e ^x] [] []	3.981071706	
9		[R↓] [R↓] [] [] []	369693099.	

Converging u^u

Formula:

If $0 < x < e$ and $u = x^{\frac{1}{x}}$, then u^u will converge at x .

Example:

Let $x = 1.5$, $u = x^{\frac{1}{x}}$

Answer:

u^u converges at 1.5 in 21 iterations. (Display = 1.499999999).

LINE	DATA	OPERATIONS	DISPLAY	REMARKS
1	x	↑ $\frac{1}{x}$ $x \div y$ x^y STO	u	
2		↑		
3		RCL x^y		Perform line 3 as many times
				as necessary to see convergence

Diverging u^u

Formula:

If $u > e^e$ ($= 1.444667861$), u^u will diverge.

Example:

Let $u = 1.45$

Answer:

u^u diverges (overflows) after 43 iterations.

LINE	DATA	OPERATIONS	DISPLAY	REMARKS
1	u	STO ↑		
2		RCL x^y		Perform line 2 as many times
				as necessary to see divergence

74 Exponentiation

Calculator limits for x^y

- For a given positive value of x , how big can y be without causing x^y to overflow?

Example:

If $x = 50$, then y can be as large as around 58 (50^{59} will overflow).

Note: The following gives an approximate, not exact answer.

LINE	DATA	OPERATIONS	DISPLAY	REMARKS
1		2 3 0 ↑		
2	x	In ÷	D	Answer is the largest integer $\leq D$.

- For a given positive value of y , how big can x be without causing x^y to overflow?

Example:

If $y = 50$, then we can take x as large as around 99 (100^{50} will overflow).

Note: The following gives an approximate, not exact answer.

LINE	DATA	OPERATIONS	DISPLAY	REMARKS
1		2 3 0 ↑		
2	y	÷ e ^x	D	Answer is the largest integer $\leq D$.

Factorial

Formula:

$$x! = x \cdot (x - 1) \cdot \dots \cdot 2 \cdot 1$$

(x is a positive integer).

Example:

$$9! = 362880.$$

Note: $0! = 1$.

LINE	DATA	OPERATIONS	DISPLAY	REMARKS
1	x	\uparrow <input type="text"/> <input type="text"/> <input type="text"/> <input type="text"/> <input type="text"/>		If $x \leq 2$, Stop
		<input type="text"/> <input type="text"/> <input type="text"/> <input type="text"/> <input type="text"/>		
2	i	\times <input type="text"/> <input type="text"/> <input type="text"/> <input type="text"/> <input type="text"/>		Perform line 2 for $i = x-1$,
		<input type="text"/> <input type="text"/> <input type="text"/> <input type="text"/> <input type="text"/>		$x-2, \dots, 2$

Factorial and Gamma Function

Stirling's approximation

Notes:

This approximation can be used for positive $x < 69$ (otherwise the answer is $> 10^{100}$).

This approximation is good for large x .

For $x < 1$, use polynomial approximation.

To compute Gamma function:

$$\Gamma(x) = (x - 1)!$$

Formula:

$$x! \cong \sqrt{2\pi x} x^x e^{-x} \left(1 + \frac{1}{12x} + \frac{1}{288x^2} - \frac{139}{51840x^3} - \frac{571}{2488320x^4} \right)$$

76 Factorial and Gamma Function

Example:

$$4.25! \cong 35.21159$$

LINE	DATA	OPERATIONS	DISPLAY	REMARKS
1	x	STO ↑ ↑ 2 x		
2		π x √x RCL 2		
3		÷ RCL x ^y x RCL		
4		e ^x 1/x x RCL 2		
5		÷ RCL x ^y x STO		
6		x ^{-y} 1/x ↑ ↑ ↑		
7		CHS 5 7 1 ↑		
8	2488320	÷ x 1 3 9		
9		↑ 5 1 8 4		
10		0 ÷ - x 2		
11		8 8 1/x + x		
12		1 2 1/x + x		
13		1 + RCL x		

Polynomial approximation

Notes:

This approximation can be used for positive $x < 69$.

This is a more accurate method (to 6 or 7 decimal places), but longer than Stirling's approximation.

Formula:

$$x! \cong 1 + b_1 x + b_2 x^2 + \dots + b_8 x^8$$

for $0 \leq x \leq 1$

$$\text{where } b_1 = -.577191652, \quad b_2 = .988205891$$

$$b_3 = -.897056937, \quad b_4 = .918206857$$

$$b_5 = -.756704078, \quad b_6 = .482199394$$

$$b_7 = -.193527818, \quad b_8 = .035868343$$

Example:

$$4.25! \cong 35.21161962$$

LINE	DATA	OPERATIONS	DISPLAY	REMARKS
1	x	↑		If $x > 1$, go to 3
2		1 STO R↓		Go to 8
3				If $x > 2$, go to 5
4		STO ↑ 1 -		Go to 8
5		↑ 1 -		
6		STO x RCL 1 - D	D	Perform line 6 until $D \leq 1$
7		x^2y STO x^2y		
8		↑ ↑ ↑		
9	.035868343	x		
10	.193527818	- x		
11	.482199394	+ x		
12	.756704078	- x		
13	.918206857	+ x		
14	.897056937	- x		
15	.988205891	+ x		
16	.577191652	- x 1 + RCL		
17		x		

Factoring Integers and Determining Primes

Prime Numbers under 100

2	13	31	53	73
3	17	37	59	79
5	19	41	61	83
7	23	43	67	89
11	29	47	71	97

With the list memorized or in sight, it is easy to factor any integer x less than 10000 (and many other integers even greater). In the following program, omit the numbers 2 and 5 from the list of primes if the integer ends in 1, 3, 7 or 9.

LINE	DATA	OPERATIONS	DISPLAY	REMARKS																														
1	x	<table border="1" style="width: 100%; border-collapse: collapse; text-align: center;"> <tr><td>STO</td><td>\sqrt{x}</td><td></td><td></td><td></td></tr> <tr><td></td><td></td><td></td><td></td><td></td></tr> <tr><td></td><td></td><td></td><td></td><td></td></tr> <tr><td></td><td></td><td></td><td></td><td></td></tr> <tr><td></td><td></td><td></td><td></td><td></td></tr> <tr><td></td><td></td><td></td><td></td><td></td></tr> </table>	STO	\sqrt{x}																														If x is even, let $P = 2$; otherwise $P = 3$.
STO	\sqrt{x}																																	
2		<table border="1" style="width: 100%; border-collapse: collapse; text-align: center;"> <tr><td></td><td></td><td></td><td></td><td></td></tr> <tr><td></td><td></td><td></td><td></td><td></td></tr> <tr><td></td><td></td><td></td><td></td><td></td></tr> <tr><td></td><td></td><td></td><td></td><td></td></tr> <tr><td></td><td></td><td></td><td></td><td></td></tr> </table>																										Max						
3		<table border="1" style="width: 100%; border-collapse: collapse; text-align: center;"> <tr><td></td><td></td><td></td><td></td><td></td></tr> <tr><td></td><td></td><td></td><td></td><td></td></tr> <tr><td></td><td></td><td></td><td></td><td></td></tr> <tr><td></td><td></td><td></td><td></td><td></td></tr> <tr><td></td><td></td><td></td><td></td><td></td></tr> </table>																											If $P \geqslant \text{Max}$, stop					
4		<table border="1" style="width: 100%; border-collapse: collapse; text-align: center;"> <tr><td>RCL</td><td></td><td></td><td></td><td></td></tr> <tr><td></td><td></td><td></td><td></td><td></td></tr> <tr><td></td><td></td><td></td><td></td><td></td></tr> <tr><td></td><td></td><td></td><td></td><td></td></tr> <tr><td></td><td></td><td></td><td></td><td></td></tr> </table>	RCL																															
RCL																																		
5	P	<table border="1" style="width: 100%; border-collapse: collapse; text-align: center;"> <tr><td>\div</td><td></td><td></td><td></td><td></td></tr> <tr><td></td><td></td><td></td><td></td><td></td></tr> <tr><td></td><td></td><td></td><td></td><td></td></tr> <tr><td></td><td></td><td></td><td></td><td></td></tr> <tr><td></td><td></td><td></td><td></td><td></td></tr> </table>	\div																									Q	Read note					
\div																																		

Note: If Q is not an integer, let P = next prime number, go to 3. If Q is a prime then both P and Q are factors, stop. Otherwise P is a factor, let P = current prime, go to 2.

Example: Factor 4807.

Answer: $4807 = 11 \times 19 \times 23$

LINE	DATA	OPERATIONS	DISPLAY	REMARKS																									
1	4807	<table border="1" style="width: 100%; border-collapse: collapse; text-align: center;"> <tr><td>STO</td><td>\sqrt{x}</td><td></td><td></td><td></td></tr> <tr><td></td><td></td><td></td><td></td><td></td></tr> <tr><td></td><td></td><td></td><td></td><td></td></tr> <tr><td></td><td></td><td></td><td></td><td></td></tr> <tr><td></td><td></td><td></td><td></td><td></td></tr> </table>	STO	\sqrt{x}																								69.33253205	$\leftarrow \text{Max}, P = 3$
STO	\sqrt{x}																												
2		<table border="1" style="width: 100%; border-collapse: collapse; text-align: center;"> <tr><td>RCL</td><td>3</td><td>\div</td><td></td><td></td></tr> <tr><td></td><td></td><td></td><td></td><td></td></tr> <tr><td></td><td></td><td></td><td></td><td></td></tr> <tr><td></td><td></td><td></td><td></td><td></td></tr> <tr><td></td><td></td><td></td><td></td><td></td></tr> </table>	RCL	3	\div																							1602.333333	$P = 7$
RCL	3	\div																											
3		<table border="1" style="width: 100%; border-collapse: collapse; text-align: center;"> <tr><td>RCL</td><td>7</td><td>\div</td><td></td><td></td></tr> <tr><td></td><td></td><td></td><td></td><td></td></tr> <tr><td></td><td></td><td></td><td></td><td></td></tr> <tr><td></td><td></td><td></td><td></td><td></td></tr> <tr><td></td><td></td><td></td><td></td><td></td></tr> </table>	RCL	7	\div																							686.7142857	$P = 11$
RCL	7	\div																											
4		<table border="1" style="width: 100%; border-collapse: collapse; text-align: center;"> <tr><td>RCL</td><td>1</td><td>1</td><td>\div</td><td></td></tr> <tr><td></td><td></td><td></td><td></td><td></td></tr> <tr><td></td><td></td><td></td><td></td><td></td></tr> <tr><td></td><td></td><td></td><td></td><td></td></tr> <tr><td></td><td></td><td></td><td></td><td></td></tr> </table>	RCL	1	1	\div																						437.	$\leftarrow Q$ is an integer, so 11 is a factor
RCL	1	1	\div																										
5		<table border="1" style="width: 100%; border-collapse: collapse; text-align: center;"> <tr><td>STO</td><td>\sqrt{x}</td><td></td><td></td><td></td></tr> <tr><td></td><td></td><td></td><td></td><td></td></tr> <tr><td></td><td></td><td></td><td></td><td></td></tr> <tr><td></td><td></td><td></td><td></td><td></td></tr> <tr><td></td><td></td><td></td><td></td><td></td></tr> </table>	STO	\sqrt{x}																								20.90454496	$\leftarrow \text{Max}, P = 11$
STO	\sqrt{x}																												
6		<table border="1" style="width: 100%; border-collapse: collapse; text-align: center;"> <tr><td>RCL</td><td>1</td><td>1</td><td>\div</td><td></td></tr> <tr><td></td><td></td><td></td><td></td><td></td></tr> <tr><td></td><td></td><td></td><td></td><td></td></tr> <tr><td></td><td></td><td></td><td></td><td></td></tr> <tr><td></td><td></td><td></td><td></td><td></td></tr> </table>	RCL	1	1	\div																						39.72727273	$P = 13$
RCL	1	1	\div																										
7		<table border="1" style="width: 100%; border-collapse: collapse; text-align: center;"> <tr><td>RCL</td><td>1</td><td>3</td><td>\div</td><td></td></tr> <tr><td></td><td></td><td></td><td></td><td></td></tr> <tr><td></td><td></td><td></td><td></td><td></td></tr> <tr><td></td><td></td><td></td><td></td><td></td></tr> <tr><td></td><td></td><td></td><td></td><td></td></tr> </table>	RCL	1	3	\div																						33.61538462	$P = 17$
RCL	1	3	\div																										
8		<table border="1" style="width: 100%; border-collapse: collapse; text-align: center;"> <tr><td>RCL</td><td>1</td><td>7</td><td>\div</td><td></td></tr> <tr><td></td><td></td><td></td><td></td><td></td></tr> <tr><td></td><td></td><td></td><td></td><td></td></tr> <tr><td></td><td></td><td></td><td></td><td></td></tr> <tr><td></td><td></td><td></td><td></td><td></td></tr> </table>	RCL	1	7	\div																						25.70588235	$P = 19$
RCL	1	7	\div																										
9		<table border="1" style="width: 100%; border-collapse: collapse; text-align: center;"> <tr><td>RCL</td><td>1</td><td>9</td><td>\div</td><td></td></tr> <tr><td></td><td></td><td></td><td></td><td></td></tr> <tr><td></td><td></td><td></td><td></td><td></td></tr> <tr><td></td><td></td><td></td><td></td><td></td></tr> <tr><td></td><td></td><td></td><td></td><td></td></tr> </table>	RCL	1	9	\div																						23.	$Q = 23$ is a prime; 19 and 23 are factors
RCL	1	9	\div																										

Example: Factor 2909.

Answer: 2909 is a prime.

Fahrenheit to Centigrade

See page 55

Feet, Inches to Centimeters

See page 56

Fibonacci Numbers

Fibonacci sequence

Formula:

In a Fibonacci sequence, each term is the sum of the two preceding terms.

$$f_i = f_{i-1} + f_{i-2}$$

f_i represents the i^{th} term in the sequence.

Example:

Develop the Fibonacci sequence with $f_1 = 1$, $f_2 = 1$

Answer:

1, 1, 2, 3, 5, 8, 13, 21, ...

LINE	DATA	OPERATIONS	DISPLAY	REMARKS
1	f_1	\uparrow 		
2	f_2	 		
3		\uparrow \uparrow $R\downarrow$ $R\downarrow$ $+$	f_i	Perform line 3 for $i = 3, 4, \dots$

Finding the n^{th} Fibonacci number

Example:

Find the 12^{th} Fibonacci number in the sequence 1, 1, 2, 3, 5, 8 ...

Answer: 144

LINE	DATA	OPERATIONS	DISPLAY	REMARKS
1	n	\uparrow 5 \sqrt{x} STO \uparrow		
2		1 $+$ 2 \div x^y		
3		RCL \div EEX 9 $+$		
4		EEX 9 $-$ 		

80 Fractions

Fractions

Improper fractions to mixed numbers

Formula:

$$\frac{a}{b} = p + \frac{q}{b}$$

where $a > b$

$q < b$

Example:

$$\frac{517}{13} = 39 \frac{10}{13}$$

LINE	DATA	OPERATIONS	DISPLAY	REMARKS
1	a	\uparrow \uparrow \square \square \square		
2	b	\uparrow $R\downarrow$ \div \square \square	D	Let p = integer part of D
3		CLX \square \square \square		
4	p	$R\downarrow$ $R\downarrow$ $R\downarrow$ x $-$	q	

Reduce fraction to lowest terms

Formula:

$$\frac{a}{b} \rightarrow \frac{p}{q}$$

Note: If $p > q$, we can simplify the fraction further, see "Improper fractions to mixed numbers", above.

Example:

$$\frac{138}{391} = \frac{6}{17}$$

LINE	DATA	OPERATIONS	DISPLAY	REMARKS
1	a	STO \square \square \square \square		
2	b	\square \square \square \square		
3		\uparrow \uparrow RCL \div \square	D	Let f be the largest integer $\leq D$
4	f	$x\div y$ CLX RCL x $-$	E	If E = 0, go to 6
		\square \square \square \square		
5		RCL $x\div y$ STO CLX +		Go to 3
		\square \square \square \square		
6	a	RCL \div \square \square \square	p	
7	b	RCL \div \square \square \square	q	

Addition and subtraction of fractions

Formulas:

$$\frac{a}{b} + \frac{c}{d} = \frac{p}{q}$$

Since

$$\frac{a}{b} - \frac{c}{d} = \frac{a}{b} + \left(\frac{-c}{d} \right),$$

Only an addition routine is necessary.

Note: If the highest common factor HCF (b, d) = 1, start from Line 6 and skip the three “**RCL** **÷**” sequences in the routine.

Example 1:

$$\frac{11}{12} + \frac{9}{34} = \frac{241}{204}$$

Example 2:

$$\frac{4}{11} - \frac{5}{31} = \frac{69}{341}$$

LINE	DATA	OPERATIONS	DISPLAY	REMARKS
1	b	STO 		
2	d	 		
3		↑ ↑ RCL ÷ 	D	Let f be the largest integer $\leq D$
4	f	x²y CLX RCL x -	E	If E = 0, go to 6
		 		
5		RCL x²y STO CLX +		Go to 3
		 		
6	a	↑ 		
7	d	x RCL ÷ 		
8	c	↑ 		
9	b	x RCL ÷ + 	p	
10	b	↑ 		
11	d	x RCL ÷ 	q	

82 Gaussian Probability Function

Multiplication and division of fractions

Formulas:

$$\frac{a}{b} \times \frac{c}{d} = \frac{ac}{db} = \frac{p}{q}$$

Since

$$\frac{a}{b} \div \frac{c}{d} = \frac{a}{b} \times \frac{d}{c} ,$$

Only a multiplication routine is necessary.

Note: The result is not necessarily reduced to lowest terms, (see “Reduce fraction to lowest terms”, above).

Example:

$$\frac{3}{19} \div \frac{2}{7} = \frac{3}{19} \times \frac{7}{2} = \frac{21}{38}$$

LINE	DATA	OPERATIONS	DISPLAY	REMARKS
1	a	<input type="button" value="↑"/> <input type="button" value=" "/> <input type="button" value=" "/> <input type="button" value=" "/> <input type="button" value=" "/>		
2	c	<input type="button" value="x"/> <input type="button" value=" "/> <input type="button" value=" "/> <input type="button" value=" "/> <input type="button" value=" "/>	p	
3	b	<input type="button" value="↑"/> <input type="button" value=" "/> <input type="button" value=" "/> <input type="button" value=" "/> <input type="button" value=" "/>		
4	d	<input type="button" value="x"/> <input type="button" value=" "/> <input type="button" value=" "/> <input type="button" value=" "/> <input type="button" value=" "/>	q	

Gamma Function

See page 75

Gaussian Probability Function

Formula:

$$f(x) = \frac{1}{\sqrt{2\pi}} e^{-\frac{x^2}{2}}$$

Example:

$$f(2.23) = .03319$$

LINE	DATA	OPERATIONS	DISPLAY	REMARKS
1	x	<input type="button" value="↑"/> <input type="button" value="x"/> <input type="button" value="2"/> <input type="button" value="÷"/> <input type="button" value="CHS"/>		
2		<input type="button" value="e<sup>x</sup>"/> <input type="button" value="π"/> <input type="button" value="2"/> <input type="button" value="x"/> <input type="button" value="√x"/>		
3		<input type="button" value="÷"/> <input type="button" value=" "/> <input type="button" value=" "/> <input type="button" value=" "/> <input type="button" value=" "/>		

Geometric Progressions

See pages 114–118

Grads to Degrees

See page 15

Gudermannian Function

See page 85

Harmonic Mean

See page 139

Harmonic Numbers*Formula:*The Harmonic numbers H_i ($i = 1, 2, \dots$) are

$$1, 1 + \frac{1}{2}, 1 + \frac{1}{2} + \frac{1}{3}, 1 + \frac{1}{2} + \frac{1}{3} + \frac{1}{4}, \dots$$

or

$$1, \frac{3}{2}, \frac{11}{6}, \frac{25}{12}, \dots$$

Example:

Display the sequence in decimal form.

Answer:

$$1, 1.5, 1.833, 2.083, \dots$$

LINE	DATA	OPERATIONS	DISPLAY	REMARKS
1		CLR		
2		1 + ↑ √/x RCL		Perform lines 2–3 for $i = 1, 2, 3, \dots$
3		+ STO	H_i	

84 Highest Common Factor

n^{th} Harmonic number

Example:

Find the 7th Harmonic number

Answer:

2.59

Note: .5772156649 is Euler's constant.

LINE	DATA	OPERATIONS	DISPLAY	REMARKS
1	n	STO ↑↑↑↑		
2		1 2 0 ↑x x		
3		x 1 2 ↑x -		
4		x 2 ↑x + x		
5	.5772156649	+ RCL ln +	H _n	

Harmonic Progressions

See page 114

Haversine

See page 157

Highest Common Factor

Definition:

The highest common factor (or greatest common divisor) of two positive integers a and b is the largest integer which divides both a and b. We write it as HCF(a, b).

Example:


$$\text{HCF}(51, 119) = 17.$$

LINE	DATA	OPERATIONS	DISPLAY	REMARKS
1	a	STO		
2	b			
3		↑↑RCL ÷	D	Let f be the largest integer $\leq D$
4	f	x \neq y CLX RCL x -	E	If E = 0, go to 6
5		RCL x \neq y STO CLX +		Go to 3
6		RCL		

Hyperbolic and Inverse Hyperbolic Functions

Register usage for the following functions:

(Same as for keyboard trigonometric functions.)



Gudermannian function

Formula:

$$\text{gd } x = 2 \arctan e^x - \frac{\pi}{2}$$

Example:

$$\text{gd } 0.345 = 19.386$$

Note: $\frac{\pi}{2} = 90^\circ$

LINE	DATA	OPERATIONS	DISPLAY	REMARKS
1	x	e ^x arc tan 2 x		
2		9 0 -		

Hyperbolic sine

Formula:

$$\sinh x = \frac{e^x - e^{-x}}{2}$$

Example:

$$\sinh 3.2 = 12.25$$

LINE	DATA	OPERATIONS	DISPLAY	REMARKS
1	x	e ^x ↑ 1/x - 2		
2		÷		

86 Hyperbolic and Inverse Hyperbolic Functions

Hyperbolic cosine

Formula:

$$\cosh x = \frac{e^x + e^{-x}}{2}$$

Example:

$$\cosh 3.2 = 12.29$$

LINE	DATA	OPERATIONS					DISPLAY	REMARKS
1	x	e ^x	↑	1/x	+	2		
2		÷						

Hyperbolic tangent

Formula:

$$\tanh x = \frac{\sinh x}{\cosh x} = \sin \text{gd } x$$

Example:

$$\tanh 3.2 = .99668$$

LINE	DATA	OPERATIONS					DISPLAY	REMARKS
1	x	e ^x	arc	tan	2	x		
2		9	0	-	sin			

Hyperbolic cotangent

Formula:

$$\coth x = \frac{1}{\tanh x}$$

Example:

$$\coth 3.2 = 1.003$$

LINE	DATA	OPERATIONS					DISPLAY	REMARKS
1	x	e ^x	arc	tan	2	x		
2		9	0	-	sin	1/x		

Hyperbolic cosecant*Formula:*

$$\operatorname{csch} x = \frac{1}{\sinh x}$$

Example:

$$\operatorname{csch} 3.2 = .08166$$

LINE	DATA	OPERATIONS					DISPLAY	REMARKS
1	x	e ^x	↑	1/x	-	2		
2		÷	1/x					

Hyperbolic secant*Formula:*

$$\operatorname{sech} x = \frac{1}{\cosh x}$$

Example:

$$\operatorname{sech} 3.2 = .081$$

LINE	DATA	OPERATIONS					DISPLAY	REMARKS
1	x	e ^x	↑	1/x	+	2		
2		÷	1/x					

Inverse Gudermannian function*Formula:*

$$\operatorname{gd}^{-1} \theta = \ln \tan \left(\frac{\pi}{4} + \frac{\theta}{2} \right)$$

Example:

$$\operatorname{gd}^{-1} 30^\circ = .549$$

$$\text{Note: } \frac{\pi}{4} = 45^\circ$$

LINE	DATA	OPERATIONS					DISPLAY	REMARKS
1	θ	↑	2	÷	4	5		
2		+	tan	ln				

88 Hyperbolic and Inverse Hyperbolic Functions

Inverse hyperbolic sine

Formula:

$$\begin{aligned}\sinh^{-1} x &= \ln \left[x + (x^2 + 1)^{\frac{1}{2}} \right] \\ &= \text{gd}^{-1} (\tan^{-1} x)\end{aligned}$$

Example:

$$\sinh^{-1} 51.777 = 4.64$$

LINE	DATA	OPERATIONS	DISPLAY	REMARKS
1	x	arc tan 2 ÷ 4		
2		5 + tan ln		

Inverse hyperbolic tangent

Formula:

$$\tanh^{-1} x = \frac{1}{2} \ln \frac{1+x}{1-x} = \text{gd}^{-1} (\sin^{-1} x)$$

Example:

$$\tanh^{-1} 0.777 = 1.038$$

LINE	DATA	OPERATIONS	DISPLAY	REMARKS
1	x	arc sin 2 ÷ 4		
2		5 + tan ln		

Inverse hyperbolic secant

Formula:

$$\operatorname{sech}^{-1} x = \ln \left[\frac{1}{x} + \left(\frac{1}{x^2} - 1 \right)^{\frac{1}{2}} \right] = \text{gd}^{-1} (\cos^{-1} x)$$

Example:

$$\operatorname{sech}^{-1} 0.777 = .74$$

LINE	DATA	OPERATIONS	DISPLAY	REMARKS
1	x	arc cos 2 ÷ 4		
2		5 + tan ln		

Inverse hyperbolic cosine*Formula:*

$$\cosh^{-1} x = \operatorname{sech}^{-1} \frac{1}{x}$$

Example:

$$\cosh^{-1} 51.777 = 4.64$$

LINE	DATA	OPERATIONS	DISPLAY	REMARKS
1	x	1/x arc cos 2 ÷		
2		4 5 + tan ln		

Inverse hyperbolic cotangent*Formula:*

$$\coth^{-1} x = \tanh^{-1} \frac{1}{x}$$

Example:

$$\coth^{-1} 51.777 = .0193$$

LINE	DATA	OPERATIONS	DISPLAY	REMARKS
1	x	1/x arc sin 2 ÷		
2		4 5 + tan ln		

Inverse hyperbolic cosecant*Formula:*

$$\operatorname{csch}^{-1} x = \sinh^{-1} \frac{1}{x}$$

Example:

$$\operatorname{csch}^{-1} 0.777 = 1.0705$$

LINE	DATA	OPERATIONS	DISPLAY	REMARKS
1	x	1/x arc tan 2 ÷		
2		4 5 + tan ln		

Hyperbolic Cosecant

See page 87

Hyperbolic Cosine

See page 86

Hyperbolic Cotangent

See page 86

Hyperbolic Secant

See page 87

Hyperbolic Sine

See page 85

Hyperbolic Tangent

See page 86

Improper Fractions

See page 80

Infinite Power Series Summation

Method:

To sum an infinite power series

$$\sum_{i=1}^{\infty} a_i x^i ,$$

1. accumulate the sum in the S register,
2. keep the value of x in the T register
3. retain the previous term in the Z register, and
4. calculate the coefficient modifier using only the X and Y registers.

Example:

Evaluate the sine integral $SI(x)$ for $x = \frac{\pi}{2}$

$$SI(x) = \int_0^x \frac{\sin u}{u} du = x - \frac{x^3}{3 \cdot 3!} + \frac{x^5}{5 \cdot 5!} - \frac{x^7}{7 \cdot 7!} + \dots$$

i takes the values 1, 3, 5, ..., 13 in the algorithm

Answer:

$$SI\left(\frac{\pi}{2}\right) = 1.37$$


LINE	DATA	OPERATIONS	DISPLAY	REMARKS
1	x	↑ ↑ ↑ STO		
2		CLX + x x		Perform lines 2–7 for i = 1,
3	i	1/x		3, 5, 7, ... until D_i does
4	i + 1	x		not change.
5	i + 2	x		
6	i + 2	x CHS ÷ ↑ ↑		
7		RCL + STO	D_i	

Numerical Integration

Method:

To approximate the area A between a curve, $y = f(x)$, and the x -axis, sum the areas of the constituent trapezoids of width I . Each trapezoid has the area

$$I \times \frac{y_i + y_{i-1}}{2}$$



Example:

Find the area bounded by

$$y = x^2 + 2x - 3,$$

$$x = -2,$$

$$x = 0,$$

$$y = 0 \text{ (the } x \text{ axis).}$$


In this case

$$x_0 = -2, x_n = 0, n = 4, I = .5$$

Replace $f(x)$ by “ $\boxed{\uparrow} \boxed{\uparrow} \boxed{\uparrow} 2 \boxed{+} \boxed{\times} 3 \boxed{-}$ ”

Answer:

7.25



LINE	DATA	OPERATIONS	DISPLAY	REMARKS
1	x_n	\uparrow [] [] [] []		
2	x_0	[-] [] [] []		
3	n	[\div] [] [] []	I	
4	x_0	[] [] [] []		
5		[f(x)] [] [] []		Replace f(x) by proper keystroke(s)
		[] [] [] []		
6		[STO] [] [] []		
7	x_0	[\uparrow] [] [] []		
8	I	[+] [] [] []		
9		[f(x)] [] [] []	y_1	
10		[RCL] [+] [] []		
11	I	[x] [2] [\div] []	D	If D > 0, go to 13
		[] [] [] []		
12		[CHS] [] [] []		
13		[STO] [] [] []		
14	x_0	[\uparrow] [] [] []		Perform lines 14–21 for i = 2, ..., n
		[] [] [] []		
15	I	[\uparrow] [] [] []		
16	i	[x] [+] [] []		
17		[f(x)] [] [] []	y_i	
18	y_{i-1}	[+] [] [] []		
19	I	[x] [2] [\div] []	E_i	If $E_i > 0$, go to 21
		[] [] [] []		
20		[CHS] [] [] []		
21		[RCL] [+] [STO] []		

Interest and Discount (Simple)

Simple amount and interest

Formulas:

The simple interest I on a principal PV for n years at the annual interest rate i is given by

$$I = PV \cdot i \cdot n$$

The simple amount FV is

$$FV = PV + I = PV(1 + i n)$$

Example:

Find the simple amount and interest on \$1000 at 5% interest for 2 years ($i = 0.05$).

Answers:

$$FV = \$1100, I = \$100$$

LINE	DATA	OPERATIONS	DISPLAY	REMARKS
1	PV	<input type="button" value="↑"/> <input type="button" value="↑"/> <input type="button" value=""/> <input type="button" value=""/> <input type="button" value=""/>		
2	i	<input type="button" value="x"/> <input type="button" value=""/> <input type="button" value=""/> <input type="button" value=""/> <input type="button" value=""/>		
3	n	<input type="button" value="x"/> <input type="button" value="EEX"/> <input type="button" value="7"/> <input type="button" value="+"/> <input type="button" value="EEX"/>		
4		<input type="button" value="7"/> <input type="button" value="-"/> <input type="button" value=""/> <input type="button" value=""/> <input type="button" value=""/>	I	
5		<input type="button" value="+"/> <input type="button" value=""/> <input type="button" value=""/> <input type="button" value=""/> <input type="button" value=""/>	FV	

Simple add-on interest

The following key sequence finds the **monthly payments** and **finance charge** for an amount PV , to be paid over t years at interest rate i .

Example:

Find the monthly payments and finance charge for a loan of \$2037.01 which will be paid over 3 years at 5%. ($i = 0.05$)

Answers:

$$TP (\text{Total of payments}) = \$2342.57$$

$$F (\text{Finance charge}) = \$305.56$$

$$n (\text{Number of payments}) = 36$$

$$PMT_1 (\text{First payment}) = \$65.12$$

$$PMT (\text{Each payment except first}) = \$65.07$$

LINE	DATA	OPERATIONS	DISPLAY	REMARKS
1	PV	STO ↑ ↑		
2	t	x		
3	i	x + . 0 0		
4		5 + EEX 7 +		
5		EEX 7 -	TP	Total of payments
6		RCL x \neq y STO x \neq y -	F	Finance charge
7		RCL ↑ ↑ 1 2		
8		↑		
9	t	x	n	Number of payments
10		÷ . 0 0 5		
11		- EEX 7 + EEX		
12		7 -	PMT	Each payment except first
13	n	↑ 1 - x RCL		
14		x \neq y -	PMT ₁	First payment

Simple discount

Formula:

The **simple discount** D on a sum S for n years at the discount rate i is given by

$$D = S n i$$

and the **present value** of S is given by

$$PV = S - D = S (1 - i n).$$

Example:

Find the simple discount on a debt of \$1500 due in 9 months at a discount rate of 6%. What is the present value of the debt? ($i = 0.06$, $n = 9/12$).

Answers:

$$D \text{ (Simple discount)} = \$67.50$$

$$PV \text{ (Present value)} = \$1432.50$$

LINE	DATA	OPERATIONS	DISPLAY	REMARKS
1	S	↑ STO		
2	i	x		
3	n	x EEX 7 + EEX		
4		7 - RCL x \neq y	D	
5		-	PV	

96 Interest (Compound)

Interest (Compound)

Formula:

The formula for finding the future value FV of PV dollars invested for t years at interest i compounded q times a year is:

$$FV = PV \left(1 + \frac{i}{q}\right)^{qt}$$

Example 1.

If \$50 is deposited at 6% interest, compounded quarterly, what will be its value after 5 years? ($i = 0.06$)

Answer:

\$67.34

LINE	DATA	OPERATIONS	DISPLAY	REMARKS
1	q	↑ STO		
2	t	x		
3	i	RCL ÷ 1 \times^y		
4	PV	x EEX 7 + EEX		
5		7 -	FV	

Example 2:

Suppose \$500 is to be initially deposited in a savings account and \$100 is added to it monthly. If the account grows at a rate of 6% compounded monthly, what is the balance at the end of each month over a 1-year period? ($i = 0.06$)

LINE	DATA	OPERATIONS	DISPLAY	REMARKS
1	.06	\uparrow 1 2 \div 1		
2		+ \uparrow \uparrow \uparrow	1.005	
3	500	\times [] [] []	502.5	After 1 month
4	100	STO + \times [] []	605.5125	After 2 months
5		RCL + \times [] []	709.0400625	After 3 months
6		RCL + \times [] []	813.0852628	After 4 months
7		RCL + \times [] []	917.6506891	After 5 months
8		RCL + \times [] []	1022.738942	After 6 months
9		RCL + \times [] []	1128.352637	After 7 months
10		RCL + \times [] []	1234.4944	After 8 months
11		RCL + \times [] []	1341.166872	After 9 months
12		RCL + \times [] []	1448.372706	After 10 months
13		RCL + \times [] []	1556.11457	After 11 months
14		RCL + \times [] []	1664.395143	After 12 months

Continuous compounding

Formula:

The future value FV of PV dollars invested at interest rate i for t years, compounded continuously is:

$$FV = PV \cdot e^{it}$$

Example:

Determine the value of \$50 deposited at 6% for 5 years, compounded continuously. ($i = 0.06$)

Answer:

\$67.49

LINE	DATA	OPERATIONS	DISPLAY	REMARKS
1	i	\uparrow [] [] []		
2	t	\times e^x [] []		
3	PV	\times EEX 7 + EEX		
4		7 - [] []	FV	

Interest Rates

True interest rate

Formula:

The true interest rate (approximate) on installment purchases may be calculated by using the direct ratio formula:

$$i_t = \frac{6 m F}{3 PV (n + 1) + F (n - 1)}$$

where n = number of payments

 m = number of payments in one year

 PV = amount financed

 F = finance charge (carrying charge).

Example:

A financial corporation advertises consumer installment loans up to \$1000 at 6% (add-on) rate. If such a loan is to be paid off in 12 equal monthly payments, what is the true interest rate? ($n = m = 12$, $F = 60$, $PV = 1000$)

Answer:

10.89%

LINE	DATA	OPERATIONS	DISPLAY	REMARKS
1		6 ↑		
2	m	x		
3	F	STO x RCL		
4	n	↑ STO 1 - x		
5		RCL 1 + 3 x		
6	PV	x + ÷ EEX 2		
7		x EEX 7 + EEX		
8		7 -		

Effective rate from nominal rate

Formula:

The effective rate equivalent of the nominal rate r compounded n times per year is:

$$\text{effective rate} = \left(1 + \frac{r}{n}\right)^n - 1$$

Example:

What is the effective rate if the nominal rate is 6% compounded monthly? ($r = 0.06$)

Answer:

6.17%

LINE	DATA	OPERATIONS	DISPLAY	REMARKS
1	n	\uparrow $1/x$ \uparrow \uparrow \uparrow		
2	y	x 1 + x^y 1		
3		- EEX 2 x EEX		
4		7 + EEX 7 -		Answer is in %

Find i (given PMT, PV, and n)

Formula:

$$i = \frac{PMT \left[1 - \left(\frac{1}{1+i} \right)^n \right]}{PV}$$

Example:

A \$30,000 mortgage is paid at \$179.86 per month for 360 months. Find the interest rate. ($PV = 30,000$; $n = 360$; $PMT = 179.86$).

Answer:

.5% after 8 iterations (monthly interest rate)


LINE	DATA	OPERATIONS	DISPLAY	REMARKS
1	n	\uparrow \uparrow \uparrow \uparrow \uparrow		
2	PMT	\uparrow \uparrow \uparrow \uparrow \uparrow		
3	PV	\div STO \uparrow \uparrow \uparrow		
4		1 + $1/x$ x^y 1		Perform lines 4–5 for $k = 1, 2, \dots$
5		$x^{-2}y$ – RCL x \uparrow \uparrow	D_k	until D_k converges (to 2 nd decimal place)
6		EEX 2 x EEX 6		
7		+ EEX 6 – \uparrow \uparrow		Answer is in %

Interpolation

Formula:

If $f(x)$ is a function of x and $x_1 < x_0 < x_2$, we can approximate $f(x_0)$ by

$$f(x_0) \cong \frac{(x_2 - x_0) f(x_1) + (x_0 - x_1) f(x_2)}{x_2 - x_1}$$



Example:

Suppose a table shows

x	$f(x)$
1.2	0.30119
1.3	0.27253

Interpolate $f(x)$ to 5 decimal places for $x = 1.27$.

Answer:

.28113 (Display = .281128)

LINE	DATA	OPERATIONS					DISPLAY	REMARKS
1	x_0	\uparrow						
2	x_1	STO	-					
3	x_2	RCL	-	÷	STO	\uparrow		
4		1	RCL	-				
5	$f(x_1)$	x						
6	$f(x_2)$	RCL	x	+				

Short-cut for interpolating tables

Example:

Find $f(21.37)$

where

x	f(x)
21.3	0.4283
21.4	0.4153

Answer:

Since 21.37 is 7/10 the distance to 21.4 and 3/10 the distance to 21.3,

$$f(21.37) = 0.7 f(21.4) + 0.3 f(21.3) = .4192$$

LINE	DATA	OPERATIONS	DISPLAY	REMARKS
1	.7	<input type="button" value="↑"/> <input type="button" value=""/> <input type="button" value=""/> <input type="button" value=""/>		
2	.4153	<input type="button" value="x"/> <input type="button" value=""/> <input type="button" value=""/> <input type="button" value=""/>		
3	.3	<input type="button" value="↑"/> <input type="button" value=""/> <input type="button" value=""/> <input type="button" value=""/>		
4	.4283	<input type="button" value="x"/> <input type="button" value="+"/> <input type="button" value=""/> <input type="button" value=""/>	.4192	

Inverse Guidermannian

See page 87

Inverse Hyperbolic Functions

See pages 88–89

Iterative Solutions to Equations

See page 63

Least Common Multiple

Formula:

The least common multiple of two positive integers a and b is the smallest positive integer that both a and b can divide.

$$\text{LCM}(a, b) = \frac{a \cdot b}{\text{HCF}(a, b)}$$

Example:

$$\text{LCM}(51, 119) = 357.$$

LINE	DATA	OPERATIONS	DISPLAY	REMARKS
1	a	STO		
2	b			
3		↑ ↑ RCL ÷	D	Let f be the largest integer $\leq D$
4	f	x ^y CLX RCL x -	E	If E = 0, go to 6
5		RCL x ^y STO CLX +		Go to 3
6		RCL ↑		
7	a	↑		
8	b	x x ^y ÷		

Linear Regression

See page 144

Loan Repayments

(See also, simple add-on interest page 94)

Down payment, monthly payments, and finance charge

Formulas:

The key sequence below computes the following values:

DP = down payment ($= S \times D$)

PV = principal ($= S - DP$)

i = monthly interest rate ($= r/12$)

n = number of payments ($= 12 t$)

PMT = monthly payment

$$= PV \left[\frac{i}{1 - (1 + i)^{-n}} \right]$$

TP = total of payments (= n × PMT)

F = finance charge (= TP - PV)

T = total paid for items (= S + F)

Using the following given values:

S = sale price

D = % down payment/100

r = annual interest rate

t = number of years

Example:

Find the monthly payments to buy a tract of land priced at \$1195.00 with annual interest 6 1/4% and 10% down, the balance to be paid over a 7-year period. (r = 0.625, D = 0.1)

Answers:

Down payment = \$119.50

Amount financed = \$1075.50

Monthly payment = \$15.85

Total of payments = \$1331.40

Finance charge = \$255.90

Total paid for land = \$1450.90

LINE	DATA	OPERATIONS	DISPLAY	REMARKS
1	S	↑ ↑		
2	D	×	DP	Down payment
3		-	PV	Amount financed
4	r	↑ 1 2 ÷ STO	i	Monthly interest rate
5		x		
6	t	↑ 1 2 x	n	Number of payments
7		CHS RCL 1 + x ^y		
8		1 x ^z y - ÷ .		
9		0 0 5 + EEX		
10		7 + EEX 7 -	PMT	Monthly payment
11		1 2 ↑		
12	t	x x	TP	Total of payments
13	PV	-	F	Finance charge
14	S	+ +	T	Total paid for item

104 Loan Repayments

Monthly payments and finance charge

Objective

Find the monthly payments and finance charge if the amount financed is PV, which will be paid over t years at annual interest rate r.

Example:

Find the equal monthly payments and finance charge if the amount financed is \$2037.01, which will be paid over 3 years at the rate of 9.25%. ($r = .0925$)

Answers:

Number of payments = 36

First payment = \$65.15

Each payment except first = \$65.01

Total of payments = \$2340.50

Finance charge = \$303.49

LINE	DATA	OPERATIONS	DISPLAY	REMARKS
1	PV			
2	r			
3				
4	t	n Number of payments	n	Number of payments
5				
6				
7				
8				
9		PMT Each payment except first	PMT	Each payment except first
10				
11	t			
12				
13		TP Total of payments	TP	Total of payments
14				
15	PV	F Finance charge	F	Finance charge
16				
17	n			
18		PMT ₁ First payment	PMT ₁	First payment

Number of payments

Formula:

The number of monthly payments required to pay for an item costing PV dollars at annual rate r if payments are close to PMT dollars is:

$$n = \log_{1+i} \frac{1}{1 - \frac{PV \cdot i}{PMT}}$$

$$i = \frac{r}{12}$$

Example:

How many payments will pay off \$4000 at 9.5% annual rate, if the payments are close to \$150 per month?

Answer:

30 monthly payments.

LINE	DATA	OPERATIONS	DISPLAY	REMARKS
1		1 ↑ [] [] [] []		
2	r	↑ 1 2 ÷ STO		
3	PV	RCL x [] []		
4	PMT	÷ - 1/x ln RCL		
5		1 + ln ÷ EEX		
6		9 + EEX 9 -		

Logarithms

Logarithm of x to the base y

Formula:

$$\log_y x = \frac{\ln x}{\ln y}$$

Example:

$$\log_7 5 = .827$$

LINE	DATA	OPERATIONS	DISPLAY	REMARKS
1	x	ln <input type="text"/> <input type="text"/> <input type="text"/> <input type="text"/> <input type="text"/>		
2	y	ln \div <input type="text"/> <input type="text"/> <input type="text"/>		

Means

See page 138–139

Mils to Degrees

See page 14

Modulo: (Y Modulo X)

See page 129

Multiple

See page 102

Normal Function

See page 82

Odometer Adjustments

See page 136

Optimization of Register Use

See page 128

Percent

Percent change

Formula:

$$\% \text{ change} = \frac{V_N - V_O}{V_O} \times 100$$

where V_O = Original Value (base number)

V_N = New Value

Example:

If sales last year were \$8.6 million, and this year were \$9.3 million, what is the percent change?

Answer:

8.14%

LINE	DATA	OPERATIONS	DISPLAY	REMARKS
1	V_O	STO		
2	V_N	RCL - RCL ÷ EEX		
3		2 x EEX 6 +		
4		EEX 6 -		

Markup percent

Formula:

To make a gross profit of G%, add A% to the cost price. To find A for a given G:

$$A = \frac{100G}{100 - G}$$

Example:

To make a profit of 30%, what is the percentage of markup?

Answer:

42.86%

LINE	DATA	OPERATIONS	DISPLAY	REMARKS
1	G	↑ EEX 2 ÷ /x		
2		1 - /x EEX 2		
3		x EEX 7 + EEX		
4		7 -	A	

108 Permutations

Percent profit

Formula:

If A% is added to the cost price, the profit will be G% of the selling price.

$$G = \frac{100A}{A + 100}$$

Example:

If we add 30% to our cost price, what percent of the selling price will be the profit?

Answer:

23.08%

LINE	DATA	OPERATIONS	DISPLAY	REMARKS
1	A	↑ EEX 2 ÷ 1/x		
2		1 + 1/x EEX 2		
3		x EEX 7 + EEX		
4		7 -	G	

Permutations

Permutations of y objects taken x at a time

Formulas:

$${}_y P_x = P(y, x) = \frac{y!}{(y-x)!} = y \cdot (y-1) \cdot \dots \cdot (y-x+1)$$

Example:

$${}_7 P_5 = 2520$$

Notes:

1. ${}_y P_0 = 1$
2. ${}_y P_1 = y$
3. ${}_y P_y = y!$
4. The following routine applies for $y \geq x \geq 2$

LINE	DATA	OPERATIONS	DISPLAY	REMARKS
1	y	<input type="button" value="↑"/> <input type="button" value=" "/> <input type="button" value=" "/> <input type="button" value=" "/> <input type="button" value=" "/>		
2	i	<input type="button" value="x"/> <input type="button" value=" "/> <input type="button" value=" "/> <input type="button" value=" "/> <input type="button" value=" "/>		Perform line 2 for $i = y - x + 1$, ..., $y - 1$
		<input type="button" value=" "/>		

Plotting Curves

Objective:

The following two routines give values of $y = f(x)$ in x increments of I for values of x between $x_0 = a$, and $x = b$ where $b > a$. $f(x)$ should be replaced by appropriate sequence of keystrokes.

Examples of $f(x)$

For	Replace $f(x)$ by
$y = x^2$	<input type="button" value="↑"/> <input type="button" value="x"/>
$y = \ln \sin x$	<input type="button" value="sin"/> <input type="button" value="ln"/>
$y = 3 \sqrt[x]{x}$	<input type="button" value="↑"/> <input type="button" value="1/x"/> <input type="button" value="x<sup>z</sup>y"/> <input button"="" type="button" value="x"/>
$y = x^4 - 2x^2 + 3x - 7$	<input type="button" value="↑"/> <input type="button" value="↑"/> <input type="button" value="↑"/> <input type="button" value="x"/> 2 <input type="button" value="-"/> <input type="button" value="x"/> 3 <input type="button" value="+"/> <input type="button" value="x"/> 7 <input type="button" value="-"/>

Routine 1:

(Saves I in the stack.) When using this routine, $f(x)$ must not destroy the stack value in the Y register.

LINE	DATA	OPERATIONS	DISPLAY	REMARKS
1	I	<input type="button" value="↑"/> <input type="button" value="↑"/> <input type="button" value=" "/> <input type="button" value=" "/> <input type="button" value=" "/>		
2	x_0	<input type="button" value="↑"/> <input type="button" value=" "/> <input type="button" value=" "/> <input type="button" value=" "/> <input type="button" value=" "/>		
3		<input type="button" value="f(x)"/> <input type="button" value=" "/> <input type="button" value=" "/> <input type="button" value=" "/> <input type="button" value=" "/>	y_0	
4		<input type="button" value="CLX"/> <input type="button" value="+"/> <input type="button" value="+"/> <input type="button" value="↑"/> <input type="button" value=" "/>	x_i	Perform lines 4–5 for $i = 1$,
5		<input type="button" value="f(x)"/> <input type="button" value=" "/> <input type="button" value=" "/> <input type="button" value=" "/> <input type="button" value=" "/>	y_i	$2, \dots, k$ until $x_k = b$

110 Plotting Curves

Example:

Plot $y = \sqrt{\ln \frac{x}{2}}$ from $x = 3$ to $x = 5$ at intervals of 0.5. Replace $f(x)$ in routine 1 by

\uparrow 2 \div ln \sqrt{x}

LINE	DATA	OPERATIONS	DISPLAY	REMARKS
1	.5	\uparrow \uparrow \quad \quad \quad		
2	3	\uparrow \quad \quad \quad \quad	3.	$\leftarrow x_0$
3		\uparrow 2 \div ln \sqrt{x} \quad	.6367614216	$\leftarrow f(x_0)$
4		CLX + + \uparrow \uparrow	3.5	$\leftarrow x_1$
5		2 \div ln \sqrt{x} \quad	.7480747208	$\leftarrow f(x_1)$
6		CLX + + \uparrow \uparrow	4.	$\leftarrow x_2$
7		2 \div ln \sqrt{x} \quad	.8325546111	$\leftarrow f(x_2)$
8		CLR + + \uparrow \uparrow	4.5	$\leftarrow x_3$
9		2 \div ln \sqrt{x} \quad	.9005166385	$\leftarrow f(x_3)$
10		CLX + + \uparrow \uparrow	5.	$\leftarrow x_4$
11		2 \div ln \sqrt{x} \quad	.957230762	$\leftarrow f(x_4)$

Routine 2:

(Saves I in S register.) When using this routine, $f(x)$ must not destroy the S register.

LINE	DATA	OPERATIONS	DISPLAY	REMARKS
1	I	STO \quad \quad \quad \quad		
2	x_0	\uparrow \quad \quad \quad \quad		
3		$f(x)$ \quad \quad \quad \quad	y_0	Replace $f(x)$ by proper keystrokes
4		CLX RCL + \uparrow \quad	x_i	Perform lines 4–5 for $i = 1,$
5		$f(x)$ \quad \quad \quad \quad	y_i	$2, \dots, k,$ until $x_k = b$

Example:

Use routine 2 to plot $y = \sqrt{\ln \frac{x}{2}}$ from $x = 3$ to $x = 5$ at intervals of 1. Replace $f(x)$ by

\uparrow 2 \div ln \sqrt{x}

LINE	DATA	OPERATIONS	DISPLAY	REMARKS
1	1	STO		
2	3	\uparrow	3.	$\leftarrow x_0$
3		\uparrow 2 \div ln \sqrt{x}	.6367614216	$\leftarrow f(x_0)$
4		CLX RCL + \uparrow	4.	$\leftarrow x_1$
5		\uparrow 2 \div ln \sqrt{x}	.8325546111	$\leftarrow f(x_1)$
6		CLX RCL + \uparrow	5.	$\leftarrow x_2$
7		\uparrow 2 \div ln \sqrt{x}	.957230762	$\leftarrow f(x_2)$

Polar to Rectangular

See page 57

Polynomial Approximation

See page 76

Polynomial Evaluation

Formulas:

$$f(x) = c_0 x^n + c_1 x^{n-1} + \dots + c_{n-1} x + c_n$$

write

$$f(x_0) = (\dots (((c_0 x_0 + c_1) x_0 + c_2) x_0 + c_3) x_0 + \dots) x_0 + c_n$$

The following algorithm may even be shortened further:

If $c_0 = 1$, both the c_0 and the \times after it may be skipped.

If $c_i = 0$, both the c_i and the $+$ after it may be skipped.

If $c_i < 0$, press $-$ instead of $+$ after c_i instead of using $[CHS]$.

Example:

If $f(x) = x^5 + 5x^4 - 3x^2 - 7x + 11$, find $f(2.5)$

Answer: 267.71875

LINE	DATA	OPERATIONS	DISPLAY	REMARKS
1	x_0	\uparrow \uparrow \uparrow		
2	C_0			
3		x		Perform lines 3–4 for $i = 1, 2, \dots, n$
4	C_i	$+$		

Power Operations

See page 70

Powers x^y —where x is any number, y is a positive integer*Examples:*

$$2^4 = 16$$

$$(3.14)^3 = 30.959$$

$$(-3.14)^3 = -30.959$$

$$(-2.34)^8 = 898.932$$

LINE	DATA	OPERATIONS	DISPLAY	REMARKS
1	y	↑		
2	x			If $x > 0$, go to 5
3		CHS		If y is even, stop
4		x^{2y}		Stop
5		x^y		

Primes

See page 78

Progressions

Formulas:

Arithmetic Progression

$$a, a + d, a + 2d, \dots, a + (n - 1)d$$

Geometric Progression

$$a, ar, ar^2, \dots, ar^{n-1}$$

Harmonic Progression

$$\frac{a}{b}, \frac{a}{b+c}, \frac{a}{b+2c}, \dots, \frac{a}{b+(n-1)c}$$

n = number of terms

a = first term in arithmetic and geometric progressions

l = last term

d = difference between two successive terms in an arithmetic progression

r = ratio between two successive terms in a geometric progression

S = sum of a progression

Step through an arithmetic progression

Formula:

$$a, a + d, a + 2d, \dots, a + (n - 1)d$$

Example:

Display the progression with $a = 0$, $d = 17$.

Answer:

$$0, 17, 34, 51, 68, 85, 102, 119, \dots$$

LINE	DATA	OPERATIONS	DISPLAY	REMARKS
1	d	\uparrow \uparrow \uparrow \square \square		
2	a	\square \square \square \square		
3		$+$ \square \square \square		Perform line 3 as many times
		\square \square \square		as desired

114 Progressions

Step through a geometric progression

Formula:

$$a, ar, ar^2, \dots, ar^{n-1}$$

Example:

Step through the powers of 8.

Answers:

8, 64, 512, 4096, 32768, ...

LINE	DATA	OPERATIONS	DISPLAY	REMARKS
1	r	↑ ↑ ↑		
2	a			
3		x		Perform line 3 as many times
				as desired

Step through an harmonic progression

Formula:

$$\frac{a}{b}, \frac{a}{b+c}, \frac{a}{b+2c}, \dots, \frac{a}{b+(n-1)c}$$

Note: A harmonic progression can be obtained by multiplying the constant a by the reciprocals of the terms of the arithmetic progression $b, b + c, b + 2c, \dots, b + (n - 1)c$. In the following algorithm, x_i ($i = 1, 2, \dots$) represents the i^{th} term of the progression.

Example:

Step through the harmonic progression where $a = 1$, $b = 2$, and $c = 3$

Answers:

.5, .2, .125, .0909, ...

LINE	DATA	OPERATIONS	DISPLAY	REMARKS
1	a	STO		
2	c	↑ ↑ ↑		
3	b	↑ $1/x$ RCL x	x_1	
4		CLX + + ↑ $1/x$		Perform lines 4–5 for $i = 2, 3, \dots$
5		RCL x	x_i	

n^{th} term of arithmetic progression*Formula:*

Given the number of terms, the last term of an arithmetic progression is given by

$$n^{\text{th}} \text{ term} = a + (n - 1) d$$

Example:

Find the 25th term of the arithmetic progression with $a = 2$, $d = 3.14$.

Answer: 77.36

LINE	DATA	OPERATIONS	DISPLAY	REMARKS
1	n	\uparrow 1 - <input type="text"/> <input type="text"/>		
2	d	x <input type="text"/> <input type="text"/> <input type="text"/> <input type="text"/>		
3	a	$+$ <input type="text"/> <input type="text"/> <input type="text"/> <input type="text"/>		

 n^{th} term of geometric progression*Formula:*

Given the number of terms, the last term of a geometric progression is given by

$$n^{\text{th}} \text{ term} = ar^{n-1}$$

Example:

Find the 14th term of the geometric progression with $a = 2$, $r = 3.14$.

Answer: 5769197.692

LINE	DATA	OPERATIONS	DISPLAY	REMARKS
1	n	\uparrow 1 - <input type="text"/> <input type="text"/>		
2	r	<input type="text"/> <input type="text"/> <input type="text"/> <input type="text"/> <input type="text"/>		If $r > 0$, go to 5
		<input type="text"/> <input type="text"/> <input type="text"/> <input type="text"/> <input type="text"/>		
3		CHS x^y <input type="text"/> <input type="text"/> <input type="text"/>		If n is odd, go to 6
		<input type="text"/> <input type="text"/> <input type="text"/> <input type="text"/> <input type="text"/>		
4		CHS $x^z y$ $x^z y$ <input type="text"/> <input type="text"/>		Go to 6
		<input type="text"/> <input type="text"/> <input type="text"/> <input type="text"/> <input type="text"/>		
5		x^y <input type="text"/> <input type="text"/> <input type="text"/> <input type="text"/>		
6	a	x <input type="text"/> <input type="text"/> <input type="text"/> <input type="text"/>		

116 Progressions

Arithmetic progression sum (knowing the last term)

Formula:

Given the last term, the sum of an arithmetic progression to n terms is

$$S = \frac{n}{2} (a + l)$$

Example:

If $a = 3.5$, $l = 25$, and $n = 11$, find the sum.

Answer:

$$S = 156.75$$

LINE	DATA	OPERATIONS					DISPLAY	REMARKS
1	a	<input type="button" value="↑"/>	<input type="button" value=""/>	<input type="button" value=""/>	<input type="button" value=""/>	<input type="button" value=""/>		
2	l	<input type="button" value="+"/>	<input type="button" value=""/>	<input type="button" value=""/>	<input type="button" value=""/>	<input type="button" value=""/>		
3	n	<input type="button" value="x"/>	<input type="button" value="2"/>	<input type="button" value="÷"/>	<input type="button" value=""/>	<input type="button" value=""/>	S	

Arithmetic progression sum (knowing the difference)

Formula:

Given the first term and the difference between two successive terms, the sum of an arithmetic progression to n terms is:

$$S = na + \frac{n(n - 1)d}{2}$$

Example:

If $a = 3.5$, $n = 11$, and $d = 2.15$, find the sum of 11 terms.

Answer:

$$S = 156.75$$

LINE	DATA	OPERATIONS					DISPLAY	REMARKS
1	n	<input type="button" value="↑"/>	<input type="button" value="↑"/>	<input type="button" value="1"/>	<input type="button" value="-"/>	<input type="button" value=""/>		
2	d	<input type="button" value="x"/>	<input type="button" value=""/>	<input type="button" value=""/>	<input type="button" value=""/>	<input type="button" value=""/>		
3	a	<input type="button" value="↑"/>	<input type="button" value="2"/>	<input type="button" value="x"/>	<input type="button" value="+"/>	<input type="button" value="x"/>		
4		<input type="button" value="2"/>	<input type="button" value="÷"/>	<input type="button" value=""/>	<input type="button" value=""/>	<input type="button" value=""/>		

Sum of geometric progression ($r < 1$)

Formula:

The sum of a geometric progression to n terms with $r < 1$ is

$$S = \frac{a(1 - r^n)}{1 - r}$$

Example:

If $a = 1$, $r = -2.1$, and $n = 6$, find the sum of 6 terms.

Answer:

$$S = -27.3439$$

LINE	DATA	OPERATIONS	DISPLAY	REMARKS
1	a	↑ ↑		
2	n	↑		
3	r	STO		If $r > 0$, go to 5
				If n is even, go to 6
4		CHS x ^y CHS x ^y x ^y		Go to 7
5		x ^y		Go to 7
6		CHS x ^y		
7		x - 1 RCL -		
8		÷		

118 Progressions

Sum of geometric progression ($r > 1$)

Formula:

The sum of geometric progression to n terms with $r > 1$ is

$$S = \frac{a(r^n - 1)}{r - 1}$$

Example:

If $a = 1$, $r = 2.1$, $n = 6$, find the sum.

Answer:

$$S = 77.06$$

LINE	DATA	OPERATIONS	DISPLAY	REMARKS
1	a	[\uparrow] [\uparrow] [] [] []		
2	n	[\uparrow] [] [] [] []		
3	r	[STO] [x^y] [x] [x^{-y}] [-]		
4		[RCL] [1] [-] [\div] []		

Sum of infinite geometric progression ($-1 < r < 1$)

Formula:

$$S = \frac{a}{1 - r}$$

Example:

If $a = 2$ and $r = .5$, find the sum.

Answer:

$$S = 4$$

LINE	DATA	OPERATIONS	DISPLAY	REMARKS
1	a	[\uparrow] [1] [\uparrow] [] []		
2	r	[-] [\div] [] []		

Quadratic Equations

Formulas:

The roots of

$$Ax^2 + Bx + C = 0$$

are

$$\frac{-B \pm \sqrt{B^2 - 4AC}}{2A} = \frac{-B}{2A} \pm \sqrt{\left(\frac{-B}{2A}\right)^2 - \frac{2C}{2A}}$$

If

$$D = (B^2 - 4AC)/4A^2$$

is positive, the roots are real; otherwise they are complex, being

$$\frac{-B}{2A} \pm \frac{\sqrt{4AC - B^2}}{2A} i.$$

Example 1:

Solve $3.142958x^2 - 6.987122x + 1.001976 = 0$

Answers:

$x = 2.069$ and $x = .154$

Example 2:

Solve $-7.231844x^2 + 2.665513x - 3.169115 = 0$

Answers:

Roots are $.1843 + .6358i$ and $.1843 - .6358i$

LINE	DATA	OPERATIONS	DISPLAY	REMARKS
1	A	<input type="button" value="↑"/> <input type="button" value="+"/> <input type="button" value="STO"/> <input type="button" value=" "/> <input type="button" value=" "/>		
2	B	<input type="button" value="RCL"/> <input type="button" value="÷"/> <input type="button" value="CHS"/> <input type="button" value="↑"/> <input type="button" value="↑"/>		
3		<input type="button" value="x"/> <input type="button" value=" "/> <input type="button" value=" "/> <input type="button" value=" "/>		
4	C	<input type="button" value="↑"/> <input type="button" value="+"/> <input type="button" value="RCL"/> <input type="button" value="÷"/> <input type="button" value="-"/>	D	If D < 0, go to 7
		<input type="button" value=" "/> <input type="button" value=" "/> <input type="button" value=" "/>		
5		<input type="button" value="√x"/> <input type="button" value="STO"/> <input type="button" value="+"/> <input type="button" value=" "/>	Root 1	
6		<input type="button" value="x²y"/> <input type="button" value="RCL"/> <input type="button" value="-"/> <input type="button" value=" "/>	Root 2	Stop
		<input type="button" value=" "/> <input type="button" value=" "/> <input type="button" value=" "/>		
7		<input type="button" value="CHS"/> <input type="button" value="√x"/> <input type="button" value="x²y"/> <input type="button" value=" "/>	Real part	
8		<input type="button" value="x²y"/> <input type="button" value=" "/> <input type="button" value=" "/>	Imag. part	

Questions and Answers

See page 163

Radians to Degrees

See page 13

Random Numbers

Objective:

This routine will use a “seed” S to generate a sequence of pseudo random numbers R_i in either of two ranges -1 to 0 (if $S < 0$) or 0 to 1 (if $S > 0$). For best results, the seed must be a ten digit decimal fraction containing all digits 0 through 9 in an arbitrary order.

Example:

If $S = .2510637948$, generate a random series.

Answers:

.280850049, .144651421, .194891209, .651845061
.90350677, .20169633, .84919357, etc.

LINE	DATA	OPERATIONS	DISPLAY	REMARKS
1		2 9 ↑ ↑ ↑		
2	S			
3		x	D_i	Perform lines 3–4 for $i = 1$,
4	f_i	-	R_i	2, 3, Let f_i = integer
				part of D_i

Application

Suppose S is a positive seed, and L is any integer. The following program will simulate

- (a) tossing a coin, if $L = 2$,
- (b) throwing a dice, if $L = 6$,
- (c) picking integers between 1 and 100, if $L = 100$.

Example:

Let $S = .2510637948$, the following sequences of numbers N_i will be generated:

- (a) ($L = 2$) 1, 1, 1, 2, 2, 1, ...
- (b) ($L = 6$) 2, 1, 2, 4, 6, 2, ...
- (c) ($L = 100$) 29, 15, 20, 66, 91, 21, ...

Note: $1 \leq N_i \leq L$

LINE	DATA	OPERATIONS	DISPLAY	REMARKS
1	L	STO 2 9 ↑ ↑		
2	S	↑		
3		R↓ x	D _i	Perform lines 3–5 for i = 1,
4	f _i	- ↑ ↑ RCL x		2, 3, Let f _i = integer
				part of D _i
5		1 +	E _i	N _i = integer part of E _i

Rectangular to Polar
See page 57

Register Operations


In the following eighteen routines, x, y, z, t and s denote the contents of registers X, Y, Z, T and S, respectively.

The following techniques may be useful when writing your own key sequence routines.

1. Clear stack; retain storage




2. Delete x



LINE	DATA	OPERATIONS	DISPLAY	REMARKS
1		CLX +		


3. Delete y

(Lower that part of the stack above x)



LINE	DATA	OPERATIONS	DISPLAY	REMARKS
1		x=z CLX +		


4. Reverse the stack



LINE	DATA	OPERATIONS				DISPLAY	REMARKS
1		x ^z y	R↓	R↓	x ^z y		

5. Fetch t or roll up


(Bring t to X, keeping the other operands in the same order)



LINE	DATA	OPERATIONS				DISPLAY	REMARKS
1		R↓	R↓	R↓			

6. Fetch t to Y

(Bring t to Y, keeping the other operands in the same order)




LINE	DATA	OPERATIONS				DISPLAY	REMARKS
1		R↓	R↓	R↓	x ^z y		

124 Register Operations


7. Fetch z to X

(Bring z to X, keeping the other operands in the same order)




LINE	DATA	OPERATIONS	DISPLAY	REMARKS
1		R↓ R↓ x→y R↓		

8. Copy x into Z and T




LINE	DATA	OPERATIONS	DISPLAY	REMARKS
1		↑ ↑ R↓ R↓		

9. Copy y into Z (T is cleared)



LINE	DATA	OPERATIONS	DISPLAY	REMARKS
1		↑ ↑ - R↓		


10. Copy y into Z and T



LINE	DATA	OPERATIONS	DISPLAY	REMARKS
1		x \rightarrow y ↑ ↑ R↓ R↓		
2		R↓		

11. Copy y and x into Z and T respectively


(Copy x and y in reverse stack order, but this is the shortest way to save both x and y in the stack)



LINE	DATA	OPERATIONS	DISPLAY	REMARKS
1		↑ ↑ CLX + R↓		

12. Copy y and x into T and Z respectively

(Copy x and y in the same stack order to Z and T)




LINE	DATA	OPERATIONS	DISPLAY	REMARKS
1		x \rightarrow y ↑ ↑ CLX +		
2		R↓ x \rightarrow z		

126 Register Operations


13. Swap x and s

(Exchange x and s, t is lost)




LINE	DATA	OPERATIONS	DISPLAY	REMARKS
1		RCL x \leftrightarrow y STO CLX +		

14. Swap y and z




LINE	DATA	OPERATIONS	DISPLAY	REMARKS
1		R↓ x \leftrightarrow y R↓ R↓ R↓		

15. Swap z and t



LINE	DATA	OPERATIONS	DISPLAY	REMARKS
1		R↓ R↓ x \leftrightarrow y R↓ R↓		


16. Swap x and t



LINE	DATA	OPERATIONS	DISPLAY	REMARKS
1		R↓ R↓ R↓ x↔y R↓		

17. Swap x and z


(Reverse contents of X, Y, Z)



LINE	DATA	OPERATIONS	DISPLAY	REMARKS
1		x↔y R↓ R↓ x↔y R↓		

18. Swap y and t

(Reverse contents of Y, Z, T)



LINE	DATA	OPERATIONS	DISPLAY	REMARKS
1		R↓ x↔y R↓ R↓ x↔y		

Register Use Optimization

Illustrative routines to show the trade offs between the number of keystrokes and number of registers used.

Example 1:

Evaluate $\frac{A \times B}{B + C}$

5 keystrokes (+ data), 3 registers

LINE	DATA	OPERATIONS	DISPLAY	REMARKS
1	B	↑ ↑		
2	C	+ ÷		
3	A	x		

Divide both the numerator and denominator by B.

$$\frac{A}{1 + \frac{C}{B}}$$

6 keystrokes (+ data), 2 registers

LINE	DATA	OPERATIONS	DISPLAY	REMARKS
1	C	↑		
2	B	÷ 1 + 1/x		
3	A	x		

Example 2:

Evaluate $\frac{A}{A + \frac{B}{C + D}}$

7 keystrokes (+ data), 4 registers

LINE	DATA	OPERATIONS	DISPLAY	REMARKS
1	A	↑		
2	B	↑		
3	C	↑		
4	D	+ ÷ + ÷		

Divide both the numerator and denominator by A.

$$\frac{1}{1 + \frac{B}{A(C + D)}}$$

8 keystrokes (+ data), 2 registers

LINE	DATA	OPERATIONS	DISPLAY	REMARKS
1	C	↑ [] [] [] []		
2	D	+ [] [] [] []		
3	A	x [] $1/x$ [] []		
4	B	x [] 1 [] + [] $1/x$ []		

Remainder Function

y modulo x

Example:

$$7 \text{ modulo } 5 = 2$$

LINE	DATA	OPERATIONS	DISPLAY	REMARKS
1	y	↑ [] [] [] []		
2	x	↑ [] - R↓ STO []		
3		÷ [] [] [] []	D	Let f be the largest integer $\leq D$
4	f	$x \leftarrow y$ CLX RCL x - []		

Resistance


Formula:

The equivalent resistance of a parallel combination of resistors is

$$R_{\text{eq}} = \frac{1}{\frac{1}{R_1} + \frac{1}{R_2} + \dots + \frac{1}{R_n}}$$

Example:

Find R_{eq} .



Answer:

$$R_{\text{eq}} = 3.$$

LINE	DATA	OPERATIONS	DISPLAY	REMARKS
1	R_1	$\frac{1}{x}$ <input type="text"/> <input type="text"/> <input type="text"/> <input type="text"/>		
2	R_i	$\frac{1}{x}$ $+$ <input type="text"/> <input type="text"/> <input type="text"/>		Perform line 2 for $i = 2, \dots, n$
3		$\frac{1}{x}$ <input type="text"/> <input type="text"/> <input type="text"/> <input type="text"/>		

Roots

x^{th} root of $y (> 0)$

Example:


$$\sqrt[5]{7} = 1.47577$$

LINE	DATA	OPERATIONS	DISPLAY	REMARKS
1	x	$\frac{1}{x}$ <input type="text"/> <input type="text"/> <input type="text"/> <input type="text"/>		
2	y	x^y <input type="text"/> <input type="text"/> <input type="text"/> <input type="text"/>		

Rounding Numbers

Objective:

Round a number A to the n^{th} decimal place in the HP-35 for $n = 0, 1, \dots, 8$ (if $n = 0$, round the number to an integer).



Examples:

1. Round 2.343678 to the 2nd decimal place.
2. Round -4.3128057 to the 5th decimal place.
3. Round -1.334455 to an integer.

Answers:

1. 2.34
2. -4.31281
3. -1.

LINE	DATA	OPERATIONS	DISPLAY	REMARKS
1	A	[\uparrow] EEX [] [] []		Let $m = 9 - n$
2	m	[] [] [] [] []		If $A < 0$, go to 5
		[] [] [] [] []		
3		[+] EEX [] [] []		
4	m	[-] [] [] [] []		Stop
		[] [] [] [] []		
5		[-] EEX [] [] []		
6	m	[+] [] [] [] []		

Secant
See page 156

Sides of Triangle
See pages 147–151

Simultaneous Linear Equations

Formulas:

$$a_1x + b_1y + c_1z = d_1$$

$$a_2x + b_2y + c_2z = d_2$$

$$a_3x + b_3y + c_3z = d_3$$

Determinant solution to those simultaneous equations:

$$x = \frac{\begin{vmatrix} d_1 & b_1 & c_1 \\ d_2 & b_2 & c_2 \\ d_3 & b_3 & c_3 \end{vmatrix}}{D}$$

$$y = \frac{\begin{vmatrix} a_1 & d_1 & c_1 \\ a_2 & d_2 & c_2 \\ a_3 & d_3 & c_3 \end{vmatrix}}{D}$$

$$z = \frac{d_1 - b_1y - a_1x}{c_1}$$

where

$$D = \begin{vmatrix} a_1 & b_1 & c_1 \\ a_2 & b_2 & c_2 \\ a_3 & b_3 & c_3 \end{vmatrix}$$

$$= a_1 b_2 c_3 + a_2 b_3 c_1 + a_3 b_1 c_2 - a_3 b_2 c_1 - a_2 b_1 c_3 - a_1 b_3 c_2$$

$$\neq 0$$

Example:

Solve $\begin{cases} 3x - 2y + z = 3 \\ 9x + 3y - 5z = -2 \\ 2x - y - 2z = -15 \end{cases}$

Answers:

$$x = 2, y = 5, z = 7$$

LINE	DATA	OPERATIONS	DISPLAY	REMARKS
1	a_1	\uparrow <input type="text"/> <input type="text"/> <input type="text"/> <input type="text"/> <input type="text"/>		
2	b_2	x <input type="text"/> <input type="text"/> <input type="text"/> <input type="text"/> <input type="text"/>		
3	c_3	x <input type="text"/> <input type="text"/> <input type="text"/> <input type="text"/> <input type="text"/>		
4	b_1	\uparrow <input type="text"/> <input type="text"/> <input type="text"/> <input type="text"/> <input type="text"/>		
5	c_2	x <input type="text"/> <input type="text"/> <input type="text"/> <input type="text"/> <input type="text"/>		
6	a_3	x <input type="text"/> $+$ <input type="text"/> <input type="text"/> <input type="text"/>		
7	c_1	\uparrow <input type="text"/> <input type="text"/> <input type="text"/> <input type="text"/> <input type="text"/>		
8	a_2	x <input type="text"/> <input type="text"/> <input type="text"/> <input type="text"/> <input type="text"/>		
9	b_3	x <input type="text"/> $+$ <input type="text"/> <input type="text"/> <input type="text"/>		
10	a_3	\uparrow <input type="text"/> <input type="text"/> <input type="text"/> <input type="text"/> <input type="text"/>		
11	b_2	x <input type="text"/> <input type="text"/> <input type="text"/> <input type="text"/> <input type="text"/>		
12	c_1	x <input type="text"/> $-$ <input type="text"/> <input type="text"/> <input type="text"/>		
13	b_3	\uparrow <input type="text"/> <input type="text"/> <input type="text"/> <input type="text"/> <input type="text"/>		
14	c_2	x <input type="text"/> <input type="text"/> <input type="text"/> <input type="text"/> <input type="text"/>		
15	a_1	x <input type="text"/> $-$ <input type="text"/> <input type="text"/> <input type="text"/>		
16	c_3	\uparrow <input type="text"/> <input type="text"/> <input type="text"/> <input type="text"/> <input type="text"/>		
17	a_2	x <input type="text"/> <input type="text"/> <input type="text"/> <input type="text"/> <input type="text"/>		
18	b_1	x <input type="text"/> $-$ <input type="text"/> STO <input type="text"/> <input type="text"/>	D	
19	d_1	\uparrow <input type="text"/> <input type="text"/> <input type="text"/> <input type="text"/> <input type="text"/>		
20	b_2	x <input type="text"/> <input type="text"/> <input type="text"/> <input type="text"/> <input type="text"/>		
21	c_3	x <input type="text"/> <input type="text"/> <input type="text"/> <input type="text"/> <input type="text"/>		
22	b_1	\uparrow <input type="text"/> <input type="text"/> <input type="text"/> <input type="text"/> <input type="text"/>		
23	c_2	x <input type="text"/> <input type="text"/> <input type="text"/> <input type="text"/> <input type="text"/>		
24	d_3	x <input type="text"/> $+$ <input type="text"/> <input type="text"/> <input type="text"/>		
25	c_1	\uparrow <input type="text"/> <input type="text"/> <input type="text"/> <input type="text"/> <input type="text"/>		
26	d_2	x <input type="text"/> <input type="text"/> <input type="text"/> <input type="text"/> <input type="text"/>		
27	b_3	x <input type="text"/> $+$ <input type="text"/> <input type="text"/> <input type="text"/>		
28	d_3	\uparrow <input type="text"/> <input type="text"/> <input type="text"/> <input type="text"/> <input type="text"/>		
29	b_2	x <input type="text"/> <input type="text"/> <input type="text"/> <input type="text"/> <input type="text"/>		
30	c_1	x <input type="text"/> $-$ <input type="text"/> <input type="text"/> <input type="text"/>		
31	b_3	\uparrow <input type="text"/> <input type="text"/> <input type="text"/> <input type="text"/> <input type="text"/>		
32	c_2	x <input type="text"/> <input type="text"/> <input type="text"/> <input type="text"/> <input type="text"/>		
33	d_1	x <input type="text"/> $-$ <input type="text"/> <input type="text"/> <input type="text"/>		
34	c_3	\uparrow <input type="text"/> <input type="text"/> <input type="text"/> <input type="text"/> <input type="text"/>		
35	d_2	x <input type="text"/> <input type="text"/> <input type="text"/> <input type="text"/> <input type="text"/>		
36	b_1	x <input type="text"/> $-$ <input type="text"/> RCL \div <input type="text"/> <input type="text"/>	x	
37	a_1	\uparrow <input type="text"/> <input type="text"/> <input type="text"/> <input type="text"/> <input type="text"/>		
38	d_2	x <input type="text"/> <input type="text"/> <input type="text"/> <input type="text"/> <input type="text"/>		
39	c_3	x <input type="text"/> <input type="text"/> <input type="text"/> <input type="text"/> <input type="text"/>		

134 Sinking Fund and Amortization

Continued

LINE	DATA	OPERATIONS	DISPLAY	REMARKS
40	d_1	\uparrow <input type="text"/> <input type="text"/> <input type="text"/> <input type="text"/> <input type="text"/>		
41	c_2	x <input type="text"/> <input type="text"/> <input type="text"/> <input type="text"/> <input type="text"/>		
42	a_3	x $+$ <input type="text"/> <input type="text"/> <input type="text"/> <input type="text"/>		
43	c_1	\uparrow <input type="text"/> <input type="text"/> <input type="text"/> <input type="text"/> <input type="text"/>		
44	a_2	x <input type="text"/> <input type="text"/> <input type="text"/> <input type="text"/> <input type="text"/>		
45	d_3	x $+$ <input type="text"/> <input type="text"/> <input type="text"/> <input type="text"/>		
46	a_3	\uparrow <input type="text"/> <input type="text"/> <input type="text"/> <input type="text"/> <input type="text"/>		
47	d_2	x <input type="text"/> <input type="text"/> <input type="text"/> <input type="text"/> <input type="text"/>		
48	c_1	x $-$ <input type="text"/> <input type="text"/> <input type="text"/> <input type="text"/>		
49	d_3	\uparrow <input type="text"/> <input type="text"/> <input type="text"/> <input type="text"/> <input type="text"/>		
50	c_2	x <input type="text"/> <input type="text"/> <input type="text"/> <input type="text"/> <input type="text"/>		
51	a_1	x $-$ <input type="text"/> <input type="text"/> <input type="text"/> <input type="text"/>		
52	c_3	\uparrow <input type="text"/> <input type="text"/> <input type="text"/> <input type="text"/> <input type="text"/>		
53	a_2	x <input type="text"/> <input type="text"/> <input type="text"/> <input type="text"/> <input type="text"/>		
54	d_1	x $-$ <input type="text"/> RCL \div <input type="text"/>	y	
55	b_1	x CHS \uparrow <input type="text"/> <input type="text"/> <input type="text"/> <input type="text"/>		
56	d_1	$+$ <input type="text"/> <input type="text"/> <input type="text"/> <input type="text"/> <input type="text"/>		
57	a_1	\uparrow <input type="text"/> <input type="text"/> <input type="text"/> <input type="text"/> <input type="text"/>		
58	x	x $-$ <input type="text"/> <input type="text"/> <input type="text"/> <input type="text"/>		
59	c_1	\div <input type="text"/> <input type="text"/> <input type="text"/> <input type="text"/> <input type="text"/>	z	

Sinking Fund and Amortization

Sinking fund

Formula:

The sinking fund method of debt repayment is based on accumulating funds through periodic deposits which also earn interest. The periodic payment PMT required to accumulate a given future value FV is given by the formula

$$PMT = FV \frac{i}{(1 + i)^n - 1}$$

where i = interest rate

n = number of payments

Example:

How much money must be deposited at the end of each year in a 15 year sinking fund to accumulate \$1000 if interest is to be paid at 5% annually? ($i = 0.05$)

Answer:

\$46.34

LINE	DATA	OPERATIONS	DISPLAY	REMARKS
1	i	\uparrow STO		
2	n	\uparrow 1 RCL + x^y		
3		1 - \div		
4	FV	x EEX 7 + EEX		
5		7 -		

Amortization

To repay a loan by amortization, a payment amount is chosen that will (1) pay the current interest on the declining balance and (2) reduce the balance by whatever amount is not needed for interest.

$$PMT = PV \left[\frac{i}{1 - (1 + i)^{-n}} \right]$$

where n = number of payments

i = periodic interest rate

Example:

Find the uniform month-end payment that will repay a loan of \$1000 in 20 months at 6% annual interest. ($i = 0.06/12$)

Answer:

\$52.67

LINE	DATA	OPERATIONS	DISPLAY	REMARKS
1	n	\uparrow 1 \uparrow		
2	i	STO + x^y $1/x$ 1		
3		x^{-y} - RCL x^{-y} \div		
4	PV	x EEX 7 + EEX		
5		7 -		

Speedometer/Odometer Adjustments

Objective:

Assume that you are an automobile passenger approaching a speedometer test section with an HP-35 in your hand. You want to calculate

TS – true speed of the car

RS – what your speedometer will register at a posted speed

DS – distance traveled after a trip

RO – the reading of the odometer after a specific distance d

Notation:

a_1 = mileage read at beginning of trip

a_2 = mileage read at end of trip

b_1 = mileage read at beginning of test section (to nearest 20th of a mile, if possible)

b_2 = mileage read at end of test section

L = length of test section

s = speedometer value

p = posted speed

q = present milage

LINE	DATA	OPERATIONS	DISPLAY	REMARKS
1	b_1	CHS ↑		
2	b_2	+		
3	L	÷ STO		
4				For TS, go to 5
				For RS, go to 6
				For DS, go to 7
				For RD, go to 9
5	s	RCL ÷	TS	Go to 4
6	p	RCL x	RS	Go to 4
7	a_2	↑		
8	a_1	- RCL ÷	DS	Go to 4
9	d	RCL x		
10	q	+	RD	Go to 4

Example:

Assume the following:

$a_1 = 2185.2$	Mileage at start of trip
$b_1 = 2219.4$	Mileage at start of test section
$b_2 = 2224.15$	Mileage at end of test section
$L = 5$ miles	Length of test section

- How fast is the car actually traveling when the speedometer registers
(a) 50 mph? (b) 70 mph? (c) 30 mph?
- What will the speedometer register if the true speed is (a) 65 mph?
(b) 55 mph?
- When the odometer reads 2236.3, how many miles have you traveled
since the start of your trip.
- You see a sign saying “DOEVILLE 48”. If your odometer now reads
2241.5, what will it register when you arrive at DOEVILLE?

LINE	DATA	OPERATIONS	DISPLAY	REMARKS
1	2219.4	CHS ↑ [] [] [] []		
2	2224.15	[+] [] [] [] []		
3	5	[÷] STO [] [] [] []		
4	50	RCL [÷] [] [] [] []	52.63157895	Answer for 1a
5	70	RCL [÷] [] [] [] []	73.68421053	Answer for 1b
6	30	RCL [÷] [] [] [] []	31.57894737	Answer for 1c
7	65	RCL [x] [] [] [] []	61.75	Answer for 2a
8	55	RCL [x] [] [] [] []	52.25	Answer for 2b
9	2236.3	[↑] [] [] [] []		
10	2185.2	[−] RCL [÷] [] [] []	53.78947368	Answer for 3
11	48	RCL [x] [] [] [] []		
12	2241.5	[+] [] [] [] []	2287.1	Answer for 4

Square of x

$$f(x) = x^2$$

Example:

$$(2.359)^2 = 5.564881$$

LINE	DATA	OPERATIONS	DISPLAY	REMARKS
1	x	[↑] [x] [] [] []		

Stack Operations

See page 122

Standard Deviation

See page 141

Statistics**Arithmetic mean***Formula:*

The arithmetic mean (average) of a set of numbers

$$\{a_1, a_2, \dots, a_n\}$$

is

$$A = \frac{1}{n} \sum_{i=1}^n a_i$$

Example:

Compute the arithmetic mean of

$$2, 3.4, 3.41, 7, 11, 23$$

Answer:

$$8.3$$

LINE	DATA	OPERATIONS	DISPLAY	REMARKS
1	a_1	\uparrow <input type="text"/> <input type="text"/> <input type="text"/> <input type="text"/>		
2	a_i	$+$ <input type="text"/> <input type="text"/> <input type="text"/> <input type="text"/>		Perform line 2 for $i = 2, \dots, n$
3	n	\div <input type="text"/> <input type="text"/> <input type="text"/> <input type="text"/>		

Geometric mean*Formula:*

The geometric mean of a set of numbers

$$\{a_1, a_2, \dots, a_n\}$$

is

$$G = \sqrt[n]{a_1 \cdot a_2 \cdot \dots \cdot a_n}$$

Example:

Compute the geometric mean of

2, 3.4, 3.41, 7, 11, 23

Answer:

5.87

LINE	DATA	OPERATIONS	DISPLAY	REMARKS
1	a_1	\uparrow <input type="text"/> <input type="text"/> <input type="text"/> <input type="text"/> <input type="text"/>		
2	a_i	x <input type="text"/> <input type="text"/> <input type="text"/> <input type="text"/> <input type="text"/>		Perform line 2 for $i = 2, 3, \dots, n$
3	n	$1/x$ $x^{2/y}$ x^y <input type="text"/> <input type="text"/>		

Harmonic mean

Formula:

The harmonic mean of a set of numbers

$$\{a_1, a_2, \dots, a_n\}$$

is

$$H = \frac{n}{\sum_{i=1}^n \frac{1}{a_i}}$$

Example:

Find the harmonic mean of

2, 3.4, 3.41, 7, 11, 23

Answer:

4.3968

LINE	DATA	OPERATIONS	DISPLAY	REMARKS
1	a_1	$1/x$ <input type="text"/> <input type="text"/> <input type="text"/> <input type="text"/> <input type="text"/>		
2	a_i	$1/x$ $+$ <input type="text"/> <input type="text"/> <input type="text"/> <input type="text"/> <input type="text"/>		Perform line 2 for $i = 2, 3, \dots, n$
3	n	$1/x$ x $1/x$ <input type="text"/> <input type="text"/>		

140 Statistics

Average and count simultaneously

Constraints

1. Averaged values must be positive integers.
2. Number of values may not exceed 999.
3. Sum of values may not exceed 10 million.

This technique is advantageous in that entries are counted as they are added. In the following routine, X_{L3} refers to the rightmost three digits visible in the display.

LINE	DATA	OPERATIONS	DISPLAY	REMARKS
1		CLR		
2	n_i	0 0 1 +		Perform line 2 for $i = 1, \dots, n$
3	X_{L3}	\div 1 - EEX 3		
4		\div		

Example:

Assume that someone is calling out the following series of integers, and you don't know how many there will be, compute the average.

37, 16, 43, 18, 29, 11, 43, 2, 61, 73, 54, 19, 28

Answer:

33.38461538

LINE	DATA	OPERATIONS	DISPLAY	REMARKS
1		CLR		
2	37001	+		
3	16001	+		
4	43001	+		
5	18001	+		
6	29001	+		
7	11001	+		
8	43001	+		
9	2001	+		
10	61001	+		
11	73001	+		
12	54001	+		
13	19001	+		
14	28001	+	434013	
15	13	\div 1 - EEX 3		
16		\div	33.38461538	

Summing two columns simultaneously

Example:

Sum the following two columns simultaneously.

x_i	y_i
14	37.2
3	190.15
438	16.4
93	.005
1736	41.81
19	11.13

Answer: SUM x = 2303 SUM y = 296.695

LINE	DATA	OPERATIONS					DISPLAY	REMARKS
1	x_1	R↓	R↓	R↓				
2	y_1	R↓	R↓					
3	x_i	+	R↓	R↓				Perform lines 3–4 for i = 2, 3, ..., n
4	y_i	+	R↓	R↓				
5		$x \bar{z} y$	CLX	+			SUM _x	
6		$x \bar{z} y$					SUM _y	

Mean, standard deviation, and sums (ungrouped data)

Formulas:

Mean:

$$\bar{x} = \frac{\sum_{i=1}^n x_i}{n}$$

Standard deviation:

$$S = \sqrt{\frac{\sum_{i=1}^n x_i^2 - \frac{\left(\sum_{i=1}^n x_i\right)^2}{n}}{n-1}}$$

Sum of values:

$$Sx = \sum_{i=1}^n x_i$$

142 Statistics

Sum of squared values:

$$SS = \sum_{i=1}^n x_i^2$$

Example:

Compute \bar{x} , S_n , Sx , SS for the set of numbers $\{2, 3.4, 3.41, 7, 11, 23\}$, we have $Sx = 49.81$, $SS = 726.1881$, $\bar{x} = 8.3$, $S = 7.9$.

Note: This routine can be used to find Sx and SS only.

LINE	DATA	OPERATIONS	DISPLAY	REMARKS
1		CLR		
2	x_i	STO + $x \rightarrow y$ RCL \uparrow		Perform lines 2–3 for $i = 1,$
				$2, \dots, n$
3		x + $x \rightarrow y$		
4			Sx	
5		$x \rightarrow y$	SS	Stop if \bar{x} and S are not needed
6		$x \rightarrow y$ \uparrow \uparrow		
7	n	STO \div	\bar{x}	
8		R \downarrow \uparrow x RCL \div		
9		– RCL 1 – \div		
10		\sqrt{x}	S	

Mean, standard deviation, and sums (grouped data)

Formulas:

Given a set of data points

$$x_1, x_2, \dots, x_n$$

with respective frequencies

$$f_1, f_2, \dots, f_n$$

$$\text{Let } k = \sum_{i=1}^n f_i.$$

Then

$$\bar{x} = \frac{\sum_{i=1}^n f_i x_i}{\sum_{i=1}^n f_i}$$

$$S = \sqrt{\frac{\sum_{i=1}^n f_i x_i^2 - \left(\sum_{i=1}^n f_i x_i\right)^2 / k}{k - 1}}$$

$$Sx = \sum_{i=1}^n f_i x_i$$

$$SS = \sum_{i=1}^n f_i x_i^2$$

Example:

Compute mean, standard deviation, and sums for

f_i	3	3	1	2	1
x_i	1	2	3	4	5

Answer:

$$k = 10$$

$$Sx = 25$$

$$SS = 81$$

$$\bar{x} = 2.5$$

$$S = 1.43$$

LINE	DATA	OPERATIONS	DISPLAY	REMARKS
1		CLR		
2	f_i	\uparrow		Perform lines 2–5 for $i = 1, \dots, n$
3	x_i	STO x \uparrow $R\downarrow$ $+$		
4		$R\downarrow$ $R\downarrow$ $R\downarrow$ RCL x		
5		$R\downarrow$ $R\downarrow$ $R\downarrow$ $+$ $x\bar{z}y$		
6			Sx	
7		$x\bar{z}y$	SS	Stop if \bar{x} and S are not needed
8		$x\bar{z}y$ \uparrow \uparrow		
9	k	STO \div	\bar{x}	
10		$R\downarrow$ \uparrow x RCL \div		
11		$-$ RCL 1 $-$ \div		
12		\sqrt{x}	S	

Cross product sum

Formula:

$$\sum_{i=1}^n x_i y_i$$

Example:

Find the cross product sums for the following:

x_i	1	2	3	4	5
y_i	2	3.4	3.41	7	11

Answer:

$$\sum_{i=1}^5 x_i y_i = 102.03$$

LINE	DATA	OPERATIONS	DISPLAY	REMARKS
1	x_i	\uparrow		
2	y_i	x		
3	x_i	\uparrow		Perform lines 3–4 for $i = 2, 3, \dots, n$
4	y_i	$x +$		

Linear regression and correlation coefficient

Objective:

- For a set of data points

$$\{(x_i, y_i), i = 1, \dots, n\}$$

fit a straight line to them having minimum least squares deviation.

- Find the correlation of x and y.

Formulas:

$$y = ax + b$$

$$a = \frac{\sum x_i y_i - \frac{\sum x_i \sum y_i}{n}}{\sum x_i^2 - \frac{(\sum x_i)^2}{n}}$$

$$b = \bar{y} - a\bar{x}$$

where

$$\bar{x} = \frac{\sum x_i}{n}$$

$$\bar{y} = \frac{\sum y_i}{n}$$

Coefficient of determination

$$r^2 = \frac{\left[\sum x_i y_i - \frac{\sum x_i \sum y_i}{n} \right]^2}{\left[\sum x_i^2 - \frac{(\sum x_i)^2}{n} \right] \left[\sum y_i^2 - \frac{(\sum y_i)^2}{n} \right]}$$

Correlation coefficient

$$r = \sqrt{r^2} \text{ (using the sign of } a)$$

Example:

x_i	26	30	44	50	62	68	74
y_i	92	85	78	81	54	51	40

Answers:

$$Sx = 354$$

$$SSx = 19956$$

$$Sy = 481$$

$$SSy = 35451$$

$$Sxy = 22200$$

$$y = -1.03x + 121.04 = ax + b$$

$$r^2 = 0.92$$

$$r = -0.957$$

146 Statistics

LINE	DATA	OPERATIONS	DISPLAY	REMARKS
1		CLR		
2	x_i	STO + x^2y RCL \uparrow		Perform lines 2–3 for $i = 1, 2, \dots, n$
3		x + x^2y		
4			Sx	
5		x^2y	SSx	
6		CLR		
7	y_i	STO + x^2y RCL \uparrow		Perform lines 7–8 for $i = 1, 2, \dots, n$
8		x + x^2y		
9			Sy	
10		x^2y	SSy	
11		CLR		
12	x_i	\uparrow		Perform lines 12–13 for $i = 1, 2, \dots, n$
13	y_i	x +		
14			Sxy	
15	Sx	STO \uparrow		
16	Sy	x		
17	n	\div – \uparrow \uparrow \uparrow		
18	SSx	\uparrow RCL \uparrow x		
19	n	\div – \div \uparrow STO	a	
20	Sy	x^2y		
21	Sx	x –		
22	n	\div	b	Stop if r^2 is not needed
23		R \downarrow RCL x		
24	SSy	\uparrow		
25	Sy	\uparrow x		
26	n	\div – \div	r^2	
27		\sqrt{x}	r	If $a > 0$, stop
28		CHS	r	

Stirling's Approximation

See page 75

Sum Two Columns Simultaneously


See page 141

Triangles (Oblique)**Law of sines**

$$\frac{a}{\sin A} = \frac{b}{\sin B} = \frac{c}{\sin C}$$

Law of cosines

$$c^2 = a^2 + b^2 - 2ab \cos C$$

*Notes:*

- When **[arc sin]**, **[arc cos]**, **[arc tan]** are used in programs, secondary values may be required instead of the principal values.
- A . B . C are in degrees

148 Triangles (Oblique)

Given a, b, C, find A, B, c.

Example:

$$\text{Given } a = 132$$


$$b = 224$$

$$C = 28^\circ 40'$$

$$= 28.66666667^\circ$$

Find A, B, and c

Answer:




$$c = 125.35$$

$$A = 30.34^\circ$$

$$B = 120.99^\circ \text{ (secondary value)}$$

LINE	DATA	OPERATIONS	DISPLAY	REMARKS
1	a	\uparrow \uparrow x \square		
2	b	\uparrow STO x $+$ $x\bar{x}y$		
3		2 x RCL x \square		
4	C	STO cos x $-$ \sqrt{x}	c	
5		\uparrow \uparrow \uparrow RCL sin		
6		STO \uparrow \square		
7	a	x $x\bar{x}y$ \div arc sin	A	
8		R↓ RCL \square		
9	b	x $x\bar{x}y$ \div arc sin	B	

Given a, b, c; find A, B, C



Example:

$$\text{Given } a = 30.3$$

$$b = 40.4$$

$$c = 62.6$$

then find A, B, and C.

Answer:

$$A = 23.66^\circ$$

$$B = 32.35^\circ$$

$$C = 123.99^\circ$$

LINE	DATA	OPERATIONS	DISPLAY	REMARKS
1	a	$\uparrow \quad \uparrow \quad x \quad \quad \quad$		
2	b	$\uparrow \quad STO \quad x \quad + \quad$		
3	c	$\uparrow \quad x \quad - \quad x \neq y \quad RCL$		
4		$x \quad 2 \quad x \quad \div \quad arc$		
5		$cos \quad \quad \quad \quad \quad$	C	
6	a	$\uparrow \quad \uparrow \quad x \quad \quad \quad$		
7	c	$\uparrow \quad STO \quad x \quad + \quad$		
8	b	$\uparrow \quad x \quad - \quad x \neq y \quad RCL$		
9		$x \quad 2 \quad x \quad \div \quad arc$		
10		$cos \quad \uparrow \quad \quad \quad \quad$	B	
11	C	$+ \quad 1 \quad 8 \quad 0 \quad x \neq y$		
12		$- \quad \quad \quad \quad \quad$	A	

Given a, A, C; find B, b, c.

Example:

$$\text{Given } a = 17.5$$

$$A = 41^\circ 14' = 41.23333333^\circ$$

$$C = 62^\circ 12' = 62.2^\circ$$


Find B, b, and c

Answer:

$$B = 76.57^\circ$$

$$b = 25.82$$

$$c = 23.486$$



LINE	DATA	OPERATIONS	DISPLAY	REMARKS
1	C	$\uparrow \quad \uparrow \quad \quad \quad \quad$		
2	A	$STO \quad + \quad 1 \quad 8 \quad 0$		
3		$x \neq y \quad - \quad \uparrow \quad \quad \quad$	B	
4		$R \downarrow \quad sin \quad \quad \quad \quad$		
5	a	$x \quad RCL \quad sin \quad STO \quad \div$	b	
6		$R \downarrow \quad sin \quad \quad \quad \quad$		
7	a	$x \quad RCL \quad \div \quad \quad \quad$	c	

150 Triangles (Oblique)

Given a, B, C; find A, b, c

Example:


$$\text{Given } a = 25.2$$

$$B = 35^\circ 20' = 35.333333333^\circ$$

$$C = 68^\circ 30' = 68.5^\circ$$

Find A, b, and c

Answer:



$$A = 76.17^\circ$$

$$b = 15.009$$

$$c = 24.1469$$

LINE	DATA	OPERATIONS	DISPLAY	REMARKS
1	C	[↑] [↑] [] [] []		
2	B	[+] [1] [8] [0] [x \div y]		
3		[-] [STO] [] [] []	A	
4		[R↓] [sin] [] [] []		
5	a	[x] [RCL] [sin] [STO] [\div]	c	
6	B	[sin] [] [] [] []		
7	a	[x] [RCL] [\div] [] []	b	

Given B, b, c; find a, A, C.

Example:

$$\text{Given } B = 33^\circ 40' = 33.66666667^\circ$$

$$b = 31.5$$

$$c = 51.8$$


Since B is acute and $b < c$, there are two solutions. Find a, A, and C.

Answer:

$$a = 56.06$$

$$A = 80.6^\circ$$

$$C = 65.729^\circ$$




Alternate Answer:

$$a_1 = 30.16$$

$$A_1 = 32.06^\circ$$

$$C_1 = 114.27^\circ$$



LINE	DATA	OPERATIONS	DISPLAY	REMARKS
1	c	\uparrow [] [] [] []		
2	B	sin [x] [] [] []		
3	b	\div [] [] [] []	D	If D > 1, no solution, stop
		[] [] [] []		
4		arc sin STO \uparrow []	C	
5	B	+ 1 8 0 x \leftrightarrow y []		
6		- [] [] [] []	A	
7		sin [] [] [] []		
8	c	x RCL sin \div []	a	If b \geq c, stop
		[] [] [] []		
9		1 8 0 RCL - []		b < c, there are two solutions.
10		STO \uparrow [] [] []	C ₁	
11	B	+ 1 8 0 x \leftrightarrow y []		
12		- [] [] [] []	A ₁	
13		sin [] [] [] []		
14	c	x RCL sin \div []	a ₁	

Given a, b, and c; find the area.

Formulas:

Given the lengths of a triangle's three sides a, b, and c

$$\text{area} = \sqrt{S(S - a)(S - b)(S - c)}$$

where

$$S = \frac{1}{2}(a + b + c)$$

152 Triangles (Oblique)

Example:


Given $a = 5.317361553$

$b = 7.089815404$

$c = 8.862269255$

Find the area.

Answer:



Area = 18.85 ($S = 10.63$)

LINE	DATA	OPERATIONS	DISPLAY	REMARKS
1	a	[↑] [] [] [] [] []		
2	b	[↑] [↑] [] [] []		
3	c	[STO] [+] [R↓] [x ² y] [R↓]		
4		[R↓] [+] [2] [÷] []	S	
5		[R↓] [R↓] [R↓] [CLX] [RCL]		
6		[R↓] [STO] [x ² y] [−] [x ² y]		
7		[RCL] [x ² y] [−] [x] [x ² y]		
8		[RCL] [x ² y] [−] [x] [RCL]		
9		[x] [√x] [] [] []	Area	

Given a, b, and C; find area

Formula:

$$\text{Area} = \frac{1}{2} a b \sin C$$

Example:

Given $a = 5.3174$


$b = 7.0898$

$C = 45^\circ$

Find the area.

Answer:

Area = 13.33.



LINE	DATA	OPERATIONS	DISPLAY	REMARKS
1	a	t x 2 ÷		
2	b			
3	c	sin x x x x x		

Given a, B, and C; find the area

Formula:

$$\text{Area} = \frac{a^2}{2} \cdot \frac{\sin B \sin C}{\sin(B+C)}$$


Example:

$$\text{Given } a = 14.625$$

$$B = 70.54^\circ$$

$$C = 62.96^\circ$$

Find the area.




Answer:

$$\text{Area} = 123.8$$

LINE	DATA	OPERATIONS	DISPLAY	REMARKS
1	B	t t x x x x		
2	C	STO + sin x \leftrightarrow y sin		
3		RCL sin x x \leftrightarrow y ÷		
4	a	t x x 2 ÷		

Trigonometric Functions

Register usage for trigonometric functions



Secondary value of arc sin x

Let p = principal value

q = secondary value

Example:

Find secondary value for $\text{arc sin}(-.7660444436)$

Answer:

230°

LINE	DATA	OPERATIONS	DISPLAY	REMARKS
1	x	arc sin [] [] []	p	
2		1 8 0 $x \leftrightarrow y$ -	q	

Secondary value of arc cos x

Let p = principal value

q = secondary value

Example:

Find the secondary value of $\text{arc cos } (.76)$.

Answer:

319.46°

LINE	DATA	OPERATIONS	DISPLAY	REMARKS
1	x	arc cos [] [] [] []	p	
2		3 6 0 $x^{2,y}$ -	q	

Secondary value of $\text{arc tan } x$

Let p = principal value

q = secondary value

Example:

Find the secondary value of $\text{arc tan } (2)$.

Answer:

243.43°

LINE	DATA	OPERATIONS	DISPLAY	REMARKS
1	x	arc tan [] [] [] []	p	
2		1 8 0 + []	q	

Cotangent

Formula:

$$\cot x = \frac{1}{\tan x}$$

Example:

$$\cot 37^\circ = 1.327$$

LINE	DATA	OPERATIONS	DISPLAY	REMARKS
1	x	tan $1/x$ [] [] []		

156 Trigonometric Functions

Cosecant

Formula:

$$\csc x = \frac{1}{\sin x}$$

Example:

$$\csc 30^\circ = 2$$

LINE	DATA	OPERATIONS	DISPLAY	REMARKS
1	x	sin 1/x		

Secant

Formula:

$$\sec x = \frac{1}{\cos x}$$

Example:

$$\sec 45^\circ = 1.41$$

LINE	DATA	OPERATIONS	DISPLAY	REMARKS
1	x	cos 1/x		

Versine

Formula:

$$\text{vers } x = 1 - \cos x$$

Example:

$$\text{vers } 38^\circ = .21$$

LINE	DATA	OPERATIONS	DISPLAY	REMARKS
1	x	cos 1 xz-y -		

Coversine*Formula:*

$$\text{covers } x = 1 - \sin x$$

Example:

$$\text{covers } 38^\circ = .38$$

LINE	DATA	OPERATIONS				DISPLAY	REMARKS
1	x	sin	1	x^2y	-		

Exsecant*Formula:*

$$\text{exsec } x = \sec x - 1 = \frac{1}{\cos x} - 1$$

Example:

$$\text{exsec } 52^\circ = .624$$

LINE	DATA	OPERATIONS				DISPLAY	REMARKS
1	x	cos	$1/x$	1	-		

Haversine*Formula:*

$$\text{hav } x = \frac{1 - \cos x}{2}$$

Example:

$$\text{hav } 42.3^\circ = .13$$

LINE	DATA	OPERATIONS				DISPLAY	REMARKS
1	x	cos	1	x^2y	-	2	
2		\div					

Arc cotangent*Formula:*

$$\text{arc cot } x = \text{arc tan } \frac{1}{x}$$

Example:

$$\text{arc cot } 0.35 = 70.71^\circ$$

LINE	DATA	OPERATIONS				DISPLAY	REMARKS
1	x	$1/x$	arc	tan			

158 T-Statistic (Paired)

Arc cosecant

Formula:

$$\text{arc csc } x = \text{arc sin } \frac{1}{x}$$

Example:

$$\text{arc csc } 3.45 = 16.849^\circ$$

LINE	DATA	OPERATIONS			DISPLAY	REMARKS
1	x	1/x	arc	sin		

Arc secant

Formula:

$$\text{arc sec } x = \text{arc cos } \frac{1}{x}$$

Example:

$$\text{arc sec } 1.1547 = 30^\circ$$

LINE	DATA	OPERATIONS			DISPLAY	REMARKS
1	x	1/x	arc	cos		

T-Statistic (Paired)

Formulas:

Given a set of paired observations:

x_i	x_1	x_2	x_3	...	x_n
y_i	y_1	y_2	y_3	...	y_n

$$D_i = x_i - y_i$$

$$\bar{D} = \frac{1}{n} \sum_{i=1}^n D_i$$

$$S_D = \sqrt{\frac{\sum D_i^2 - \frac{(\sum D_i)^2}{n}}{n-1}}$$

$$S_{\bar{D}} = \frac{S_D}{\sqrt{n}}$$

and test statistic

$$t = \frac{\bar{D}}{S_{\bar{D}}}$$

Example:

Compute t for the following:

x _i	14	17.5	17	17.5	15.4
y _i	17	20.7	21.6	20.9	17.2

Answer:

$$t = -7.1554$$

LINE	DATA	OPERATIONS	DISPLAY	REMARKS
1		CLR		
2	x _i	↑		Perform lines 2–4 for i = 1, 2, ..., n
3	y _i	- STO + x ² y RCL		
4		↑ x + x ² y		
5		↑ ↑		
6	n	÷ STO	̄D	
7		R↓ ↑ x		
8	n	↑ R↓ ÷ - R↓		
9		R↓ 1 - x ÷		
10		√x	S _D	
11		RCL x ² y ÷	t	

Vector Cross Product

Formula:

If $\vec{x} = (x_1, x_2, x_3)$ and $\vec{y} = (y_1, y_2, y_3)$ are two vectors, then cross product \vec{z} is also a vector.

$$\vec{z} = \vec{x} \times \vec{y}$$

$$\begin{aligned}
 &= (x_2 y_3 - x_3 y_2, x_3 y_1 - x_1 y_3, x_1 y_2 - x_2 y_1) \\
 &= (z_1, z_2, z_3)
 \end{aligned}$$

Example:

If $\vec{x} = (2.34, 5.17, 7.43)$
 $\vec{y} = (.072, .231, .409)$

Find $\vec{x} \times \vec{y}$

Answer:

$$\vec{x} \times \vec{y} = (.3982, -.4221, .1683)$$

LINE	DATA	OPERATIONS	DISPLAY	REMARKS
1	x_2	\uparrow 		
2	y_3	STO x 		
3	y_2	\uparrow 		
4	x_3	\uparrow R↓ x - 	z_1	
5		R↓ 		
6	y_1	x 		
7		RCL 		
8	x_1	STO x - 	z_2	
9		RCL 		
10	y_2	x 		
11	x_2	\uparrow 		
12	y_1	x - 	z_3	

Vector Dot Product

Formulas:

Given two vectors \vec{x} , \vec{y} in an n-dimensional vector space

$$\vec{x} = (x_1, x_2, \dots, x_n)$$

$$\vec{y} = (y_1, y_2, \dots, y_n)$$

the dot product is

$$\vec{x} \cdot \vec{y} = x_1 y_1 + x_2 y_2 + \dots + x_n y_n$$

Example:

If $\vec{x} = (2.34, 5.17, 7.43, 9.11, 11.41)$

$$\vec{y} = (.072, .231, .409, .703, .891)$$

then $\vec{x} \cdot \vec{y} = 20.97226$

LINE	DATA	OPERATIONS	DISPLAY	REMARKS
1	x_1	\uparrow <input type="text"/> <input type="text"/> <input type="text"/> <input type="text"/> <input type="text"/>		
2	y_1	\times <input type="text"/> <input type="text"/> <input type="text"/> <input type="text"/> <input type="text"/>		
3	x_i	\uparrow <input type="text"/> <input type="text"/> <input type="text"/> <input type="text"/> <input type="text"/>		Perform lines 3–4 for $i = 2, 3, \dots, n$
4	y_i	\times <input type="text"/> $+$ <input type="text"/> <input type="text"/> <input type="text"/>		

Versine

See page 156

Weekday

DAY OF THE WEEK FOR ANY DATE (Since September 14, 1752)

d = day of month

m = month, with January and February being the 13th and 14th months of the previous year.

y = year (4 digits)

162 Weekday

Output is read as follows:

- 0 – Saturday
- 1 – Sunday
- 2 – Monday
- 3 – Tuesday
- 4 – Wednesday
- 5 – Thursday
- 6 – Friday

Example:

On what day was February 29, 1972?

Answer:

Tuesday (d = 29, m = 14, y = 1971)

LINE	DATA	OPERATIONS	DISPLAY	REMARKS
1	d	↑		
2	m	↑		
3	y	STO R↓ 1 + 1		
4		3 x 5 ÷	E ₁	Let e ₁ = integer part of E ₁
5		CLX		
6	e ₁	+ RCL x ² y STO x ² y		
7		↑ ↑ ↑ 5 x		
8		4 ÷	E ₂	Let e ₂ = integer part of E ₂
9		CLX		
10	e ₂	RCL +		For 20 th century date, go to 17
11		STO 1 0 0 ÷	E ₃	Let e ₃ = integer part of E ₃
12		CLX		
13	e ₃	CHS RCL + STO 4		
14		0 0 ÷	E ₄	Let e ₄ = integer part of E ₄
15		CLX		
16	e ₄	RCL +		Go to 18
17		6 +		
18		↑ ↑ 7 STO ÷	E ₅	Let e ₅ = integer part of E ₅
19		CLX		
20	e ₅	RCL x -		

Appendix

Questions you may have wanted answered about your HP-35.

- 1. Question:** What are the maximum and minimum light displays on the HP-35?

Answer: Maximum: .8.88888888888 after all decimal points light up on low battery power!

Minimum: .

- 2. Question:** A googol is 10^{100} . We can key this in as 10×10^{99} on the HP-35, right? Its reciprocal can be keyed in as $.1 \times 10^{-99}$, right?

Answer:

Wrong. 10 **EEX** 99 → 9.999999999 99
.1 **EEX** **CHS** 99 → 0.

- 3. Question:** The display is blanked out during computations. Is the keyboard also locked out?

Answer: Yes

- 4. Question:** What is the longest response on the HP-35?

Answer: $\tan x$ where $x = 9.99999999999999999$ (about 2 seconds).

- 5. Question:** Do:

1 **↑** 2 **↑** 3 **↑** **CLx**

Now, if we press **÷** or **x^y** (that is $\frac{3}{0}$ or 0^3) a flashing zero will be displayed. Will the stack be dropped?

Answer: Yes, if we divide!:

÷ **R↓** → 2. **R↓** → 1. **R↓** → 1.

No, if we do a x^y !:

x^y **R↓** → 3. **R↓** → 2. **R↓** → 1.

164 Appendix

6. Question: Pressing which key will reset the arc prefix?

Answer: Any key except **[arc]**, **[sin]**, **[cos]**, or **[tan]**.

7. Question: Do: 6 **[EE]** 16. How would you make the whole number negative?

Answer: **[x_y]** **[x_y]** **[CHS]**.

8. Question: When is **[ENTER ↑]** desirable before entering data, but **NOT** just after a **[CLX]**, **[STO]**, or **[ENTER ↑]**?

Answer: Sometimes after negating a result by **[CHS]**:
 $- \sqrt{5} + 6$ is 5 **[√x]** **[CHS]** **[↑]** 6 **[+]**

9. Question: How can three successive **[tan]** presses be used to calculate the tangent of 30° (accurate to about 7 decimal places)?

Answer: 30 **[arc]** **[tan]** **[tan]** **[tan]** $\rightarrow .5773502648$

10. Question: When may it be desirable to press **[CLX]** if the display is already zero?

Answer: When we get zero as the result of an operation, but don't want to raise it in the stack upon entering data.

11. Question: When may it be desirable to enter 0 if x is already zero?

Answer: If an algorithm requires the integer part of a number (already in the X register) such as .4, we would do: **[CLX]** 0, which would insure that this zero would be raised in the stack if the algorithm requires it.

12. **Question:** Do: $\pi \sin \pi \cos \div \pi \tan -$
 $\rightarrow .0548861507$
 $\rightarrow .0548861507$
 $\rightarrow 1. -11; \text{why not } 0?$

Answer: See problem 20.

13. **Question:** The calculator gives overflow for the tangent of 90° , right?

Answer: Wrong, but nearly so:

$$90 \tan \rightarrow 9.99999999\ 99$$

We can still add to it:

$$\text{EEEX} \ 90 \ + \rightarrow 9.999999991\ 99$$

14. **Question:** Enter any three digits in the form $d.dd \times 10^{-7}$:

$$d \ . \ d \ d \ \text{EEEX} \ \text{CHS} \ 7.$$

How do you add 1 to this number with one keystroke?
 Your number now has the form: 1.000000ddd, how do you subtract 1 from it with one keystroke?

Answer: e^x and \ln

(All this means that $\ln(1 + x)$ approaches x as x approaches 0.)

15. **Question:** Display $.00026 \times 10^{-2}$ in correct scientific notation.

Answer: .00026 $\text{EEEX} \ \text{CHS} \ 2 \ x\bar{z}y \ x\bar{z}y \rightarrow 2.6 -06$

16. **Question:** If we do “5 $\text{EEEX} \ \text{arc} \ 8$ ”, does 8 appear as an exponent or does it push up the stack?

Answer: Exponent.

17. **Question:** For which numbers will the HP-35 **not** give reciprocals?

Answer: 0. and any number with an exponent of +99 having a value other than ± 1 .

166 Appendix

18. **Question:** How can you insure an integer result for x^y , if x and y both contain integers?

Answer: x^y [EEEX] 9 [+][EEEX] 9 [-]

19. **Question:** X contains 26.56505118. You wish to get tan x, but accidentally you hit the [arc] key. How can you cancel the [arc] prefix, and get tan x?

Answer: $x_{\bar{z}}y$ $x_{\bar{z}}y$ [tan] → .4999999999

20. **Question:** Numbers which start with .0d, where d is any digit other than zero, are displayed with only 9 significant digits, although ten significant digits are used in calculations. (This includes numbers .01 through .0999999999). The number is not rounded to 9 significant digits. Can you give some examples to show this fact?

Answer: Example 1:

23 [$\frac{1}{x}$] → .0434782608 10 [x] → .4347826087

Example 2:

1.000000001 [EEEX] [CHS] 2 → 1.000000001 -02
 $x_{\bar{z}}y$ $x_{\bar{z}}y$ → .01 10 [x] → .1000000001

Example 3:

[EEEX] 9 [\uparrow] 9 [+][EEEX] 11 [\div] [STO] .01
[$\frac{1}{x}$] → 100 [RCL] → .01 [$\frac{1}{x}$] → 99.9999991

Example 4:

1.000000001 [EEEX] [CHS] 2 [\uparrow] 23 [$\frac{1}{x}$] [+]
→ .0534782608 10 [\div] → .5347826088 -03

21. **Question:** Compute $e^\pi - \pi^e$.

Answer: [π] [e^x] 1 [e^x] [π] [x^y] [-] → .68153493

22. Question: **EEX CHS** 8 **STO** 30

RCL **-** **↑** → 29.99999999 **sin** → .4999999998
↓ **RCL** **+** **↑** → 30. **sin** → .5
↓ **RCL** **+** **↑** → 30.00000001 **sin** → .4999999998
↓ **RCL** **+** → 30.00000002 **sin** → .5000000002

Answer: Can be explained by the accuracy statement on page 21 of the operating manual.

23. Question: 2050667999 **arc** **tan** → 89.99999995

EEX CHS 8 **+** → 89.99999996 **tan**

→ 861719914.7

Answer: Can be explained by the accuracy statement on page 21 of the operating manual, an extreme “end case”.

24. Question: For values:

89.9999997 to 90.	tan	→ 9.99999999	99
89.99999992 to 89.99999996	tan	→ 861719914.7	
89.99999987 to 89.99999991	tan	→ 457788704.7	
89.99999980 to 89.99999986	tan	→ 311685926.6	
89.99999975 to 89.99999979	tan	→ 240151451.6	

Answer: Can be explained by the accuracy statement on page 21 of the operating manual, an extreme “end case”.

25. Question: How many decimal points light up on low battery power?

Answer: Only 14—the point following the true decimal point does not come on!

26. Question: **EEX** 9 **↑** .5 **+** .5 **-** → 1000000001. Why?

Answer: **EEX** 9 or 10000000000 **+** .5 = 1000000000.5 which is rounded to 1000000001, and 1000000001 **-** .5 = 1000000000.5 which is again rounded to 1000000001.

27. **Question:** How would you display all 1's, 2's, ... 9's?

- Answer:**
- 1's: 9 e^x e^x 9 EEX 88 \div
 - 2's: 9 e^x e^x 4.5 EEX 77 \div
 - 3's: 9 e^x e^x 3 EEX 66 \div
 - 4's: 9 e^x e^x 2.25 EEX 55 \div
 - 5's: 9 e^x e^x 1.8 EEX 44 \div
 - 6's: 9 e^x e^x 1.5 EEX 33 \div
 - 7's: 7.777777777 EEX 77
 - 8's: 9 e^x e^x 1.125 EEX 11 \div
 - 9's: 9 e^x e^x

28. **Question:** To divide the displayed value by 2: \uparrow 2 \div . What other three keystrokes will do the same?

- Answer:** e^x \sqrt{x} \ln , for $|x| \leq 227$.

29. **Question:** CHS 9 \uparrow 8 \div EEX 97 \times $1/x \rightarrow -8.888888889-98$

Change this number to $-8.888888888-98$ in a shorter number of keystrokes than keying in the new value.

Note: we must add 1×10^{-107} to the number in the display. If we try to add $.000000001 \times 10^{-98}$, we also get into trouble.

- Answer:** EEX 98 STO \times EEX CHS 9 + RCL \div

30. **Question:** How would you display the number e (2.718281828):

- in the shortest number of keystrokes? (2 keystrokes)
- as the limit of $\left(1 + \frac{1}{n}\right)^n$ as n gets large? (7 keystrokes)
- as the function of a digit? (5 keystrokes)
- using the π key, but not e^x
or x^y ? (13 keystrokes)
- as a continued fraction? (21 keystrokes)

- Answer:**
- (a) 1 e^x or if $x = 0$: e^x e^x
 - (b) $\text{EE} \times$ 6 \uparrow $1/x$ 1 $+$ x^y
 - (c) 5 $1/x$ 5 e^x x^y
 - (d) π \uparrow \ln \div \uparrow \ln \div \uparrow \ln \div \uparrow \ln \div
 - (e) 18 $1/x$ 14 $+$ $1/x$ 10 $+$ $1/x$ 6 $+$ $1/x$ 1 $+$
 $1/x$ 2 \times 1 $+$

31. Question: Can you display all ten L.E.D. digits in an orderly sequence? (5 keystrokes):

Answer: 8 \uparrow 81 \div $\rightarrow .0987654321$

32. Question: Determine the Golden Ratio ϕ to 11 significant digits, given $\phi = (\sqrt{5} + 1)/2$ and $\phi - \frac{1}{\phi} = 1$.

Answer: 5 \sqrt{x} 1 $+$ 2 \div $\rightarrow 1.618033989 \uparrow$
 $1/x \rightarrow .6180339887 \boxed{-} \rightarrow 1.$
 $\phi = 1.6180339887$

Index

- Amortization 135
Angle Conversions 11
Angles of Triangle 147–151
Annuities 15
Antilog of x (common) 16
Appendix 163
Arc Cosecant 158
Arc Cosine, Secondary 154
Arc Cotangent 157
Arc Secant 158
Arc Sine, Secondary 154
Arc Tangent, Secondary 155
Area by Integration 92
Area of Triangle 151–153
Arithmetic Mean 138
Arithmetic Progressions 113, 115–116
Average 138, 140–142
Azimuth to Bearing 12

Base Conversion 17
Bearing to Azimuth 12
Bernoulli Numbers 21
Binomial Coefficient 23

Centigrade to Fahrenheit 55
Centimeters to Feet, Inches 56
Chi-square Evaluation 22
Combinations 23
Complex Number Operations 24
Complex Trigonometric and Hyperbolic Functions 36–54
Compound Interest 96
Conversions 55
Coordinate Conversions 57
Correlation Coefficient 144
Cosecant 156
Cotangent 155
Coversine 157
Cross Product 160
Cross Product Sum 144

Decimal Degrees to Degrees, Minutes, Seconds 11
Degrees, Minutes, Seconds to Decimal Degrees 11
Degrees to Grads 15
Degrees to Mils 14
Degrees to Radians 14
Discount 95
Dot Product 161
Double Precision Product 59

Equation Solving (iterative techniques) 63–68
Euler Numbers 69
Exponentiation 70–74

Factorial 75
Factorial and Gamma Functions 75–77
Factoring Integers and Determining Primes 78
Fahrenheit to Centigrade 55
Feet, Inches to Centimeters 56
Fibonacci Numbers 79
Fractions 80

Gamma Function 75
Gaussian Probability Function 82
Geometric Progressions 113–118
Grads to Degrees 15
Gudermannian Function 85

Harmonic Mean 139
Harmonic Numbers 83
Harmonic Progressions 114
Haversine 157
Highest Common Factor 84
Hyperbolic Functions 85

Improper Fractions 80
Infinite Power Series Summation 90
Integration, Numerical 92
Interest and Discount (simple) 94
Interest (compound) 96
Interest Rates 98
Interpolation 100
Inverse Gudermannian 87
Inverse Hyperbolic Functions 88–89
Iterative Solutions to Equations 63–68

Least Common Multiple 102
Linear Regression 144
Loan Repayments 94, 102–105
Logarithms 106

Markup Percent 107
Means 138–139
Mils to Degrees 14
Modulo: (Y Modulo X) 129
Multiple 102

Normal Function 82

Odometer Adjustments 136

Percent 107
Permutations 108
Plotting Curves 109–111
Polar to Rectangular 57
Polynomial Approximation 76
Polynomial Evaluation 111
Power Operations 70
Power Series 90
Powers 112
Primes 78
Progressions 113–118
Profit Percent 108

Quadratic Equations 119
Questions and Answers 163–169

Radians to Degrees 13
Random Numbers 120
Rectangular to Polar 57
Register Operations 122–127
Register Use Optimization 128
Remainder Function 129
Resistance 130
Root of Polynomial 68
Roots 130
Rounding Numbers 131

Secant 156
Sides of Triangle 147–151
Simultaneous Linear Equations 132
Sine Integral, example 91
Sinking Fund and Amortization 134
Speedometer/Odometer Adjustments 136
Square of X 137
Stack Operations 122–127
Standard Deviation 141
Statistics 138–146
Stirling's Approximation 75
Sum Two Columns Simultaneously 141

Triangles (oblique) 147–153
Trigonometric Functions 154–158
T-Statistic (paired) 158

Vector Cross Product 160
Vector Dot Product 161
Versine 156

Weekday 161



Sales, service and support in 172 centers in 65 countries
ADVANCED PRODUCTS DIVISION
10900 Wolfe Road, Cupertino, CA 95014
Offices in principal cities throughout the United States