## HEWLETT-PACKARD

# HP-41C

## STAT PAC



## NOTICE

Hewlett-Packard Company makes no express or implied warranty with regard to the keystroke procedures and program material offered or their merchantability or their fitness for any particular purpose. The keystroke procedures and program material are made available solely on an "as is" basis, and the entire risk as to their quality and performance is with the user. Should the keystroke procedures or program material prove defective, the user (and not Hewlett-Packard Company nor any other party) shall bear the entire cost of all necessary correction and all incidental or consequential damages. Hewlett-Packard Company shall not be liable for any incidental or consequential damages in connection with or arising out of the furnishing, use, or performance of the keystroke procedures or program material.

## INTRODUCTION

The programs in the Stat Pac have been drawn from the fields of general statistics, analysis of variance, regression, test statistics, and distribution functions.

Each program in this pac is represented by one program in the Application Module and a section in this manual. The manual provides a description of the program with relevant equations, a set of instructions for using the program, and one or more example problems, each of which includes a list of the keystrokes required for its solution.

Before plugging in your Application Module, turn your calculator off, and be sure you understand the section "Inserting and Removing Application Modules." Before using a particular program, take a few minutes to read "Format of User Instructions" and "A Word About Program Usage."

You should first familiarize yourself with a program by running it once or twice while following the complete User Instructions in the manual. Thereafter, the program's prompting should provide the necessary instructions, including which variables are to be input, which keys are to be pressed, and which values will be output. A quick-reference card with a brief description of each program's operating instructions has been provided for your convenience.

We hope the Stat Pac will assist you in the solution of numerous problems in your discipline. If you have technical problems with this Pac, refer to your HP-41 owner's handbook for information on Hewlett-Packard "technical support" or "programming assistance."

**Note:** Application modules are designed to be used in all HP-41 model calculators. The term "HP-41C" is used throughout the rest of this manual, unless otherwise specified, to refer to all HP-41 calculators.

## CONTENTS

Introduction	
Contents	2
Inserting and Removing Application Modules	ļ
Format of User Instructions	,
A Word About Program Usage	3

## Programs

## **General Statistics**

Basic Statistics for Two Variables	10
Moments, Skewness and Kurtosis (For Grouped or Ungrouped Data) This program provides geometric descriptions (symmetry, relative peakness and flatness) of a distribution.	<b>14</b>

## **Analysis of Variance**

Analysis of Variance (One Way) This program is used to test the observed differences among k sample means.	.18
Analysis of Variance (Two Way, No Replications) The row effects and the column effects are tested independently in the analysis of the total variability of a set of data.	.22
Analysis of Covariance (One Way) This program tests the effect of one variable separately from the effect of the second variable.	.26
Curve Fitting	
Curve Fitting	.32
Multiple Linear Regression Linear regression for two or three independent variables, using least squares method.	.38
Polynomial Regression This program fits a set of data points to either a cubic or a parabolic	.44

polynomial, using least squares method.

## **Test Statistics**

t Statistics
Chi-Square Evaluation
Contingency Table
Spearman's Rank Correlation Coefficient
Distribution Functions
Normal and Inverse Normal Distribution

Chi-Square Distribution
This program evaluates the chi-square density. A series approxi-
mation is used to evaluate the cumulative distribution.

Appendix A: Program Data	 
<b>Appendix B: Program Labels</b>	 <b>74</b>

## **INSERTING AND REMOVING APPLICATION MODULES**

Before you insert an application module for the first time, familiarize yourself with the following information.

Up to four application modules can be plugged into the ports on the HP-41C. While plugged in, the names of all programs contained in the module can be displayed by pressing **CATALOG** 2.

#### CAUTION

Always turn the HP-41C off before inserting or removing any plug-in extensions or accessories. Failure to turn the HP-41C off could damage both the calculator and the accessory.

Here is how you should insert application modules:

1. Turn the HP-41C off! Failure to turn the calculator off could damage both the module and the calculator.



2. Remove the port covers. Remember to save the port covers, they should be inserted into the empty ports when no extensions are inserted.



3. With the application module label facing downward as shown, insert the application module into any port **after** the last memory module presently inserted.



- 4. If you have additional application modules to insert, place them into any port after the last memory module. For example, if you have a memory module inserted in port 1, you can insert application modules in any of ports 2, 3, or 4. Never insert an application module into a lower numbered port than a memory module. Be sure to place port covers over unused ports.
- 5. Turn the calculator on and follow the instructions given in this book for the desired application functions.

To remove application modules:

- 1. Turn the HP-41C off! Failure to do so could damage both the calculator and the module.
- 2. Grasp the desired module handle and pull it out as shown.



3. Place a port cap into the empty port.

#### **Mixing Memory Modules and Application Modules**

Any time you wish to insert other extensions (such as the HP-82104A Card Reader, or the HP-82143A Printer) the HP-41C has been designed so that the memory modules are in lower numbered ports.

So, when you are using both memory modules and application modules, the memory modules must always be inserted into the lower numbered ports and the application module into any port after the last memory module. When mixing memory and application modules, the HP-41C allows you to leave gaps in the port sequence. For example, you can plug a memory module into port 1 and an application module into port 4, leaving ports 2 and 3 empty.

## FORMAT OF USER INSTRUCTIONS

The completed User Instruction Form—which accompanies each program is your guide to operating the programs in this Pac.

The form is composed of five labeled columns. Reading from left to right, the first column, labeled STEP, gives the instruction step number.

The INSTRUCTIONS column gives instructions and comments concerning the operations to be performed.

The INPUT column specifies the input data, the units of data if applicable, or the appropriate alpha response to a prompted question. Data Input keys consists of 0 to 9 and the decimal point (the numeric keys), EEX (enter exponent), and CHS (change sign).

The FUNCTION column specifies the keys to be pressed after keying in the corresponding input data.

The DISPLAY column specifies prompts, intermediate and final answers and their units, where applicable.

Above the DISPLAY column is a box which specifies the minimum number of registers necessary to execute the program. Refer to pages 73 and 117 in the Owner's Handbook for a complete description of how to size calculator memory.

## A WORD ABOUT PROGRAM USAGE

#### Catalog

When an Application Module is plugged into a port of the HP-41C, the contents of the Module can be reviewed by pressing CATALOG 2 (the Extension Catalog). Executing the CATALOG function lists the name of each global label in the module, as well as functions of any other extensions which might be plugged in. Remember that the catalog function lists the extension in port 1 first, followed by the extensions in ports 2-4.

#### **ALPHA and USER Mode Notation**

This manual uses a special notation to signify ALPHA mode. Whenever a statement on the User Instruction Form is printed in gold, the ALPHA key must be pressed before the statement can be keyed in. After the statement is input, press ALPHA again to return the calculator to its normal operating mode, or to begin program execution. For example,  $XEO \ SBSTAT$  means press the following keys:  $XEO \ ALPHA \ SBSTAT \ ALPHA$ .

In USER mode, when referring to the top two rows of keys (the keys have been re-defined), this manual will use the symbols (A, C, E, A) and (R/S) on the User Instruction Form and in the keystroke solutions to sample problems.

#### **Using Optional Printer**

When the optional printer is plugged into the HP-41C along with this Applications Module, all results will be printed automatically. You may also want to keep a permanent record of the values input to a certain program. A convenient way to do this is to set the Print Mode switch to NORMAL before running the program. In this mode, all input values and the corresponding keystrokes will be listed on the printer, thus providing a record of the entire operation of the program.

#### **Downloading Module Programs**

If you wish to trace execution, to modify, to record on magnetic cards, or to print a program in this Application Module, it must first be copied into the HP-41C's program memory. For information concerning the HP-41C COPY function, see the Owner's Handbook. It is *not* necessary to copy a program in order to run it.

## **Program Interruption**

These programs have been designed to operate properly when run from beginning to end, without turning the calculator off (remember, the calculator may turn itself off). If the HP-41C is turned off, it may be necessary to set flag 21 (SF 21) to continue proper execution.

## Use of Labels

You should generally avoid writing programs into the calculator memory that use program labels identical to those in your Application Module. In case of a label conflict, the label within program memory has priority over the label within the Application Pac program. All program labels used in this Pac are listed in appendix B, "Program Labels."

## **Key Assignments**

If you have customized your keyboard with the ASN function, those reassignments will take precedence over the local labels A, C, and E used in this Pac.

## Flag 03

If flag 03 is set when a Stat Pac program is executed, the statistical registers may not be cleared and incorrect results may occur.

## **BASIC STATISTICS FOR TWO VARIABLES**

This program calculates means, standard deviations, covariance, correlation coefficient, coefficients of variation, sums of data points, sum of multiplication of data points, and sums of squares of data points derived from a set of ungrouped data points  $\{(x_i, y_i), i = 1, 2, ..., n\}$ , or grouped data points  $\{(x_i, y_i), i = 1, 2, ..., n\}$ , or grouped data points  $\{(x_i, y_i), i = 1, 2, ..., n\}$ .

$$\begin{array}{rcl} \mbox{means } \bar{x} = \frac{1}{n} \sum_{i=1}^{n} x_i & \overline{y} = \frac{1}{n} \sum_{i=1}^{n} y_i \\ \mbox{standard deviations } s_x = & \sqrt{\frac{\sum x_i^2 - n \overline{x}^2}{n - 1}} & (\text{of sample}) \\ \mbox{or } s_x' = & \sqrt{\frac{\sum x_i^2 - n \overline{x}^2}{n}} & (\text{of population}) \\ \mbox{s}_y = & \sqrt{\frac{\sum y_i^2 - n \overline{y}^2}{n - 1}} & (\text{of sample}) \\ \mbox{or } s_y' = & \sqrt{\frac{\sum y_i^2 - n \overline{y}^2}{n}} & (\text{of population}) \\ \mbox{covariance} & s_{xy} = \frac{1}{n - 1} \left( \sum x_i y_i - \frac{1}{n} \sum x_i \sum y_i \right) & (\text{of sample}) \\ \mbox{or } s_{xy'}' = \frac{1}{n} \left[ \sum x_i y_i - \frac{1}{n} \sum x_i \sum y_i \right] & (\text{of population}) \\ \mbox{correlation coefficient } \gamma_{xy} = \frac{s_{xy}}{s_x s_y} \end{array}$$

Coefficients of variation 
$$V_x = \frac{s_x}{\overline{x}} \cdot 100$$
,  $V_y = \frac{s_y}{\overline{y}} \cdot 100$ 

Note n is a positive integer and n > 1.

r.

				<b>SIZE:</b> 012
STEP	INSTRUCTIONS	INPUT	FUNCTION	DISPLAY
	Ungrouped Data			
1.	Initialize the program.		xeq ΣBSTAT	ΣBSTAT
2.	Repeat step 2~3 for i=1,2,,n. Input: $x_i$ $y_i$	X <sub>i</sub> Yi	ENTER•)	(i)
3.	If you made a mistake in inputting $x_{\rm k}$ and $y_{\rm k},$ then correct by	Х <sub>к</sub> Ук	ENTER•) C	(k-1)
4.	Go to step 8 for basic statistic calculations.			
	Grouped Data			
5.	Initialize the program.		xeq ΣBSTG	ΣBSTG
6. 7	Repeat step 6~7 for i=1,2,,n. Input: $x_i$ $y_i$ $f_i$ If you made a mistake in inputting	X <sub>i</sub> Y <sub>i</sub> f <sub>i</sub>	ENTER+ ENTER+ A	$(\Sigma f_i)$
7.	$x_k$ , $y_k$ , and $f_k$ , then correct by	X <sub>k</sub> Y <sub>k</sub> f <sub>k</sub>	ENTER+) ENTER+) C	$(\Sigma f_i - f_k)$
8.	To calculate basic statistics: $\overline{x}$ $\overline{y}$ $s_x$ $s_x'$ $s_y'$ $s_y'$ $v_x^*$ $v_y^*$ $s_{xy}$ $s_{xy}'$ $y_x'$ $y_x'$ $\sum x_i$ $\sum y_i$ $\sum x_i^2$ $\sum y_i^2$		E R/S R/S R/S R/S R/S R/S R/S R/S R/S R/S	$\begin{array}{l} XBAR = (\bar{x}) \\ YBAR = (\bar{y}) \\ SX = (S_x) \\ SX = (s'_x) \\ SY = (s_y) \\ SY = (s_y) \\ SY = (s'_y) \\ VX = (v_x) \\ VY = (v_y) \\ SXY = (S_{xy}) \\ SXY = (S_{xy}) \\ SXY = (\Sigma_{x}) \\ SXY = (\Sigma_{x}) \\ \SigmaY = (\Sigma_{x}) \\ \SigmaY = (\Sigma_{x}) \\ \SigmaY = (\Sigma_{x}) \\ \SigmaY2 = (\Sigma_{x})^{2} \\ \SigmaY2 = (\Sigma_{y})^{2} \end{array}$
9.	Repeat step 8 if you want the results again. To use the same program for another set of data, initialize the program by $\rightarrow$ then go to step 2 or step 6.		<b>a</b>	ΣBSTAT or ΣBSTG
11.	To use the other program, go to step 1 or step 5.			

**NOTE:** "DATA ERROR" will be displayed if  $\bar{x}$  or  $\bar{y}$  is zero. Press **Press** and proceed.

## Example 1:

For the following set of data, find the means, standard deviations, covariance, correlation coefficient, coefficients of variation, and the sums.

x <sub>i</sub>	26	30	44	50	62	68	74	
y <sub>i</sub>	92	85	78	81	54	51	40	

Voustralias	Diamlary
Reystrokes:	Display:
XEQ ALPHA SIZE ALPHA 012	
XEQ ALPHA SBSTAT ALPHA	Σ <b>BSTAT</b>
26 ENTER+) 92 A	
100 ENTER+ 100 A	
100 ENTER+ 100 C	
30 ENTER+) 85 A	
44 [ENTER+] 78 A	
50 ENTER+) 81 A	
62 [ENTER+] 54 A	
68 ENTER+ 51 A	
74 ENTER+) 40 A	7.00
E	XBAR = 50.57
R/S	YBAR=68.71
R/S	SX=18.50
R/S	SX.=17.13
(R/S)	SY=20.00
	SY.=18.51
R/S	VX=36.58
R/S	VY=29.10
	SXY=-354.14
(R/S)	SXY.=-303.55
R/S	GXY=-0.96
R/S	ΣX=354.00
R/S	ΣY=481.00
	ΣXY=22200.00
	ΣX2=19956.00
R/S	ΣY2=35451.00

## Example 2:

Apply the program to the following set of grouped data.

Xi	4.8	5.2	3.8	4.4	4.1	
Уi	15.1	11.5	14.3	13.6	12.8	
f <sub>i</sub>	1	3	1	6	2	

Keystrokes:	Display:
XEQ ALPHA SIZE ALPHA 012	
XEQ ALPHA SBSTG ALPHA	Σ <b>BSTG</b>
4.8 ENTER+ 15.1 ENTER+ 1 A	
5.2 ENTER+) 11.5 ENTER+) 3 A	
3.8 ENTER+ 14.3 ENTER+ 1 A	
4.4 ENTER+ 13.6 ENTER+ 6 A	
4.1 ENTER+ 12.8 ENTER+ 2 A	13.00
Ε	XBAR = 4.52
R/S	YBAR=13.16
R/S	SX=0.45
R/S	SX.=0.43
R/S	SY=1.11
R/S	SY.=1.07
R/S	VX=9.93
R/S	VY=8.42
R/S	SXY=-0.31
R/S	SXY.=-0.28
R/S	GXY=-0.62
R/S	Σ <b>X</b> = <b>58.80</b>
R/S	Σ <b>Y</b> =171.10
R/S	Σ <b>XY</b> =770.22
R/S	$\Sigma X2 = 268.38$
R/S	Σ <b>Y2=2266.69</b>

## MOMENTS, SKEWNESS AND KURTOSIS (FOR GROUPED OR UNGROUPED DATA)

For grouped or ungrouped data, moments are used to describe sets of data, skewness is used to measure the lack of symmetry in a distribution, and kurtosis is the relative peakness or flatness of a distribution. For a given set of data

$$\{x_1, x_2, ..., x_n\}$$
:

$$1^{\text{st}} \text{ moment} \qquad \bar{\mathbf{x}} = \frac{1}{n} \sum_{i=1}^{n} \mathbf{x}_i$$

$$2^{nd} \text{ moment} \quad m_2 = \frac{1}{n} \sum x_i^2 - \bar{x}^2$$

- 3<sup>rd</sup> moment  $m_3 = \frac{1}{n} \sum x_i^3 \frac{3}{n} \bar{x} \sum x_i^2 + 2 \bar{x}^3$
- 4<sup>th</sup> moment  $m_4 = \frac{1}{n} \sum x_i^4 \frac{4}{n} \overline{x} \sum x_i^3 + \frac{6}{n} \overline{x}^2 \sum x_i^2 3\overline{x}^4$

Moment coefficient of skewness

$$\gamma_1 = \frac{m_3}{m_2^{3/2}}$$

Moment coefficient of kurtosis

$$\gamma_2 = \frac{m_4}{m_2^2}$$

This program also provides the option for calculating those statistics for grouped data (using similar formulas as for ungrouped data):

data	X <sub>1</sub>	X <sub>2</sub>	 Xm
frequency	f <sub>1</sub>	f2	 f <sub>m</sub>

Note that for this case, 1<sup>st</sup> moment

$$\bar{\mathbf{x}} = \frac{\sum_{i=1}^{m} \mathbf{f}_{i} \mathbf{x}_{i}}{\sum_{i=1}^{m} \mathbf{f}_{i}}$$

## **Reference:**

Theory and Problems of Statistics, M.R. Spiegel, Schaum's Outline, McGraw-Hill, 1961

				<b>SIZE</b> : 012
STEP	INSTRUCTIONS	INPUT	FUNCTION	DISPLAY
1.	<b>Ungrouped Data</b> Initialize the program.		(χεο) ΣMMTUG	ΣMMTUG
2.	Repeat step $2 \sim 3$ for $i=1,2,\ldots,n$ . Input $x_i$ .	Xi	A	(i)
3.	If you made a mistake in inputting $x_{{\mbox{\tiny K}}},$ then correct by	Xĸ	C	(k-1)
4.	Go to step 8 for moments calculations.			
	Grouped Data			
5.	Initialize the program.		xeq ΣMMTGD	ΣMMTGD
6.	Repeat step $6 \sim 7$ for $j=1,2,,m$ Input: $x_j$ $f_i$	X <sub>i</sub> f <sub>i</sub>	ENTER+)	(j)
7.	If you made a mistake in inputting $x_h$ and $f_h$ , then correct by	X <sub>h</sub> f <sub>h</sub>	ENTER+) C	(h-1)
8.	Calculate moments etc.: $\bar{x}$ $m_2$ $m_3$ $m_4$ $\gamma_1$ $\gamma_2$		E R/S R/S R/S R/S R/S	$XBAR = (\overline{x})$ $M2 = (m_2)$ $M3 = (m_3)$ $M4 = (m_4)$ $GM1 = (\gamma_1)$ $GM2 = (\gamma_2)$
9.	Repeat step 8 if you want the results again.			
10.	To use the same program for another set of data, initialize the program by $\rightarrow$ then go to step 2 or step 6.		<b>—</b> A	ΣMMTUG or ΣMMTGD
11.	To use the other program, go to step 1 or step 5.			

## **Examples:**

1. Ungrouped data

$$\bar{\mathbf{x}} = 4.21, \, \mathbf{m}_2 = 1.39, \, \mathbf{m}_3 = 0.39, \, \mathbf{m}_4 = 5.49$$

 $\gamma_1 = 0.24, \, \gamma_2 = 2.84$ 

Keystrokes:	Display:
XEQ ALPHA SIZE ALPHA 012	
$\begin{array}{c} \textbf{XEQ}  \textbf{ALPHA}  \boldsymbol{\Sigma} \text{ MMTUG}  \textbf{ALPHA} \end{array}$	ΣMMTUG
2.1 A 3.5 A 4.0 A 4.0 C	
4.2 A 6.5 A 4.1 A 3.6 A	
5.3 A 3.7 A 4.9 A	9.00
E	XBAR = 4.21
R/S	M2=1.39
R/S	M3=0.39
R/S	M4=5.49
R/S	GM1=0.24
R/S	GM2=2.84

## 2. Grouped data

i	1	2	3	4	5
Xi	3	2	4	6	1
f,	4	5	3	2	1

$$\bar{x} = 3.13, m_2 = 1.98, m_3 = 2.14, m_4 = 11.05$$

$$\gamma_1 = 0.77, \gamma_2 = 2.81$$

#### **Keystrokes: Display:** XEQ ALPHA SIZE ALPHA 012 **(XEQ) (ALPHA)** $\Sigma$ MMTGD **(ALPHA)** *SMMTGD* 3 ENTER+ 4 A 2 ENTER+ 5 A 4 ENTER+ 4 A 4 ENTER+ 4 C 4 ENTER+ 3 A 6 ENTER+ 2 A 1 ENTER+ 1 A 5.00 E XBAR=3.13 R/S M2=1.98 R/S M3=2.14 R/S M4=11.05 R/S GM1=0.77 R/S GM2=2.81

## ANALYSIS OF VARIANCE (ONE WAY)

The one-way analysis of variance is used to test if observed differences among k sample means can be attributed to chance or whether they are indicative of actual differences among the corresponding population means. Suppose the i<sup>th</sup> sample has  $n_i$  observations (samples may have equal or unequal number of observations). The null hypothesis we want to test is that the k population means are all equal. This program generates the complete ANOVA table.

1. Mean of observations in the  $i^{th}$  sample (i = 1, 2, ..., k)

$$\overline{\mathbf{x}}_{i} = \frac{1}{n_{i}} \sum_{j=1}^{n_{i}} \mathbf{x}_{ij}$$

2. Standard deviation of observations in the i<sup>th</sup> sample

$$s_i = \left[ \left( \sum_{j=1}^{n_i} x_{ij}^2 - n_i \bar{x}_i^2 \right) / (n_i - 1) \right]^{\frac{1}{2}}$$

3. Sum of observations in the i<sup>th</sup> sample

$$Sum_i = \sum_{j=1}^{n_i} x_{ij}$$

4. Total sum of squares

TSS = 
$$\sum_{i=1}^{k} \sum_{j=1}^{n_i} x_{ij}^2 - \frac{\left(\sum_{i=1}^{k} \sum_{j=1}^{n_i} x_{ij}\right)^2}{\sum_{i=1}^{k} n_i}$$

5. Treatment sum of squares

$$TrSS = \sum_{i=1}^{k} \frac{\left(\sum_{j=1}^{n_i} x_{ij}\right)^2}{n_i} - \frac{\left(\sum_{i=1}^{k} \sum_{j=1}^{n_i} x_{ij}\right)^2}{\sum_{i=1}^{k} n_i}$$

6. Error sum of squares

$$ESS = TSS - TrSS$$

7. Treatment degrees of freedom

$$df_1 = k - 1$$

8. Error degrees of freedom

$$df_2 = \sum_{i=1}^k n_i - k$$

9. Total degrees of freedom

$$df_3 = df_1 + df_2 = \sum_{i=1}^k n_i - 1$$

10. Treatment mean square

$$TrMS = \frac{TrSS}{df_1}$$

11. Error mean square

$$EMS = \frac{ESS}{df_2}$$

12. The F ratio

$$F = \frac{\text{TrMS}}{\text{EMS}} \text{(with degrees of freedom df_1, df_2)}$$

#### **Reference:**

J.E. Freund, Mathematical Statistics, Prentice Hall, 1962.

				<b>SIZE</b> : 020
STEP	INSTRUCTIONS	INPUT	FUNCTION	DISPLAY
1.	Initialize the program.		XEQ SAOVONE	ΣΑΟVONE
2.	Repeat step $2\sim5$ for $i=1,2,\ldots,k$ .			
3.	Repeat step 3~4 for j=1,2,,n <sub>i</sub> . Input x <sub>ij</sub> .	x <sub>ij</sub>	A	(j)
4.	If you made a mistake in inputting $\mathbf{x}_{im},$ then correct by	X <sub>im</sub>	C	(m-1)
5.	Calculate: mean $\bar{x}_i$ standard deviation s <sub>i</sub> sum Sum <sub>i</sub>		R/S R/S R/S	$\begin{array}{c} XBAR = (\overline{x}_i) \\ S = (s_i) \\ SUM = (Sum_i) \end{array}$
6.	To calculate ANOVA Table: TSS TrSS ESS df <sub>i</sub> df <sub>2</sub> df <sub>3</sub> TrMS EMS F		E R/S R/S R/S R/S R/S R/S R/S	$TSS = (TSS)$ $TRSS = (TrSS)$ $ESS = (ESS)$ $DF1 = (df_1)$ $DF2 = (df_2)$ $DF3 = (df_3)$ $TRMS = (TrMS)$ $EMS = (EMS)$ $F = (F)$
7.	Repeat step 6 if you want the results again.			
8.	For another set of data, initialize the program by $\rightarrow$ then go to step 2.		<b>A</b>	ΣΑΟVONE

## Example:

The following random samples of achievement test scores were obtained from students at four different schools:

j	1	2	3	4	5	6	7
School 1	88	99	96	68	85		
School 2	78	62	98	83	61	88	
School 3	80	61	74	92	78	54	77
School 4	71	65	90	46			

Calculate the ANOVA table and test the null hypothesis that the differences among the sample means can be attributed to chance. Use significance level  $\alpha = 0.01$ .

Keystrokes:	Display:
XEQ ALPHA SIZE ALPHA 020	
XEQ ALPHA SAOVONE ALPHA	ΣΑΟΥΟΝΕ
88 A 99 A 96 A 68 A	
85 A	5.00
R/S	XBAR = 87.20
R/S	S=12.15
R/S	SUM=436.00
78 A 62 A 98 A 83 A	
61 A 88 A	6.00
R/S	XBAR = 78.33
R/S	S=14.62
R/S	SUM=470.00
80 A 61 A 74 A 92 A	
78 A 54 A 77 A	7.00
	XBAR = 73.71
	S=12.61
R/S	SUM=516.00
71 A 66 A 66 C 65 A	
90 A 46 A	4.00
R/S	XBAR = 68.00
R/S	S=18.13
R/S	SUM=272.00
E	TSS=4530.00
R/S	TRSS=930.44
	ESS=3599.56
	DF1=3.00
	DF2=18.00
R/S	DF3=21.00
R/S	TRMS=310.15
R/S	EMS = 199.98
(R/S)	F=1.55

## **ANOVA** Table

	SS	df	MS	F
Treatments	930.44	3	310.15	1.55
Error	3599.56	18	199.98	
Total	4530.00	21		

Since F = 1.55 does not exceed  $F_{.01,3,18} = 5.09$ , the null hypothesis can not be rejected. Thus we have no evidence to conclude that the means of the scores for the four schools are significantly different.

#### :

## ANALYSIS OF VARIANCE (TWO WAY, NO REPLICATIONS)

The analysis of variance is the analysis of the total variability of a set of data (measured by their total sum of squares) into components which can be attributed to different sources of variation.

The two way analysis of variance tests the row effects and the column effects independently. This program will generate the ANOVA table for the case such that (1) each cell only has one observation and (2) the row and column effects do not interact.

#### **Equations:**

1. Sums

Row 
$$RS_i = \sum_{j} x_{ij}$$
  $i = 1, 2, ..., r$ 

Column 
$$CS_j = \sum_{i} x_{ij}$$
  $j = 1, 2, ..., c$ 

2. Sums of squares

Total TSS = 
$$\Sigma \Sigma x_{ij}^2 - (\Sigma \Sigma x_{ij})^2/rc$$

Row RSS = 
$$\sum_{i} \left(\sum_{j} x_{ij}\right)^2 / c - (\Sigma \Sigma x_{ij})^2 / rc$$

Column CSS = 
$$\sum_{j} \left( \sum_{i} x_{ij} \right)^2 / r - (\Sigma \Sigma x_{ij})^2 / rc$$

Error ESS = TSS - RSS - CSS

3. Degrees of freedom

Row 
$$df_1 = r - 1$$

Column  $df_2 = c - 1$ 

Error 
$$df_3 = (r - 1) (c - 1)$$

4. F ratios

Row 
$$F_1 = \frac{RSS}{df_1} / \frac{ESS}{df_3}$$
  
Column  $F_2 = \frac{CSS}{df_2} / \frac{ESS}{df_3}$ 

## **Reference:**

Dixon and Massey, Introduction to Statistical Analysis, McGraw-Hill, 1969.

				<b>SIZE</b> : 018
STEP	INSTRUCTIONS	INPUT	FUNCTION	DISPLAY
1.	Initialize the program.		χέο Σαοντωο	ΣΑΟντωο
	Row-Wise			
2.	Repeat step $2\sim5$ for $i=1,2,\ldots,r$ .			
3.	Repeat step $3 \sim 4$ for $j=1,2,,c$ . Input $x_{ij}$	X <sub>ij</sub>	A	(j)
4.	If you made a mistake in inputting $x_{\rm im},$ then correct by	X <sub>im</sub>	C	(m-1)
5.	Calculate row sum and initialize the program for the next row.		R/S	SUM=(RS <sub>i</sub> )
6.	After completion of the last row, (row r) initialize the program for column-wise data entry.		R/S	COLUMN-WISE
	Column-Wise			
7.	Repeat step $7 \sim 10$ for $j = 1, 2,, c$ .			
8.	Repeat step 8~9 for $i=1,2,,r$ . Input $x_{ij}$ .	X <sub>ij</sub>	A	(i)
9.	If you made a mistake in inputting $x_{\rm hj},$ then correct by	X <sub>hj</sub>	C	(h-1)
10.	Calculate column sum and initialize the program for the next column.		R/S	SUM=(CS <sub>i</sub> )
11.	Calculate ANOVA Table: RSS CSS TSS ESS df <sub>1</sub> df <sub>2</sub> df <sub>3</sub> $F_1$ $F_2$ Descent attack of a figure at the		E R/S R/S R/S R/S R/S R/S	$RSS = (RSS) \\ CSS = (CSS) \\ TSS = (TSS) \\ ESS = (ESS) \\ DF1 = (df_1) \\ DF2 = (df_2) \\ DF3 = (df_3) \\ F1 = (F_1) \\ F2 = (F_2) \\ \end{cases}$
12.	Repeat step 11 if you want the results again			
13.	For another set of data, initialize the program by $\rightarrow$ then go to step 2			ΣΑΟVΤWO

## Example:

Apply this program to analyze the following set of data.



## **Keystrokes:**

**Display:** 

XEQ ALPHA SIZE ALPHA 018	
XEQ ALPHA SAOVTWO ALPHA	ΣΑΟΥΤWO
7 A 6 A 8 A 7 A	4.00
R/S	SUM=28.00
2 A 4 A 4 A 4 A	4.00
(R/S)	SUM=14.00
4 A 7 A 7 C 6 A 5 A	
3 <b>A</b>	4.00
R/S	SUM=18.00
R/S	COLUMN-WISE
7 A 2 A 4 A	3.00
R/S	SUM=13.00
6 A 4 A 6 A	3.00
R/S	SUM=16.00
8 A 4 A 5 A	3.00
R/S	SUM=17.00
7 A 4 A 3 A	3.00
R/S	SUM=14.00
E	RSS=26.00
R/S	CSS=3.33
R/S	TSS=36.00
R/S	ESS=6.67
R/S	DF1=2.00
R/S	DF2=3.00
R/S	DF3=6.00
R/S	F1=11.70
R/S	F2=1.00

ANOVA

	SS	df	F ratio
Row	26.00	2	11.70
Column	3.33	3	1.00
Error	6.67	6	
Total	36.00		

## ANALYSIS OF COVARIANCE (ONE WAY)

The one way analysis of covariance program tests the effect of one variable separately from the effect of a second variable, if the second variable represents an actual measurement for each individual (rather than a category).

Suppose  $(x_{ij}, y_{ij})$  represents the j<sup>th</sup> observation from the i<sup>th</sup> population (i = 1,2, ..., k, j = 1,2, ..., n<sub>i</sub>). Note that samples may have equal or unequal number of observations. The analysis of covariance tests for a difference in means of residuals. The residuals are the differences of the observations and a regression quantity based on the associated second variable. The analysis of covariance procedure is based on the separations of the sums of squares and the sums of products into several portions. This program will generate the complete ANOCOV table.

#### **Equations:**

1. Sums and sums of squares

$$Sx_{i} = \sum_{j} x_{ij} (i = 1, 2, ..., k)$$
$$TSSx = \Sigma \Sigma x_{ij}^{2} - \frac{(\Sigma \Sigma x_{ij})^{2}}{\sum_{i} n_{i}}$$
$$ASSx = \sum_{i} \frac{\left(\sum_{j} x_{ij}\right)^{2}}{n_{i}} - \frac{(\Sigma \Sigma x_{ij})^{2}}{\sum_{i} n_{i}}$$

$$WSSx = TSSx - ASSx$$

2. Degrees of freedom

$$df_1 = k - 1$$
$$df_2 = \sum_{i} n_i - k$$

3. Mean squares and F statistic

$$AMSx = \frac{ASSx}{df_1}$$

$$WMSx = \frac{WSSx}{df_2}$$

$$F_x = \frac{AMSx}{WMSx}$$
 with degrees of freedom df\_1, df\_2

By changing  $x_{ij}$  to  $y_{ij}$ , similar formulas for  $y_{ij}$  can be obtained.

4. Sums of products

$$TSP = \Sigma\Sigma x_{ij} y_{ij} - \frac{(\Sigma\Sigma x_{ij}) (\Sigma\Sigma y_{ij})}{\sum_{i} n_{i}}$$
$$ASP = \sum_{i} \frac{\left(\sum_{j} x_{ij}\right) \left(\sum_{j} y_{ij}\right)}{n_{i}} - \frac{(\Sigma\Sigma x_{ij}) (\Sigma\Sigma y_{ij})}{\sum_{i} n_{i}}$$

$$WSP = TSP - ASP$$

5. Residual sums of squares

$$TSS\hat{y} = TSSy - \frac{(TSP)^2}{TSSx}$$
$$WSS\hat{y} = WSSy - \frac{(WSP)^2}{WSSx}$$
$$ASS\hat{y} = TSS\hat{y} - WSS\hat{y}$$

6. Residual degrees of freedom

$$df_3 = k - 1$$
$$df_4 = \sum_{i} n_i - k - 1$$

## 28 Analysis of Covariance

7. Residual mean squares and F statistic

$$AMS\hat{y} = \frac{ASSy}{df_3}$$
$$WMS\hat{y} = \frac{WSS\hat{y}}{df_4}$$

$$F = \frac{AMS\hat{y}}{WMS\hat{y}}$$
 with degrees of freedom df<sub>3</sub>, df<sub>4</sub>

. . . . .

#### **ANOCOV** Table

						Resid	duals	
	degrees of freedom	SSx	SP	SSy	degrees of freedom	SSŷ	MSŷ	F statistic
Among means	df₁	ASSx	ASP	ASSy	df <sub>3</sub>	ASSŷ	AMSŷ	F
Within groups	df₂	WSSx	WSP	WSSy	df₄	WSSŷ	WMSŷ	
Total		TSSx	TSP	TSSy		TSSŷ		

#### **Remarks:**

- $F_x$  can be used to test if the X means are equal (ANOVA for X).
- F<sub>y</sub> can be used to test if the Y means (not making use of the X values) are equal (ANOVA for unadjusted Y).

#### **Reference:**

Dixon and Massey, Introduction to Statistical Analysis, McGraw-Hill, 1969.

				<b>SIZE</b> : 026
STEP	INSTRUCTIONS	INPUT	FUNCTION	DISPLAY
1.	Initialize the program.		xeq ΣANOCOV	∑ANOCOV (pse) NEW I=1.00
2.	Repeat step 2 $\sim$ 6 for i=1,2,,k.			
3.	Repeat step $3 \sim 4$ for $j = 1, 2,, n_j$ . Input $x_{ij}$ and $y_{ij}$ .	x <sub>ij</sub> y <sub>ij</sub>		(j)
4.	If you made a mistake in inputting $x_{\rm im}$ and $y_{\rm im},$ then correct by	X <sub>im</sub> Y <sub>im</sub>		(m-1)
5.	Calculate the $i^{\text{th}}$ sums: $Sx_i \\ Sy_i$		R/S R/S	$SX = (Sx_i)$ $SY = (Sy_i)$
6.	Initialize for new i.		R/S	NEW I=(i)
7.	To calculate ANOCOV Table: TSSx ASSx WSSx TSSy ASSy WSSy df <sub>1</sub> df <sub>2</sub> Fx Fy TSP ASP WSP TSS $\hat{y}$ WSS $\hat{y}$ df <sub>3</sub> df <sub>4</sub> AMS $\hat{y}$ WMS $\hat{y}$ F		E R/S R/S R/S R/S R/S R/S R/S R/S	$\begin{split} &TSSX = (TSSx)\\ &ASSX = (ASSx)\\ &WSSX = (WSSx)\\ &TSSY = (TSSy)\\ &ASSY = (ASSy)\\ &WSSY = (WSSy)\\ &DF1 = (df_1)\\ &DF2 = (df_2)\\ &FX = (Fx)\\ &FY = (Fy)\\ &TSP = (TSP)\\ &ASP = (ASP)\\ &WSP = (WSP)\\ &TSSY = (WSS\hat{y})\\ &MSSY = (WSS\hat{y})\\ &DF3 = (df_3)\\ &DF4 = (df_4)\\ &AMSY = (AMS\hat{y})\\ &WMSY = (WMS\hat{y})\\ &F = (F) \end{split}$
9.	results again. For another set of data, initialize the program by $\rightarrow$ then go to step 2.		<b>a</b>	ンANOCOV (pse) NEW I=1.00

## **Example:**

			j			
			1	2	З	4
		х	3	2	1	2
	1	у	10	8	8	11
		х	4	3	3	5
•	2	у	12	12	10	13
		х	1	2	3	1
	3	у	6	5	8	7

 $(k = 3, n_1 = n_2 = n_3 = 4)$ 

#### **Keystrokes:**

XEQ ALPHA SIZE ALPHA 026 XEQ ALPHA SANOCOV ALPHA 3 [ENTER+] 10 [A] 2 [ENTER+] 8 [A] 5 (ENTER+) 5 (A) 5 (ENTER+) 5 (C) 1 ENTER+ 8 A 2 ENTER+ 11 A R/S R/S R/S 4 ENTER+ 12 A 3 ENTER+ 12 A 3 [ENTER+] 10 A 5 [ENTER+] 13 A R/S R/S R/S 1 ENTER+ 6 A 2 ENTER+ 5 A 3 ENTER+ 8 A 1 ENTER+ 7 A R/S R/S R/S E R/S R/S R/S R/S R/S R/S R/S R/S

#### Display:

∑ANOCOV (Pse) NEW I=1.00 4.00

SX=8.00 SY=37.00 NEW I=2.00

4.00 SX=15.00 SY=47.00 NEW I=3.00

4.00 SX = 7.00 SY = 26.00 NEW I = 4.00 TSSX = 17.00 ASSX = 9.50 WSSX = 7.50 TSSY = 71.67 ASSY = 55.17 WSSY = 16.50 DF1 = 2.00 DF2 = 9.00 FX = 5.70

R/S	FY=15.05
R/S	TSP=27.00
R/S	ASP=20.75
R/S	WSP=6.25
R/S	TSSY.=28.78
R/S	WSSY.=11.29
R/S	ASSY. =17.49
R/S	DF3=2.00
R/S	DF4=8.00
R/S	AMSY.=8.75
R/S	WMSY. = 1.41
R/S	F=6.20

## ANOCOV Table

					Residuals			
	df	SSx	SP	SSy	df	SSŷ	MSŷ	F
Among means	2	9.50	20.75	55.17	2	17.49	8.75	6.20
Within groups	9	7.50	6.25	16.50	8	11.29	1.41	
Total		17.00	27.00	71.67		28.78		

#### **CURVE FITTING**

For a set of data points  $(x_i, y_i)$ , i = 1, 2, ..., n, this program can be used to fit the data to any of the following curves:

- 1. Straight line (linear regression); y = a + bx.
- 2. Exponential curve;  $y = ae^{bx}$  (a > 0).
- 3. Logarithmic curve;  $y = a + b \ln x$ .
- 4. Power curve;  $y = ax^b$  (a> 0).

The regression coefficients a and b are found from solving the following system of linear equations:

$$\begin{bmatrix} n & \Sigma X_i \\ \Sigma X_i & \Sigma X_i^2 \end{bmatrix} \begin{bmatrix} A \\ b \end{bmatrix} = \begin{bmatrix} \Sigma Y_i \\ \Sigma Y_i X_i \end{bmatrix}$$

where the variables are defined as follows:

Regression	A	Xi	Y
Linear	а	x <sub>i</sub>	y,
Exponential	In a	×i	In y <sub>i</sub>
Logarithmic	а	ln x <sub>i</sub>	y,
Power	In a	In x <sub>i</sub>	In y <sub>i</sub>

The coefficient of determination is:

$$R^{2} = \frac{A\Sigma Y_{i} + b\Sigma X_{i} Y_{i} - \frac{1}{n} (\Sigma Y_{i})^{2}}{\Sigma (Y_{i}^{2}) - \frac{1}{n} (\Sigma Y_{i})^{2}}$$

#### Linear Regression







#### **Remarks:**

- The program applies the least square method, either to the original equations (straight line and logarithmic curve) or to the transformed equations (exponential curve and power curve).
- Negative and zero values of x<sub>i</sub> will cause a machine error for logarithmic curve fits. Negative and zero values of y<sub>i</sub> will cause a machine error for exponential curve fits. For power curve fits, both x<sub>i</sub> and y<sub>i</sub> must be positive, non-zero values.
- As the differences between x and/or y values become small, the accuracy of the regression coefficients will decrease.

				<b>SIZE</b> : 016
STEP	INSTRUCTIONS	INPUT	FUNCTION	DISPLAY
1.	Initialize the program. ● for STRAIGHT LINE→ or ● for EXPONENTIAL CURVE→ or ● for LOGARITHMIC CURVE→ or ● for POWER CURVE→		xeq ΣLIN (xeq Σexp (xeq Σlog (xeq Σpow	ΣLIN ΣEXP ΣLOG ΣPOW
2.	$ \begin{array}{l} \mbox{Repeat step 2-3 for I} = 1,2 \ldots,n. \\ \mbox{Input } x_i \\ y_i \end{array} $	Xi Vi	ENTER+ A	(i)
3.	If you made a mistake in inputting $x_k$ and $y_k$ , then correct by	Хк Ук	ENTER+ C	(k – 1)
4.	Calculate R <sup>2</sup> and regression coefficients a and b.		E R/S R/S	$R2 = (R^2)$ a = (a) b = (b)
5.	Calculate estimated y from regression. Input x.	x	R/S	$Y_{\cdot} = (\hat{y})$
6.	Repeat step 5 for different x's.			
7.	Repeat step 4 if you want the results again.			
8.	To use the same program for another set of data, initialize the program by $\rightarrow$		A	ΣLIN or ΣEXP or ΣLOG or ΣPOW
	then go to step 2.			
9.	To use another program, go to step 1.			

## Example 1:

Fit the following set of data into a straight line.

$\mathbf{X}_{i}$	40.5	38.6	37.9	36.2	35.1	34.6
y,	104.5	102	100	97.5	95.5	94
### Solution:

a = 33.53, b = 1.76 R<sup>2</sup> = 0.99 i.e., y = 33.53 + 1.76 x For x = 37,  $\hat{y}$  = 98.65 For x = 35,  $\hat{y}$  = 95.13

Display:
ΣLIN
6.00
R2=0.99
a=33.53
b=1.76
Y.=98.65
Y.=95.13

# Example 2:

Fit the following set of data into an exponential curve.

Xi	.72	1.31	1.95	2.58	3.14	
y <sub>i</sub>	2.16	1.61	1.16	.85	0.5	

#### Solution:

a = 3.45, b = -0.58y = 3.45 e<sup>-0.58x</sup> R<sup>2</sup> = 0.98 For x = 1.5,  $\hat{y}$  = 1.44 For x = 2,  $\hat{y}$  = 1.08

Keystrokes:	Display:
XEQ ALPHA SIZE ALPHA 016	
XEQ ALPHA DEXP ALPHA	ΣΕΧΡ
.72 ENTER+) 2.16 A	
1.31 ENTER+ 1.61 A	
1.95 ENTER+ 1.16 A	
2.58 ENTER+ .85 A	
3.15 ENTER+ .05 A	
3.15 ENTER+ .05 C	
3.14 ENTER+ 0.5 A	5.00
E	R2=0.98
R/S	a=3.45
R/S	b=-0.58
1.5 <b>R/S</b>	Y.=1.44
2.0 <b>R/S</b>	Y.=1.08

## Example 3:

Fit the following set of data into a logarithmic curve.

Xi	3	4	6	10	12	
y,	1.5	9.3	23.4	45.8	60.1	

# Solution:

a = -47.02, b = 41.39y =  $-47.02 + 41.39 \ln x$ R<sup>2</sup> = 0.98For x = 8,  $\hat{y} = 39.06$ For x = 14.5,  $\hat{y} = 63.67$ 

# **Keystrokes:**

-
XEQ ALPHA SIZE ALPHA 016
XEQ ALPHA ELOG ALPHA
3 ENTER+) 1.5 A
4 ENTER+ 9.3 A
6 ENTER+) 23.4 A
10 ENTER+) 45.8 A
12 ENTER+) 6.01 A
12 ENTER+) 6.01 C
12 ENTER+) 60.1 A

# **Display:**



E	R2=0.98
R/S	a=-47.02
R/S	b=41.39
8 <b>R/S</b>	Y.=39.06
14.5 <b>R/S</b>	Y.=63.67

# Example 4:

Fit the following set of data into a power curve.

$\mathbf{x}_{i}$	10	12	15	17	20	22	25	27	30	32	35	
y,	0.95	1.05	1.25	1.41	1.73	2.00	2.53	2.98	3.85	4.59	6.02	

# Solution:

a = .03, b = 1.46 y = .03x<sup>1.46</sup> R<sup>2</sup> = 0.94 For x = 18,  $\hat{y} = 1.76$ For x = 23,  $\hat{y} = 2.52$ 

# **Keystrokes:**

**Display:** 

$\begin{array}{c} \textbf{XEQ}  \textbf{ALPHA}  \textbf{SIZE}  \textbf{ALPHA}  \textbf{016} \\ \hline \textbf{XEQ}  \textbf{ALPHA}  \boldsymbol{\Sigma} \textbf{POW}  \textbf{ALPHA} \end{array}$	ΣΡΟΨ
10 [ENTER+] 0.95 [A]	
12 [ENTER+] 1.05 [A]	
15 [ENTER+] 1.25 [A]	
17 ENTER+) 1.41 A	
20 ENTER+) 1.73 A	
22 ENTER+) 2.00 A	
25 ENTER+) 2.53 A	
27 ENTER+) 2.98 A	
30 ENTER+ 3.85 A	
32 ENTER+) 4.59 A	
35 ENTER+) 60.2 A	
35 ENTER+) 60.2 C	
35 ENTER+) 6.02 A	11.00
E	R2=0.94
R/S	a=0.03
R/S	b=1.46
18 <b>R/S</b>	Y.=1.76
23 <b>R/S</b>	Y.=2.52

### **MULTIPLE LINEAR REGRESSION**

### **Three Independent Variables**

For a set of data points  $\{(x_i, y_i, z_i, t_i), i = 1, 2, ..., n\}$ , this program fits a linear equation of the form:

$$\mathbf{t} = \mathbf{a} + \mathbf{b}\mathbf{x} + \mathbf{c}\mathbf{y} + \mathbf{d}\mathbf{z}$$

by the least squares method.

Regression coefficients a, b, c, and d are calculated by solving the following system of equations. Gauss's elimination method with partial pivoting is used.

$$\begin{bmatrix} n & \Sigma x_i & \Sigma y_i & \Sigma z_i \\ \Sigma x_i & \Sigma (x_i)^2 & \Sigma (x_i y_i) & \Sigma (x_i z_i) \\ \Sigma y_i & \Sigma (y_i x_i) & \Sigma (y_i)^2 & \Sigma (y_i z_i) \\ \Sigma z_i & \Sigma (x_i z_i) & \Sigma (y_i z_i) & \Sigma (z_i)^2 \end{bmatrix} \quad \begin{bmatrix} a \\ b \\ c \\ d \end{bmatrix} = \begin{bmatrix} \Sigma t_i \\ \Sigma x_i t_i \\ \Sigma y_i t_i \\ \Sigma z_i t_i \end{bmatrix}$$

The coefficient of determination  $R^2$  is defined as:

$$R^{2} = \frac{a\Sigma t_{i} + b\Sigma x_{i}t_{i} + c\Sigma y_{i}t_{i} + d\Sigma z_{i}t_{i} - \frac{1}{n} (\Sigma t_{i})^{2}}{\Sigma (t_{i}^{2}) - \frac{1}{n} (\Sigma t_{i})^{2}}$$

#### **Two Independent Variables**

For a set of data points  $\{(x_i, y_i, t_i), i = 1, 2, ..., n\}$ , this program fits a linear equation of the form:

$$\mathbf{t} = \mathbf{a} + \mathbf{b}\mathbf{x} + \mathbf{c}\mathbf{y}$$

by the least squares method.

Regression coefficients a, b, and c are calculated by solving the following system of equations. Gauss's elimination method with partial pivoting is used.

$$\begin{bmatrix} n & \Sigma x_i & \Sigma y_i \\ \Sigma x_i & \Sigma (x_i)^2 & \Sigma x_i y_i \\ \Sigma y_i & \Sigma y_i x_i & \Sigma (y_i)^2 \end{bmatrix} \quad \begin{bmatrix} a \\ b \\ c \end{bmatrix} = \begin{bmatrix} \Sigma t_i \\ \Sigma x_i t_i \\ \Sigma y_i t_i \end{bmatrix}$$

The coefficient of determination  $R^2$  is defined as:

$$R^{2} = \frac{a\Sigma t_{i} + b\Sigma x_{i}t_{i} + c\Sigma y_{i}t_{i} - \frac{1}{n} (\Sigma t_{i})^{2}}{\Sigma (t_{i}^{2}) - \frac{1}{n} (\Sigma t_{i})^{2}}$$

## **Remarks:**

- If the coefficient matrix has determinant equal to zero, indicating no solution or more than one solution, "DATA ERROR" will be displayed.
- There is no restriction on the maximum number of data points n, but the following minimum condition for n must be satisfied:
  - $n \ge 3$  for the case of two independent variables  $n \ge 4$  for the case of three independent variables

#### **Refernce:**

HP-67/97 Math Pac I, program MA1-07

				<b>SIZE</b> : 045
STEP	INSTRUCTIONS	INPUT	FUNCTION	DISPLAY
-	Three Independent Variables			
1.	Initialize the program.		xeq ΣMLRXYZ	ΣMLRXYZ
2.	$\begin{array}{l} \mbox{Repeat step } 2{\sim}3 \mbox{ for } i{=}1,2,\ldots,n. \\ \mbox{Input: } x_i \\ y_i \\ z_i \\ t_i \end{array}$	X <sub>i</sub> Y <sub>i</sub> Z <sub>i</sub> t <sub>i</sub>	ENTER+) ENTER+) ENTER+) A	(i)
3.	If you made a mistake in inputting $x_k$ , $y_k$ , $z_k$ , and $t_k$ , then correct by	X <sub>k</sub> Yk Zk t <sub>k</sub>	ENTER+) ENTER+) ENTER+) C	(k-1)
4.	Calculate R <sup>2</sup> and regression coefficients a,b,c, and d.		E R/S R/S R/S R/S	R2 = (R2)a = (a)b = (b)c = (c)d = (d)
5. 6.	Calculate estimated t from regression. Input: x y z Repeat step 5 for different (x,y,z)'s.	x y z	ENTER+) ENTER+) (R/S)	T.=(î)
7.	To recall sums used in calculation: $\Sigma x_i$ $\Sigma y_i$ $\Sigma z_i$ $\Sigma t_i$ $\Sigma x_i^2$ $\Sigma y_i^2$ $\Sigma z_i^2$ $\Sigma t_i^2$ $\Sigma x_i y_i$ $\Sigma x_i z_i$ $\Sigma x_i t_i$ $\Sigma y_i z_i$ $\Sigma y_i z_i$ $\Sigma y_i t_i$ $\Sigma z_i t_i$		RCL 32 RCL 33 RCL 34 RCL 41 RCL 35 RCL 38 RCL 30 RCL 30 RCL 30 RCL 37 RCL 39 RCL 43 RCL 43	$\begin{array}{c} (\Sigma x_i) \\ (\Sigma y_i) \\ (\Sigma z_i) \\ (\Sigma t_i) \\ (\Sigma x_i^2) \\ (\Sigma y_i^2) \\ (\Sigma z_i^2) \\ (\Sigma z_i^2) \\ (\Sigma z_i^2) \\ (\Sigma x_i y_i) \\ (\Sigma x_i z_i) \\ (\Sigma x_i z_i) \\ (\Sigma y_i z_i) \\ (\Sigma y_i z_i) \\ (\Sigma y_i z_i) \\ (\Sigma z_i t_i) \end{array}$
8.	Repeat step 4 if you want the results again.			
9.	To use the program for another set of data, initialize the program by $\rightarrow$ then go to step 2.		<b>A</b>	ΣMLRXYZ

#### Example 1:

For the following set of data, find the regression line with three independent variables. i.e. t = a + bx + cy + dz

, i	1	2	3	4	5	
Xi	7	1	11	11	7	
y,	25	29	56	31	52	
Zi	6	15	8	8	6	
ti	60	52	20	47	33	

#### Solution:

The regression line is described by t = 103.45 - 1.28x - 1.04y - 1.34z.

 $R^2 = 1.00$ For x = 7, y = 25, z = 6,  $\hat{t} = 60.50$ For x = 1, y = 29, z = 15,  $\hat{t} = 52.00$ 

### **Keystrokes:**

#### **Display:**

XEQ ALPHA SIZE ALPHA 045	
XEQ ALPHA SMLRXYZ ALPHA	SMLRXYZ
7 ENTER+) 25 ENTER+)	
6 ENTER+) 60 A	
1 ENTER+ 29 ENTER+	
15 ENTER+ 52 A	
11 ENTER+ 56 ENTER+	
8 ENTER+ 20 A	
11 ENTER+ 31 ENTER+	
8 [ENTER+] 47 [A]	
6 ENTER+ 33 A	
C ENTERN 33 C	
6 ENTERA 22 A	5.00
F	$B_{2} = 1.00$
B/S	a = 103.45
B/S	h = -1.28
R/S	c = -1.04
	d=-1.34
7 [ENTER+] 25 [ENTER+] 6 [R/S]	T.=60.50
1 [ENTER+] 29 [ENTER+] 15 [R/S]	T.=52.00

				<b>SIZE</b> : 045
STEP	INSTRUCTIONS	INPUT	FUNCTION	DISPLAY
1	Two Independent Variables			
1.				
2.	Repeat step 2~3 for i=1,2,,n. Input: $x_i$ $y_i$ $t_i$	x <sub>i</sub> y <sub>i</sub> t <sub>i</sub>	ENTER• ENTER• A	(i)
3.	If you made a mistake in inputting $x_k, y_k$ , and $t_k$ , then correct by	x <sub>k</sub> y <sub>k</sub> t <sub>k</sub>	ENTER+ ENTER+ C	(k-1)
4.	Calculate R <sup>2</sup> and regression coefficients a, b, and c.		E R/S R/S R/S	$R2 = (R^{2}) a = (a) b = (b) c = (c)$
5.	Calculate estimated t from regression. Input: x y	x y	ENTER+) R/S	T.=(î)
6. 7.	Repeat step 5 for different $(x,y)$ 's. To recall sums used in calculation: $\Sigma x_i$ $\Sigma y_i$ $\Sigma t_i$ $\Sigma x_i^2$ $\Sigma y_i^2$ $\Sigma t_i^2$ $\Sigma x_i y_i$ $\Sigma x_i y_i$ $\Sigma x_i t_i$ $\Sigma y_i t_i$		RCL 32 RCL 33 RCL 41 RCL 35 RCL 38 RCL 30 RCL 36 RCL 42 RCL 43	$\begin{array}{c} (\Sigma x_{i}) \\ (\Sigma y_{i}) \\ (\Sigma y_{i}) \\ (\Sigma t_{i}) \\ (\Sigma x_{i}^{2}) \\ (\Sigma y_{i}^{2}) \\ (\Sigma t_{i}^{2}) \\ (\Sigma x_{i} y_{i}) \\ (\Sigma x_{i} t_{i}) \\ (\Sigma y_{i} t_{i}) \end{array}$
8.	Repeat step 4 if you want the results again.			
9.	To use the program for another set of data, initialize the program by $\rightarrow$ then go to step 2.		A	ΣMLRXY

## Example 2:

For the following set of data, find the regression line with two independent variables. i.e. t = a + bx + cy

	1	2	3	4
Xi	1.5	0.45	1.8	2.8
y,	0.7	2.3	1.6	4.5
ti	2.1	4.0	4.1	9.4

## Solution:

The regression line is t = -0.10 + 0.79x + 1.63y

 $R^2 = 1.00$ For x = 2, y = 3,  $\hat{t} = 6.37$ For x = 1.5, y = 0.7  $\hat{t} = 2.23$ 

T

## **Keystrokes:**

### **Display:**

XEQ ALPHA SIZE ALPHA 045	
$\begin{array}{c} \textbf{XEQ}  \textbf{ALPHA}  \boldsymbol{\Sigma} \textbf{MLRXY}  \textbf{ALPHA} \end{array}$	ΣMLRXY
1.5 ENTER+ 0.7 ENTER+ 2.1 A	
0.46 ENTER+ 2.3 ENTER+ 4.0 A	
0.46 ENTER+ 2.3 ENTER+ 4.0 C	
0.45 ENTER+ 2.3 ENTER+ 4.0 A	
1.8 ENTER+ 1.6 ENTER+ 4.1 A	
2.8 ENTER+ 4.5 ENTER+ 9.4 A	4.00
E	R2=1.00
R/S	a=-0.10
R/S	b=0.79
R/S	c=1.63
2 ENTER+ 3 R/S	T.=6.37
1.5 ENTER+) 0.7 R/S	T.=2.23

## **POLYNOMIAL REGRESSION**

#### **Cubic Regression**

For a set of data points  $(x_i, y_i)$ , i = 1, 2, ..., n, this program fit a cubic equation of the form:

$$y = a + bx + cx^2 + dx^3$$

by the least squares method.

Regression coefficients a, b, c and d are calculated by solving the following system of equations. Gauss's elimination method with partial pivoting is used.

n	$\Sigma x_i$	$\Sigma x_i^{\ 2}$	$\Sigma x_i^3$	a		$\Sigma y_i$
$\Sigma x_i$	$\Sigma x_i^{\ 2}$	$\Sigma x_i^{\ 3}$	$\Sigma x_i^4$	b	_	$\Sigma x_i y_i$
$\Sigma x_i^2$	$\Sigma x_i^{\ 3}$	$\Sigma x_i^4$	$\Sigma x_i^{5}$	c	=	$\Sigma x_i^{\ 2} y_i$
$\Sigma x_i^3$	$\Sigma x_i^4$	$\Sigma x_i^{5}$	$\Sigma x_i^{6}$	d		$\Sigma x_i^{\;3}y_i$

The coefficient of determination is:

$$R^{2} = \frac{a\Sigma y_{i} + b\Sigma x_{i}y_{i} + c\Sigma x_{i}^{2}y_{i} + d\Sigma x_{i}^{3}y_{i} - \frac{1}{n} (\Sigma y_{i})^{2}}{\Sigma (y_{i}^{2}) - \frac{1}{n} (\Sigma y_{i})^{2}}$$

#### **Parabolic Regression**

For a set of data points  $(x_i, y_i)$ , i = 1, 2, ..., n, this program fits a parabola of the form:

$$y = a + bx + cx^2$$

by the least squares method.

Regression coefficients a, b, and c are calculated by solving the following system of equations. Gauss's elimination method with partial pivoting is used.

$$\begin{bmatrix} n & \Sigma x_i & \Sigma x_i^2 \\ \Sigma x_i & \Sigma x_i^2 & \Sigma x_i^3 \\ \Sigma x_i^2 & \Sigma x_i^3 & \Sigma x_i^4 \end{bmatrix} \begin{bmatrix} a \\ b \\ c \end{bmatrix} = \begin{bmatrix} \Sigma y_i \\ \Sigma x_i y_i \\ \Sigma x_i^2 y_i \end{bmatrix}$$

The coefficient of determination is:

$$R^{2} = \frac{a\Sigma y_{i} + b\Sigma x_{i}y_{i} + c\Sigma x_{i}^{2}y_{i} - \frac{1}{n} (\Sigma y_{i})^{2}}{\Sigma (y_{i}^{2}) - \frac{1}{n} (\Sigma y_{i})^{2}}$$

### **Remarks:**

- If the coefficient matrix has determinant equal to zero, indicating no solution or more than one solution, *DET=0* will be displayed.
- There is no restriction on the maximum number of data points n, but the following minimum condition for n must be satisfied:

$$n \ge 3$$
 for Parabolic Regression  
 $n \ge 4$  for Cubic Regression

#### **Reference:**

HP-67/97 Math Pac I, program MA1-07

				<b>SIZE</b> : 045
STEP	INSTRUCTIONS	INPUT	FUNCTION	DISPLAY
	Cubic Regression			
1.	Initialize the program.		XEQ SPOLYC	ΣPOLYC
2.	Repeat step 2~3 for i=1,2,,n. Input: $x_i$	Xi Vi	ENTER+) A	(i)
3.	If you made a mistake in inputting			
	$\boldsymbol{x}_k$ and $\boldsymbol{y}_k,$ then correct by	Х <sub>к</sub> Ук	ENTER+) C	(k-1)
4.	Calculate R <sup>2</sup> and regression coefficients a,b,c, and d.		E R/S R/S R/S R/S	R2 = (R2)a = (a)b = (b)c = (c)d = (d)
5.	Calculate estimated y from regression. Input x.	x	R/S	Y.=(ŷ)
6.	Repeat step 5 for different x's.			
7.	To recall sums in calculation: $\Sigma x_i$ $\Sigma x_i^2$ $\Sigma x_i^3$ $\Sigma x_i^4$ $\Sigma x_i^5$ $\Sigma x_i^6$ $\Sigma y_i$ $\Sigma x_i y_i$ $\Sigma x_i^2 y_i$ $\Sigma x_i^3 y_i$		RCL       32         RCL       33         RCL       34         RCL       37         RCL       39         RCL       40         RCL       41         RCL       42         RCL       43         RCL       44	$\begin{array}{c} (\Sigma x_i) \\ (\Sigma x_i^2) \\ (\Sigma x_i^3) \\ (\Sigma x_i^4) \\ (\Sigma x_i^5) \\ (\Sigma x_i^6) \\ (\Sigma y_i) \\ (\Sigma x_i y_i) \\ (\Sigma x_i^2 y_i) \\ (\Sigma x_i^3 y_i) \end{array}$
8.	Repeat step 4 if you want the results again.			
9.	To use the program for another set of data, initialize the program by $\rightarrow$ then go to step 2.			ΣPOLYC

### Example 1:

For the following set of data, perform a cubic regression, i.e., find suitable coefficients for:

$y = a + bx + cx^2 + dx^3$						
, i	1	2	3	4	5	
х	.8	1	1.2	1.4	1.6	
У	24	20	10	13	12	

## Solution:

 $y = 47.94 - 9.76x - 41.07x^{2} + 20.83x^{3}$   $R^{2} = 0.87$ For x = 1,  $\hat{y} = 17.94$ For x = 1.4,  $\hat{y} = 10.94$ 

Keystrokes:	Display:		
XEQ ALPHA SIZE ALPHA 045			
XEQ ALPHA SPOLYC ALPHA	<b>SPOLYC</b>		
.8 ENTER+) 24 A			
1 ENTER+) 20 A			
1.3 ENTER+ 10 A			
1.3 ENTER+ 10 C			
1.2 ENTER+ 10 A			
1.4 ENTER+) 13 A			
1.6 ENTER+ 12 A	5.00		
E	R2=0.87		
R/S	a=47.94		
R/S	b=-9.76		
R/S	c=-41.07		
R/S	d=20.83		
1 <b>R/S</b>	Y.=17.94		
1.4 <b>R/S</b>	Y.=10.94		

				<b>SIZE</b> : 045
STEP	INSTRUCTIONS	INPUT	FUNCTION	DISPLAY
1.	Parabolic Regression Initialize the program.		xeo ΣPOLYP	ΣΡΟLΥΡ
2.	Repeat step $2 \sim 3$ for $i=1,2,,n$ . Input: $x_i$ $y_i$	Xi Vi	ENTER+) A	(i)
3.	If you made a mistake in inputting $x_k$ and $y_k$ , then correct by	Х <sub>к</sub> Ук	ENTER+) C	(k-1)
4.	Calculate R <sup>2</sup> and regression coefficients a,b, and c		E R/S R/S R/S	R2=(R <sup>2</sup> ) a=(a) b=(b) c=(c)
5.	Calculate estimated y from regression. Input x.	x	R/S	Y.=(ŷ)
6. 7.	Repeat step 5 for different x's. To recall sums in calculation: $\Sigma x_i$ $\Sigma x_i^2$ $\Sigma x_i^3$ $\Sigma x_i^4$ $\Sigma y_i$ $\Sigma x_i^2 y_i$ $\Sigma x_i^2 y_i$		RCL 32 RCL 33 RCL 34 RCL 37 RCL 41 RCL 42 RCL 43	$\begin{array}{c} (\Sigma x_{i}) \\ (\Sigma x_{i}^{2}) \\ (\Sigma x_{i}^{3}) \\ (\Sigma x_{i}^{4}) \\ (\Sigma y_{i}) \\ (\Sigma x_{i} y_{i}) \\ (\Sigma x_{i} y_{i}) \end{array}$
8.	Repeat step 4 if you want the results again.			
9.	To use the program for another set of data, initialize the program by $\rightarrow$ then go to step 2.		<b>—</b> (A)	ΣΡΟLΥΡ

## Example 2:

For the following set of data, perform a parabolic regression, i.e., find suitable coefficients for:

 $y = a + bx + cx^2$ 

Ţ	1	2	3	4	5	6	7
х	1	2	3	4	5	6	7
у	5	12	34	50	75	84	128

## Solution:

 $y = -4.00 + 6.64x + 1.64x^{2}$   $R^{2} = 0.98$ For x = 2,  $\hat{y} = 15.86$ For x = 4,  $\hat{y} = 48.86$ 

Key	stro	kes:
-----	------	------

**Display:** 

XEQ ALPHA SIZE ALPHA 045	
	ZPULTP
2 ENTER+ 12 A	
3 ENTER+ 34 A	
4 ENTER+ 50 A	
5 ENTER+ 75 A	
6 ENTER+ 84 A	
7 ENTER+) 128 A	7.00
E	R2=0.98
R/S	a=-4.00
R/S	b=6.64
R/S	c=1.64
2 <b>R</b> / <b>S</b>	Y.=15.86
4 <b>R</b> /S	Y.=48.86

#### t **STATISTICS**

#### **Paired t Statistic**

Given a set of paired observations from two normal populations with means  $\mu_1$ ,  $\mu_2$  (unknown)

let

$$D_{i} = x_{i} - y_{i}$$

$$\overline{D} = \frac{1}{n} \sum_{i=1}^{n} D_{i}$$

$$s_{D} = \sqrt{\frac{\Sigma D_{i}^{2} - \frac{1}{n} (\Sigma D_{i})^{2}}{n - 1}}$$

The test statistic

$$t = \frac{\overline{D}}{s_{D}} \cdot \sqrt{n}$$

which has n - 1 degrees of freedom (df) can be used to test the null hypothesis

$$H_0: \mu_1 = \mu_2$$

#### **Reference:**

Statistics in Research, B. Ostle, Iowa State University Press, 1963.

### t Statistic For Two Means

Suppose  $\{x_1, x_2, ..., x_{n1}\}$  and  $\{y_1, y_2, ..., y_{n2}\}$  are independent random samples from two normal populations having means  $\mu_1$ ,  $\mu_2$  (unknown) and the same unknown variance  $\sigma^2$ .

We want to test the null hypothesis

$$\mathbf{H}_0: \boldsymbol{\mu}_1 - \boldsymbol{\mu}_2 = \mathbf{d}$$

Define

$$\overline{\mathbf{x}} = \frac{1}{n_1} \sum_{i=1}^{n_1} \mathbf{x}_i$$

$$\overline{y} = \frac{1}{n_2} \sum_{i=1}^{n_2} y_i$$

$$t = \frac{\overline{x} - \overline{y} - d}{\sqrt{\frac{1}{n_1} + \frac{1}{n_2}}} \sqrt{\frac{\Sigma x_i^2 - n_1 \overline{x}^2 + \Sigma y_i^2 - n_2 \overline{y}^2}{n_1 + n_2 - 2}}$$

We can use this t statistic which has the t distribution with  $n_1 + n_2 - 2$  degrees of freedom (df) to test the null hypothesis  $H_0$ .

#### **Reference:**

Statistical Theory and Methodology in Science and Engineering, K.A. Brownlee, John Wiley & Sons, 1965.

				<b>SIZE</b> : 015
STEP	INSTRUCTIONS	INPUT	FUNCTION	DISPLAY
	Paired t Statistic			
1.	Initialize the program.		XEQ <u>SPTST</u>	ΣPTST
2.	Repeat step 2~3 for i=1,2,,n.			
	Input: x <sub>i</sub>	X <sub>i</sub>		(i)
0	yi If you made a mistaka in innutting	yri I	(A)	(1)
J.	x, and $v_{\mu}$ , then correct by	Xĸ		
		Уĸ	C	(k-1)
4.	To calculate the test statistic:			
	D		E	DBAR = (D)
	s <sub>D</sub> t		R/S	T=(t)
	df		R/S	DF=(df)
5.	Repeat step 4 if you want the results again.			
6.	To use the same program for another set of data, initialize the program by $\rightarrow$ then go to step 2.			ΣPTST
	t Statistic for Two Means			
7.	Initialize the program.		XEQ STSTAT	ΣΤSTAT
8.	Repeat step 8~9 for i=1,2,,n1 Input $x_i$ .	x,	A	(i)
9.	If you made a mistake in inputting $x_{\boldsymbol{k}},$ then correct by	Xĸ	C	(k–1)
10.	Initialize for the 2 <sup>nd</sup> array of data		R/S	0.00
11.	Repeat step 11 $\sim$ 12 for j=1, 2, . $n_{2^{*}}$ Input $y_{j}.$	, Уі	A	(j)
12.	If you made a mistake in inputting $\boldsymbol{y}_{\rm h},$ then correct by	У'n	C	(h-1)
13.	Input d to calculate test statistic:			<b>T</b> (1)
	t df	d	E R/S	T = (t) DF=(df)
14.	Repeat step 13 if you want to calculate the test statistic for a different value of d.			
15.	To use the same program for another set of data, initialize the program by $\rightarrow$ then go to step 8.		A	STSTAT
Exam	ple 1:			

r

٦

Xi	14	17.5	17	17.5	15.4
y <sub>i</sub>	17	20.7	21.6	20.9	17.2

 $\overline{D} = -3.20$   $s_D = 1.00$  t = -7.16df = 4.00

# **Keystrokes:**

Display:

ΣPTST
5.00
DBAR =-3.20
SD=1.00
T=-7.16
DF=4.00

# Example 2:

# **Keystrokes:**

Display:

**ΣTSTAT** 

8.00 0.00

10.00 T=1.73 DF=16.00

XEQ ALPHA SIZE ALPHA 015 XEQ ALPHA STSTAT ALPHA
79 A 84 A
99 A 99 C 108 A
114 A 120 A
103 A 122 A 120 A
R/S
91 A 103 A 90 A 113 A
108 A 87 A 100 A 80 A
99 A 54 A
0 E
R/S

# **CHI-SQUARE EVALUATION**

This program calculates the value of the  $\chi^2$  statistic for the goodness of fit test by the equation

$$\chi_1^2 = \sum_{i=1}^n \frac{(O_i - E_i)^2}{E_i}$$
 with df = n - 1

where

 $O_i$  = observed frequency

 $E_i$  = expected frequency

n = number of classes

If the expected values are equal

$$\left( E = E_i = \frac{\Sigma O_i}{n} \text{ for all } i \right)$$

then

$$\chi_2^2 = \frac{n\Sigma O_i^2}{\Sigma O_i} - \Sigma O_i$$

#### **Remarks:**

 In order to apply the goodness of fit test to a set of given data, combining some classes may be necessary to make sure that each expected frequency is not too small (say, not less than 5).

### **Reference:**

Mathematical Statistics, J.E. Freund, Prentice Hall, 1962.

				<b>SIZE</b> : 008
STEP	INSTRUCTIONS	INPUT	FUNCTION	DISPLAY
	Unequal Expected Frequency			
1.	Initialize the program.		XEQ <b>SXSQEV</b>	ΣXSQEV
2.	$\begin{array}{l} \mbox{Repeat step } 2{\sim}3 \mbox{ for } i{=}1{,}2{,}{,}n. \\ \mbox{Input: } 0_i \\ E_i \end{array}$	Oi Ei	(ENTER+)	(i)
3.	If you made a mistake in inputting $O_{\rm k}$ and $E_{\rm k},$ then correct by	0ĸ Eĸ	ENTER•) C	(k-1)
4.	Calculate $\chi_1^2$ .		E	$XSQ = (\chi_1^2)$
5.	To use the same program for another set of data, initialize the program by $\rightarrow$ then go to step 2.			ΣXSQEV
	Equal Expected Frequency			
6.	Initialize the program.		XEQ <u>SEEFXSQ</u>	ΣEEFXSQ
7.	Repeat step 7~8 for $i=1,2,,n$ . Input: $O_i$	0 i	A	(i)
8.	If you made a mistake in inputting $O_n,$ then correct by	0 <sub>h</sub>	C	(h-1)
9.	Calculate: $\chi^2_2$ E		E R/S	$\begin{array}{c} XSQ = (\chi_2^2) \\ E = (E) \end{array}$
10.	Repeat step 9 if you want the results again.			
11.	To use the same program for another set of data, initialize the program by $\rightarrow$ then go to step 7.			ΣEEFXSQ

# Examples 1:

Find the value of  $\chi^2$  statistic for the goodness of fit for the following data set:

Keystrokes:	Display:
XEQ ALPHA SIZE ALPHA 008	
XEQ ALPHA SXSQEV ALPHA	<b><i>SXSQEV</i></b>
8 ENTER+) 9.6 A	
50 ENTER+ 46.75 A	
47 ENTER• 51.85 A	
56 ENTER+) 54.4 A	
5 ENTER+) 8.25 A	
88 ENTER+ 88 A	
88 ENTER+) 88 C	
14 ENTER+) 9.15 A	6.00
E	XSQ=4.84

## Example 2:

The following table shows the observed frequencies in tossing a die 120 times.  $\chi^2$  can be used to test if the die is fair.

Note: Assume that the expected frequencies are equal.

number	1	2	3	4	5	6	
frequency O <sub>i</sub>	25	17	15	23	24	16	
	$\chi_2^2$ E	e = 5. = 20.	00 00				

Since 5.00 is less than 11.07, the data does not support the statement that the die is "unfair" (5% significance level).

Keystrokes:	Display:
XEQALPHASIZEALPHA008XEQALPHASEEFXSQALPHA	Σ <b>EEFXSQ</b>
25 A 17 A 15 A 22 A 22 C	
23 A 24 A 16 A	6.00
E	XSQ=5.00
R/S	E=20.00

# **CONTINGENCY TABLE**

Contingency tables can be used to test the null hypothesis that two variables are independent.

This program calculates the  $\chi^2$  statistic for testing the independence of the two variables. Also Pearson's coefficient of contingency C<sub>c</sub>, which measures the degree of association between the two variables, is calculated.

j	1	2	 k	Totals
1	X <sub>11</sub>	X <sub>12</sub>	 X <sub>1k</sub>	R <sub>1</sub>
2	X <sub>21</sub>	X <sub>22</sub>	 X <sub>2k</sub>	R <sub>2</sub>
Totals	C <sub>1</sub>	C <sub>2</sub>	 Cĸ	Т

## 2 x k CONTINGENCY TABLE

#### 3 x k CONTINGENCY TABLE

÷

.

j	1	2	 k	Totals
1	<b>X</b> <sub>11</sub>	X <sub>12</sub>	 X <sub>1k</sub>	R <sub>1</sub>
2	X <sub>21</sub>	X <sub>22</sub>	 X <sub>2k</sub>	R <sub>2</sub>
3	<b>X</b> <sub>31</sub>	<b>X</b> <sub>32</sub>	 X <sub>3k</sub>	R <sub>3</sub>
Totals	C <sub>1</sub>	C <sub>2</sub>	 Cĸ	Т

## **Equations:**

Row sum 
$$R_i = \sum_{j=1}^{k} x_{ij}$$
  $i = 1, 2 \text{ (for } 2 \times k)$   
 $i = 1, 2, 3 \text{ (for } 3 \times k)$ 

Column sum 
$$C_j = \sum_{i=1}^{n} x_{ij}$$
  $j = 1, 2, ..., k$   
 $n = 2 (for 2 \times k)$   
 $n = 3 (for 3 \times k)$ 

Total T = 
$$\sum_{i=1}^{n} \sum_{j=1}^{k} x_{ij}$$
  $n = 2 (for 2 \times k)$   
 $n = 3 (for 3 \times k)$ 

Chi-square statistic

$$\chi^{2} = \sum_{i=1}^{n} \sum_{j=1}^{k} \frac{(x_{ij} - E_{ij})^{2}}{E_{ij}} \text{ with } df = (n - 1) (k - 1)$$

$$= T \left( \sum_{i=1}^{n} \sum_{j=1}^{k} \frac{x_{ij}^{2}}{R_{i} C_{j}} \right) - T \qquad \begin{array}{c} n = 2 (\text{for } 2 \times k) \\ n = 3 (\text{for } 3 \times k) \end{array}$$

Contingency coefficient

$$C_{\rm c} = \sqrt{\frac{\chi^2}{T + \chi^2}}$$

## **Reference:**

B. Ostle, Statistics in Research, Iowa State University Press, 1972.

				<b>SIZE:</b> 015
STEP	INSTRUCTIONS	INPUT	FUNCTION	DISPLAY
	2×k			
1.	Initialize the program.		xeq ΣCTKK	ΣСТКК
2.	Repeat step 2 $\sim$ 5 for j=1,2,,k. input: $x_{1j}$ $x_{2j}$	X <sub>1j</sub> X <sub>2j</sub>	ENTER+) A	(j)
3.	(Optional) Calculate column sum C <sub>j</sub> .		R/S	$CS = (C_i)$
4.	If you made a mistake in inputting $x_{1h}$ and $x_{2h}$ , then correct by	X <sub>1h</sub> X <sub>2h</sub>	ENTER+) C	(h-1)
5.	(Optional) Calculate column sum C <sub>n</sub> (correction).		R/S	$CS = (-C_h)$
6.	Go to step 12 for contingency table calculations.			
	3×k			
7.	Initialize the program.		XEQ SCTKKK	ΣСТККК
8.	Repeat step 8~11 for j=1,2,,k. input: $x_{1j}$	X <sub>1j</sub>		
	X <sub>2j</sub> X <sub>3i</sub>	Х <sub>2j</sub> Х <sub>3i</sub>	A	(i)
9.	(Optional) Calculate column sum C <sub>j</sub> .		R/S	$CS = (C_j)$
10.	If you made a mistake in inputting			
	$x_{1h}$ , $x_{2h}$ , and $x_{3h}$ , then correct by	X <sub>1h</sub>		
		X <sub>2h</sub> X <sub>3h</sub>	C	(h-1)

STEP	INSTRUCTIONS	INPUT	FUNCTION	DISPLAY
11.	(Optional) Calculate column sum $C_h$ (correction).		R/S	$CS = (-C_h)$
12.	Calculate: Test statistic $\chi^2$ Coefficient C <sub>c</sub> Row sum 1 R <sub>1</sub> Row sum 2 R <sub>2</sub> Row sum 3 R <sub>3</sub> (3×k only) Total T		E R/S R/S R/S R/S R/S	$XSQ = (\chi^{2})$ $CC = (C_{c})$ $R1 = (R_{1})$ $R2 = (R_{2})$ $R3 = (R_{3})$ T = (T)
13.	Repeat step 12 if you want the results again.			
14.	To use the same program for another set of data, initialize by $\rightarrow$		<b>a</b>	ΣCTKK or ΣCTKKK
15.	then go to step 2 or step 8. To use the other program, go to			
	step 1 or step 7.			

# Example 1:

Find the test statistic  $\chi^2$  and coefficient of contingency C<sub>c</sub> for the following set of data.

	1	2	3
A	2	5	4
В	3	8	7

Keystrokes:	Display:
XEQ ALPHA SIZE ALPHA 015	
XEQ ALPHA SCTKK ALPHA	ΣCTKK
2 ENTER+ 3 A	1.00
R/S	CS=5.00
5 ENTER+ 8 A 4 ENTER+ 7 A	3.00
E	XSQ=0.02
R/S	CC=0.03
R/S	R1=11.00
R/S	R2=18.00
R/S	T=29.00

olay:

# Example 2:

Find test statistic  $\chi^2$  and coefficient of contingency  $C_{\rm c}$  for the following set of data.



# **Keystrokes:**

Display:

XEQ ALPHA SIZE ALPHA 015	
ΧΕΟ ΑLPHA ΣCTKKK ΑLPHA	ΣCTKKK
36 ENTER+ 31 ENTER+ 58 A	1.00
[R/S]	CS=125.00
67 ENTER+) 60 ENTER+) 87 A	
4 ENTER+) 49 ENTER+) 80 A	
4 ENTER+ 49 ENTER+ 80 C	
49 [ENTER+] 49 [ENTER+] 80 [A]	
58 ENTER+ 54 ENTER+ 68 A	4.00
E	XSQ=3.36
R/S	CC=0.07
R/S	R1=210.00
R/S	R2=194.00
(R/S)	R3=293.00
	T=697.00

## SPEARMAN'S RANK CORRELATION COEFFICIENT

Spearman's rank correlation coefficient is a measure of rank correlation under the following circumstance: n individuals are ranked from 1 to n according to some specified characteristic by 2 observers, and we wish to know if the 2 rankings are substantially in agreement with one another.

Spearman's rank correlation coefficient is defined by

$$r_{s} = 1 - \frac{6\sum_{i=1}^{n} D_{i}^{2}}{n(n^{2} - 1)}$$

where  $n = number of paired observations (x_i, y_i)$  $D_i = rank (x_i) - rank (y_i) = R_i - S_i$ 

If the X and Y random variables from which these n pairs of observations are derived are independent, then  $r_s$  has zero mean and a variance equal to

$$\frac{1}{n-1}$$

A test for the null hypothesis

 $H_0$ : X, Y are independent

is made using

$$z = r_s \sqrt{n-1}$$

which is approximately a standardized normal variable (for large n, say  $n \ge 10$ ).

If the null hypothesis of independence is not rejected, we can infer that the population correlation coefficient  $\rho(x, y) = 0$ , but dependence between the variables does not necessarily imply that  $\rho(x, y) \neq 0$ .

#### Note:

 $-1 \le r_s \le 1$  $r_s = 1$  indicates complete agreement in order of the ranks and  $r_s = -1$  indicates complete agreement in the opposite order of the ranks.

#### **Reference:**

Nonparametric Statistical Inference, J. D. Gibbons, McGraw Hill, 1971.

-

				<b>SIZE</b> : 003
STEP	INSTRUCTIONS	INPUT	FUNCTION	DISPLAY
1.	Initialize the program.		xeq ΣSPEAR	ΣSPEAR
2.	$\begin{array}{l} \mbox{Repeat step } 2{\sim}3 \mbox{ for } i{=}1{,}2{,}{,}n. \\ \mbox{Input: } R_i \\ S_i \end{array}$	R <sub>i</sub> S <sub>i</sub>	(ENTER+)	(i)
3.	If you made a mistake in inputting $R_{\rm k}$ and $S_{\rm k},$ then correct by	R⊾ S⊾	ENTER+) C	(k-1)
4.	Calculate: r <sub>s</sub> z		E R/S	$RS=(r_s)$ Z=(z)
5.	Repeat step 4 if you want the results again.			
6.	For another set of data, initialize the program by $\rightarrow$ then go to step 2.			ΣSPEAR

# **Example:**

The following data set is the result of two tests in a class; find  $r_{s} \mbox{ and } z.$ 

			-	
	Xi	Уi	К <sub>і</sub>	S <sub>i</sub>
Student	Math Grade	Stat Grade	Rank of x <sub>i</sub>	Rank of y <sub>i</sub>
1	82	81	6	7
2	67	75	14	11
3	91	85	3	4
4	98	90	1	2
5	74	80	11	8
6	52	60	15	15
7	86	94	4	1
8	95	78	2	9
9	79	83	9	6
10	78	76	10	10
11	84	84	5	5
12	80	69	8	13
13	69	72	13	12
14	81	88	7	3
15	73	61	12	14
	1	1	1	

64 Spearman's Rank Correlation Coefficient

Keystrokes:	Display:
XEQ ALPHA SIZE ALPHA 003	
XEQ ALPHA SSPEAR ALPHA	ΣSPEAR
6 ENTER+) 7 A	
14 ENTER+) 11 A	
3 ENTER+ 4 A	
1 ENTER+ 2 A	
11 ENTER+ 8 A	
5 ENTER+ 5 A	
5 ENTER+ 5 C	
15 ENTER+ 15 A	
4 ENTER+ 1 A	
2 ENTER+ 9 A	
9 ENTER+ 6 A	
10 ENTER+ 10 A	
5 ENTER+ 5 A	
8 ENTER+) 13 A	
13 ENTER+ 12 A	
7 ENTER+ 3 A	
12 [ENTER+] 14 A	15.00
E	RS=0.76
R/S	Z=2.85

Notes

### NORMAL AND INVERSE NORMAL DISTRIBUTION

This program evaluates the standard normal density function f(x) and the normal integral Q(x) for given x. If Q is given, x can also be found. The standard normal distribution has mean 0 and standard deviation 1.

### **Equations:**

1. Standard normal density

$$f(x) = \frac{1}{\sqrt{2\pi}} e^{-\frac{x^2}{2}}$$



2. Normal integral

$$Q(x) = \frac{1}{\sqrt{2\pi}} \int_{-x}^{\infty} e^{-\frac{t^2}{2}} dt$$

Polynomial approximation is used to calculate Q(x) for given x.

Define R = f(x)  $(b_1t + b_2t^2 + b_3t^3 + b_4t^4 + b_5t^5) + \epsilon(x)$ where  $|\epsilon(x)| < 7.5 \times 10^{-8}$ 

$$t = \frac{1}{1 + r |x|}$$
,  $r = 0.2316419$ 

$$b_1 = .319381530, \qquad b_2 = -.356563782$$
  

$$b_3 = 1.781477937, \qquad b_4 = -1.821255978$$
  

$$b_5 = 1.330274429$$

Then Q(x) = 
$$\begin{cases} R & \text{if } x \ge 0\\ 1 - R & \text{if } x < 0 \end{cases} \text{ with error } |\epsilon(x)| < 7.5 \times 10^{-8} \end{cases}$$

3. Inverse normal

For a given 0 < Q < 1, x can be found such that

$$Q = \frac{1}{\sqrt{2\pi}} \int_{x}^{\infty} e^{-\frac{t^2}{2}} dt$$

The following rational approximation is used:

Define y = t - 
$$\frac{c_0 + c_1 t + c_2 t^2}{1 + d_1 t + d_2 t^2 + d_3 t^3} + \epsilon(Q)$$

where  $|\epsilon(Q)| < 4.5 \times 10^{-4}$ 

$$t = \begin{cases} \sqrt{\ln \frac{1}{Q^2}} & \text{if } 0 < Q \le 0.5 \\ \sqrt{\ln \frac{1}{(1-Q)^2}} & \text{if } 0.5 < Q < 1 \end{cases}$$

 $\begin{array}{ll} c_0 = 2.515517 & d_1 = 1.432788 \\ c_1 = 0.802853 & d_2 = 0.189269 \\ c_2 = 0.010328 & d_3 = 0.001308 \end{array}$ 

Then x =  $\begin{cases} y & \text{if } 0 < Q \leq 0.5 \\ -y & \text{if } 0.5 < Q < 1 \end{cases}$  with error  $|\epsilon(Q)| < 4.5 \times 10^{-4}$ 

#### **Reference:**

Abramowitz and Stegun, *Handbook of Mathematical Functions*, National Bureau of Standards, 1970.

				<b>SIZE</b> : 019
STEP	INSTRUCTIONS	INPUT	FUNCTION	DISPLAY
1.	Initialize the program.		xeq SNORMD	ΣNORMD
2.	Input x to calculate f(x).	х	С	F = (f(x))
3.	Input x to calculate Q(x).	х	E	Q = (Q(x))
4.	Input $Q(x)$ to calculate x.	Q(x)	A	X = (x)
5.	Repeat any of the above steps if desired.			

# Example 1:

Find f(x) and Q(x) for x = 1.18 and x = -2.28.

Keystrokes:	Display:
XEQ ALPHA SIZE ALPHA 019	
XEQ ALPHA SNORMD ALPHA	2NORMD
1.18 C	F=0.20
1.18 E	Q=0.12
2.28 CHS E	Q=0.99
2.28 CHS C	F=0.03

# Example 2:

Given Q = 0.12 and Q = 0.95, find x.

(If you have run through Example 1, then you can proceed; otherwise you have to initialize the program as described in Example 1).

Display:	
X=1.18	
X=-1.65	

Notes

### **CHI-SQUARE DISTRIBUTION**

This program evaluates the chi-square density

$$f(x) = \frac{1}{2^{\frac{\nu}{2}} \Gamma\left(\frac{\nu}{2}\right)} x^{\frac{\nu}{2} - 1} e^{-\frac{x}{2}}$$

where  $x \ge 0$ 

 $\nu$  is the degrees of freedom.



Series expansion is used to evaluate the cumulative distribution

$$P(x) = \int_{0}^{x} f(t) dt$$
$$= \left(\frac{x}{2}\right)^{-\frac{\nu}{2}} \frac{e^{-\frac{x}{2}}}{\Gamma\left(\frac{\nu+2}{2}\right)} \left[1 + \sum_{k=1}^{\infty} \frac{x^{k}}{(\nu+2)(\nu+4)\dots(\nu+2k)}\right]$$

The program calculates successive partial sums of the above series. When two consecutive partial sums are equal, the value is used as the sum of the series.
### **Remarks:**

- Program requires  $\nu < 141$ . If  $\nu > 141$ , erroneous overflow will result.
- If both x and v are large, f(x) may result in an overflow error.
- If  $\nu$  is even,

$$\Gamma\left(\frac{\nu}{2}\right) = \left(\frac{\nu}{2} - 1\right) \, !$$

If  $\nu$  is odd,

$$\Gamma\left(\frac{\nu}{2}\right) = \left(\frac{\nu}{2} - 1\right) \left(\frac{\nu}{2} - 2\right) \dots \left(\frac{1}{2}\right) \Gamma\left(\frac{1}{2}\right)$$
$$\Gamma\left(\frac{1}{2}\right) = \sqrt{\pi}$$

#### **Reference:**

Abramowitz and Stegun, *Handbook of Mathematical Functions*, National Bureau of Standards, 1970.

				<b>SIZE</b> : 007
STEP	INSTRUCTIONS	INPUT	FUNCTION	DISPLAY
1.	Initialize the program.		XEQ SCHISQD	ΣCHISQD
2.	Input degrees of freedom $\nu_{\rm e}$	ν	A	$(\Gamma( u/2))$
3.	Input x to calculate f(x).	х	C	F = (f(x))
4.	Input x to calculate P(x).	x	E	P = (P(x))
5.	Repeat step 3 or step 4 if desired.			
6.	For a different $\nu$ , go to step 2.			

### 72 Chi-Square Distribution

# **Examples:**

- 1. If degrees of freedom  $\nu = 20$ , find f(x), P(x) for x = 9.6 and x = 15.
- 2. If  $\nu = 3$ , find f(x) and P(x) for x = 7.82.

## **Keystrokes:**

Display:

XEQ ALPHA SIZE ALPHA 007	
$\begin{array}{c} \textbf{XEQ}  \textbf{ALPHA}  \boldsymbol{\Sigma} \textbf{CHISQD}  \textbf{ALPHA} \end{array}$	ΣCHISQD
20 🔺	362880.00
9.6 C	F=0.02
9.6 E	P=0.03
15 E	P=0.22
15 C	F=0.06
3 🔺	0.89
7.82 C	F=0.02
7.82 E	P=0.95

# APPENDIX A PROGRAM DATA

		#REG. TO	DATA		DISPLAY
	PROGRAM	СОРУ	REGISTERS		FORMAT
÷	Basic Statistics for Two Variables	50	$00 \sim 11$	$00 \sim 03, 21, 27, 29$	FIX 2
2	Moments. Skewness, and Kurtosis	36	00 ~ 11	$00 \sim 03, 21, 27, 29$	FIX 2
ဂဲ	Analysis of Variance (One Way)	29	$00 \sim 19$	$00 \sim 03, 21, 27, 29$	FIX 2
4	Analysis of Variance (Two Way)	S	$00 \sim 17$	$00 \sim 03, 21, 27, 29$	FIX 2
<u>ى</u>	Analysis of Covariance (One Way)	60	$00 \sim 25$	$00 \sim 03, 21, 27, 29$	FIX 2
ю	Curve Fitting	8	$00 \sim 15$	$00 \sim 03, 21, 27, 29$	FIX 2
۲.	Multiple Linear Regression	157	00 ~ 44	$00 \sim 03, 21, 27, 29$	FIX 2
œ	Polynomial Regression	102	00 ~ <del>4</del>	$00 \sim 03, 21, 27, 29$	FIX 2
<u>ю</u>	t Statistics	29	$00 \sim 14$	$00 \sim 03, 21, 27, 29$	FIX 2
₽.	Chi-Square Evaluation	21	$00 \sim 07$	$00 \sim 03, 21, 27, 29$	FIX 2
÷.	Contingency Table	ŝ	$00 \sim 14$	$00 \sim 03, 21, 27, 29$	FIX 2
₽	Spearman's Rank Correlation Coefficient	13	00 ~ 02	$00 \sim 03, 21, 27, 29$	FIX 2
₽. 13	Normal and Inverse Normal Distribution	47	$00 \sim 18$	$00 \sim 03, 21, 27, 29$	FIX 2
14.	Chi-Square Distribution	21	$90 \sim 00$	$00 \sim 03, 21, 27, 29$	FIX 2

# APPENDIX B PROGRAM LABELS

*A	ΣΑΟΥΤWΟ	ΣLIN	ΣPOLYP
жB	ΣΒSΤΑΤ	ΣLOG	ΣΡΟΨ
жве	ΣBSTG	ΣMLRXY	ΣPTST
жС	ΣCHISQD	ΣMLRXYZ	ΣSPEAR
*MD	ΣСТКК	ΣMMTGO	<b>ΣTSTAT</b>
*MT	ΣСТККК	ΣΜΜΤUG	ΣΧSQEV
ΣΑΝΟϹΟΥ	ΣΕΕΓΧSQ	ΣNORMD	
ΣΑΟVONE	ΣΕΧΡ	ΣPOLYC	

The labels in this list are not in the same order as they appear in the catalog listing for the module.

Notes



Portable Computer Division 1000 N.E. Circle Blvd., Corvallis, OR 97330, U.S.A.

European Headquarters 150, Route du Nant-d'Avril P.O. Box, CH-1217 Meyrin 2 Geneva-Switzerland HP-United Kingdom (Pinewood) Gb-Nine Mile Ride, Wokingham Berkshire RG11 3LL

00041-90030 Rev. E English-IK-IPC

Printed in Singapore 8/84