

HEWLETT  PACKARD

**HP-65**  
**STAT PAC 1**

JANUARY, 1975

The program material contained herein is supplied without representation or warranty of any kind. Hewlett-Packard Company therefore assumes no responsibility and shall have no liability, consequential or otherwise, of any kind arising from the use of this program material or any part thereof.

## CONTENTS

Introduction . . . . .	3
Format of User Instructions . . . . .	4
Entering a Program . . . . .	7
1. Mean, Standard Deviation, Standard Error . . . . .	8
2. Mean, Standard Deviation, Standard Error (Grouped Data) . . . . .	10
3. Permutation and Combination . . . . .	12
4. Arithmetic, Geometric, Harmonic and Generalized Means . . . . .	14
5. Sums for Two Variables . . . . .	16
6. Basic Statistics (Two Variables) . . . . .	18
7. Moments, Skewness and Kurtosis (For Grouped or Ungrouped Data) . . . . .	20
8. Random Number Generator . . . . .	22
9. Analysis of Variance (One Way) . . . . .	24
10. Normal Distribution . . . . .	26
11. Inverse Normal Integral . . . . .	28
12. Chi-Square Distribution . . . . .	30
13. t Distribution . . . . .	32
14. F Distribution . . . . .	34
15. Bivariate Normal Distribution . . . . .	36
16. Logarithmic Normal Distribution . . . . .	38
17. Weibull Distribution . . . . .	40
18. Binomial Distribution . . . . .	42
19. Negative Binomial Distribution . . . . .	44
20. Hypergeometric Distribution . . . . .	46
21. Poisson Distribution . . . . .	48
22. Linear Regression . . . . .	49
23. Exponential Curve Fit . . . . .	52
24. Power Curve Fit . . . . .	54
25. Logarithmic Curve Fit . . . . .	56
26. Least Squares Regression of $y = cx^a + dx^b$ . . . . .	58
27. Multiple Linear Regression . . . . .	60
28. Parabolic Curve Fit . . . . .	62
29. Paired t Statistic . . . . .	64
30. t Statistic for Two Means . . . . .	66
31. Chi-Square Evaluation . . . . .	68
32. 2 x k Contingency Table . . . . .	70
33. Bartlett's Chi-Square Statistic . . . . .	72
34. Spearman's Rank Correlation Coefficient . . . . .	74
35. Mann-Whitney Statistic . . . . .	76
36. Kendall's Coefficient of Concordance . . . . .	78
37. Biserial Correlation Coefficient . . . . .	80
Program Listings . . . . .	83-123



## INTRODUCTION

Programs for your HP-65 Stat Pac 1 have been selected from the areas of general statistics, distribution functions, curve fitting and test statistics.

Each program includes a general description, formulas used in the program solution, numerical examples, and user instructions. Program listings and register allocations are given in the back of the Pac.

Some related individual programs were combined on one card when it seemed they might be useful together. In this way more programs could be included in the Pac.

We hope you find the HP-65 Stat Pac 1 a useful tool for your computational work, and welcome your comments, requests and suggestions—these are our most important source of future user-oriented programs.

#### 4 Format of User Instructions

## FORMAT OF USER INSTRUCTIONS

The following is an example of a set of User Instructions.

LINE	INSTRUCTIONS	DATA	KEYS	DISPLAY
1	Enter program			
2	Clear registers		A	
3	Perform 3–4 for i=1, ..., n	a <sub>i</sub>	↑	
4		b <sub>i</sub>	B	
5			C	Answer
	(To run a new case, go to 2)			

To follow the instructions, start with line 1 and read from left to right, performing indicated operations as you proceed. Lines having no numbers contain special notes to the user and are inside parentheses in the INSTRUCTIONS column. The message “To run a new case, go to 2” following line 5 in the above example is a special note.

Lines are read in sequential order except where the INSTRUCTIONS column directs otherwise. For example, “go to 2” means to jump to line 2. Repeated processes—used in most cases for a long string of input/output data—are outlined with a bold border together with a “Perform” instruction. In the above example, “Perform 3–4 for i = 1, ..., n” means to execute the loop (line 3 and line 4) n times. The first time, the dummy variable i takes the value 1; the second time i takes the value 2; etc.

Normally, as in the above example, the first instruction is “Enter program” which means load the preprogrammed magnetic card (for instructions of loading a card, see “Entering A Program” on P. 7). Some instructions are self-contained and can be carried out by just reading the INSTRUCTIONS column alone, e.g., “Enter program”. But some instructions depend on the information supplied by the DATA and/or KEYS columns. In line 2 of the example above, “Clear registers” appears in the INSTRUCTIONS column and **A** appears in the KEYS column, which means you have to clear the working registers by pressing the **A** key.

The DATA column specifies the input data to be supplied. Invalid arguments which result in division by zero, finding square root of a negative number, etc. will result in flashing zeros. Arguments out of the designated program range will result in incorrect answers or flashing zeros. When a computed value exceeds the calculator range, an overflow or underflow occurs and halts the program.

The KEYS column specifies the keys to be pressed. **↑** is the symbol used to denote the **ENTER** key. All other key designations are identical to those appearing on the HP-65. Ignore any blank positions in the KEYS column.

The DISPLAY column may show counters, intermediate or final results. In line 5 of the example, the answer will be displayed after pressing the **C** key.



## ENTERING A PROGRAM

From the card case supplied with this application pac, select a program card.

Set W/PRGM-RUN switch to RUN.

Turn the calculator ON. You should see 0.00


Gently insert the card (printed side up) in the right, lower slot as shown. When the card is part way in, the motor engages it and passes it out the left side of the calculator. Sometimes the motor engages but does not pull the card in. If this happens, push the card a little farther into the machine. Do not impede or force the card; let it move freely. (The display will flash if the card reads improperly. In this case, press **CLX** and reinsert the card.)



When the motor stops, remove the card from the left side of the calculator and insert it in the upper "window slot" on the right side of the calculator.

The program is now stored in the calculator. It remains stored until another program is entered or the calculator is turned off.



**MEAN, STANDARD DEVIATION, STANDARD ERROR**

Given a set of data points

$$\{x_1, x_2, \dots, x_n\}$$

the program computes the following statistics:

$$\text{mean } \bar{x} = \frac{1}{n} \sum_{i=1}^n x_i$$

$$\text{standard deviation } s_x = \sqrt{\frac{\sum x_i^2 - n\bar{x}^2}{n-1}}$$

$$\left( \text{or } s_x' = \sqrt{\frac{\sum x_i^2 - n\bar{x}^2}{n}} \right)$$

$$\text{standard error of the mean } s_{\bar{x}} = \frac{s_x}{\sqrt{n}}$$

$$\left( \text{or } s_{\bar{x}}' = \frac{s_x'}{\sqrt{n}} \right)$$

**Notes:** 1.  $n$ ,  $\sum x_i$ ,  $\sum x_i^2$  are in registers  $R_1$ ,  $R_2$ ,  $R_3$ .

2. To remove erroneous data, key in that data value and press **E**. “ $\Sigma-$ ” is the operational inverse of “ $\Sigma+$ ”.
3.  $n$  is a positive integer and  $n > 1$ .
4. Due to roundoff errors, flashing zeros may be returned for the standard deviation when it is very small relative to the mean.

**Example:**

The set of numbers  $\{2, 3.4, 7, 11, 23, 3.41\}$  has

$$\bar{x} = 8.30$$

$$s_x = 7.91 \quad s_x' = 7.22$$

$$s_{\bar{x}} = 3.23 \quad s_{\bar{x}}' = 2.95$$

LINE	INSTRUCTIONS	DATA	KEYS	DISPLAY
1	Enter program			
2	Initialize		RTN R/S	
3	Perform 3 for $i = 1, 2, \dots, n$	$x_i$	A	$i$
	(Correct erroneous data $x_k$ )	$x_k$	E	
4	Compute $\bar{x}$		B	$\bar{x}$
5	Compute $s_x$		C	$s_x$
	(optional)		R/S	$s_x'$
6	Compute $s_{\bar{x}}$		D	$s_{\bar{x}}$
	(optional)		R/S	$s_{\bar{x}}'$
	(For a new case, go to 2)			

## MEAN, STANDARD DEVIATION, STANDARD ERROR (GROUPED DATA)



Given a set of data points

$$x_1, x_2, \dots, x_n$$

with respective frequencies

$$f_1, f_2, \dots, f_n$$

the program computes the following statistics:

$$\text{mean } \bar{x} = \frac{\sum f_i x_i}{\sum f_i}$$

$$\text{standard deviation } s_x = \sqrt{\frac{\sum f_i x_i^2 - (\sum f_i) \bar{x}^2}{\sum f_i - 1}}$$

$$\left( \text{or } s_x' = \sqrt{\frac{\sum f_i x_i^2 - (\sum f_i) \bar{x}^2}{\sum f_i}} \right)$$

$$\text{standard error } s_{\bar{x}} = \frac{s_x}{\sqrt{\sum f_i}}$$

$$\left( \text{or } s_{\bar{x}}' = \frac{s_x'}{\sqrt{\sum f_i}} \right)$$

- Notes:**
- $\sum f_i$ ,  $\sum f_i x_i$ ,  $\sum f_i x_i^2$ ,  $n$  are in registers  $R_1$ ,  $R_2$ ,  $R_3$ ,  $R_4$ .
  - To remove erroneous data  $x_k$ ,  $f_k$ :  
 $x_k \quad \boxed{\uparrow} \quad f_k \quad \boxed{E}$   
“ $\Sigma-$ ” is the operational inverse of “ $\Sigma+$ ”.
  - $n$  is a positive integer and  $n > 1$ .

**Example:**

$x_i$	2	3.4	7	11	23	3.41
$f_i$	5	3	4	2	3	1

$$\bar{x} = 7.92$$

$$s_x = 7.52 \quad s_x' = 7.31$$

$$s_{\bar{x}} = 1.77 \quad s_{\bar{x}}' = 1.72$$

LINE	INSTRUCTIONS	DATA	KEYS	DISPLAY
1	Enter program			
2	Initialize		RTN R/S	
3	Perform 3–4 for $i = 1, 2, \dots, n$	$x_i$	$\uparrow$	
4		$f_i$	A	i
	(Correct erroneous data $x_k, f_k$ )	$x_k$	$\uparrow$	
		$f_k$	E	
5	Compute $\bar{x}$		B	$\bar{x}$
6	Compute $s_x$		C	$s_x$
	(optional)		R/S	$s_x'$
7	Compute $s_{\bar{x}}$		D	$s_{\bar{x}}$
	(optional)		R/S	$s_{\bar{x}}'$
	(For a new case, go to 2)			

**PERMUTATION AND COMBINATION**

PERMUTATION AND COMBINATION

**STAT 1-03A** $mP_n$  $mC_n$ 

$$mP_n = \frac{m!}{(m-n)!} = m(m-1) \dots (m-n+1)$$

$$mC_n = \frac{m!}{(m-n)! n!} = \frac{m(m-1) \dots (m-n+1)}{1 \cdot 2 \cdot \dots \cdot n}$$

where  $m, n$  are integers and  $0 \leq n \leq m$ .

**Notes:** 1.  $mP_0 = 1$ ,  $mP_1 = m$ ,  $mP_m = m!$

$$2. \quad mC_0 = mC_m = 1$$

$$3. \quad mC_1 = mC_{m-1} = m$$

$$4. \quad mC_n = mC_{m-n}$$


**Examples:**

1.  ${}_{27}P_5 = 9687600.00$

2.  ${}_{73}C_4 = 1088430.00$

LINE	INSTRUCTIONS	DATA	KEYS	DISPLAY
1	Enter program			
2	Compute ${}_mP_n$	m	↑	
3		n	A	${}_mP_n$
4	Compute ${}_mC_n$	m	↑	
5		n	B	${}_mC_n$

# ARITHMETIC, GEOMETRIC, HARMONIC AND GENERALIZED MEANS



Arithmetic mean

$$A = \frac{a_1 + \dots + a_n}{n}$$

Geometric mean

$$G = (a_1 a_2 \dots a_n)^{\frac{1}{n}}$$

Harmonic mean

$$H = \frac{n}{\frac{1}{a_1} + \frac{1}{a_2} + \dots + \frac{1}{a_n}}$$

Generalized mean

$$M(t) = \left( \frac{1}{n} \sum_{k=1}^n a_k^t \right)^{\frac{1}{t}}$$

**Notes:** 1.  $a_k > 0$ ,  $k = 1, 2, \dots, n$

- 2.  $M(1) = A$
- $M(-1) = H$

**Examples:**

The set of numbers  $\{2, 3.4, 3.41, 7, 11, 23\}$  has


$$A = 8.30$$

$$G = 5.87$$

$$H = 4.40$$

$$M(1) = 8.30$$

LINE	INSTRUCTIONS	DATA	KEYS	DISPLAY
1	Enter program			
2	Initialize		RTN R/S	
3	If $M(t)$ is desired	t	R/S	
4	Perform 4 for $k=1, 2, \dots, n$	$a_k$	A	k
5	Compute A		B	A
6	Compute G		C	G
7	Compute H		D	H
8	Compute $M(t)$		E	$M(t)$

**SUMS FOR TWO VARIABLES**

This program computes sums for a set of given data

$$\{(x_i, y_i), i = 1, 2, \dots, n\}.$$

$n, \Sigma x_i, \Sigma x_i^2, \Sigma y_i, \Sigma y_i^2, \Sigma x_i y_i$  are in registers R<sub>1</sub> through R<sub>6</sub>.

This program can be used in conjunction with *Stat 1-22A, Linear Regression*, to fit a linear regression line or *Stat 1-06A, Basic Statistic (Two Variables)*, to obtain means, standard deviations, covariance and correlation coefficient.

**Example:**

$x_i$	26	30	44	50	62	68	74
$y_i$	92	85	78	81	54	51	40

$$n = 7.00$$

$$\Sigma x_i = 354.00$$

$$\Sigma x_i^2 = 19956.00$$

$$\Sigma y_i = 481.00$$

$$\Sigma y_i^2 = 35451.00$$

$$\Sigma x_i y_i = 22200.00$$

LINE	INSTRUCTIONS	DATA	KEYS	DISPLAY
1	Enter program			
2	Initialize		f REG	
3	Perform 3–4 for $i=1, 2, \dots, n$	$x_i$	$\uparrow$	
4		$y_i$	A	i
	(Correct erroneous data $x_k, y_k$ )	$x_k$	$\uparrow$	
		$y_k$	B	
5			C	n
6			R/S	$\Sigma x_i$
7			R/S	$\Sigma x_i^2$
8			R/S	$\Sigma y_i$
9			R/S	$\Sigma y_i^2$
10			R/S	$\Sigma x_i y_i$
	(To run a new case, go to 2)			

**BASIC STATISTICS (TWO VARIABLES)**

BASIC STATISTICS (TWO VARIABLES)

 $\bar{x}, \bar{y}$  $s_x, s_y$  $s_{xy}$  $r_{xy}$ **STAT 1-06A**

This program must be used in conjunction with *Stat 1–05A, Sums for Two Variables*, to compute means, standard deviations, covariance and correlation coefficient derived from a set of data points

$$\{(x_i, y_i), i = 1, 2, \dots, n\} .$$

$$\text{means } \bar{x} = \frac{1}{n} \sum_{i=1}^n x_i \quad \bar{y} = \frac{1}{n} \sum_{i=1}^n y_i$$

$$\begin{aligned} \text{standard deviations } s_x &= \sqrt{\frac{\sum x_i^2 - n\bar{x}^2}{n-1}} \\ \left( \text{or } s_x' \right) &= \sqrt{\frac{\sum x_i^2 - n\bar{x}^2}{n}} \\ s_y &= \sqrt{\frac{\sum y_i^2 - n\bar{y}^2}{n-1}} \end{aligned}$$

$$\left( \text{or } s_y' = \sqrt{\frac{\sum y_i^2 - n\bar{y}^2}{n}} \right)$$

$$\text{covariance } s_{xy} = \frac{1}{n-1} \left( \sum x_i y_i - \frac{1}{n} \sum x_i \sum y_i \right)$$

$$\left( \text{or } s_{xy}' = \frac{1}{n} \left[ \sum x_i y_i - \frac{1}{n} \sum x_i \sum y_i \right] \right)$$

$$\text{correlation coefficient } r_{xy} = \frac{s_{xy}}{s_x s_y} = \frac{s_{xy}'}{s_x' s_y'}$$

**Note:** n is a positive integer and n > 1.

**Example:**

$x_i$	26	30	44	50	62	68	74
$y_i$	92	85	78	81	54	51	40

$$\bar{x} = 50.57, \quad \bar{y} = 68.71$$

$$s_x = 18.50, \quad s_y = 20.00$$

$$s_x' = 17.13, \quad s_y' = 18.51$$

$$s_{xy} = -354.14$$

$$s_{xy}' = -303.55$$

$$r_{xy} = -0.96$$

LINE	INSTRUCTIONS	DATA	KEYS	DISPLAY
1	Enter program <i>Stat 1–05A</i>			
2	Initialize		f REG	
3	Perform 3–4 for $i = 1, 2, \dots, n$	$x_i$	$\uparrow$	
4		$y_i$	A	i
	(Correct erroneous data $x_k, y_k$ )	$x_k$	$\uparrow$	
		$y_k$	B	
5	Enter program <i>Stat 1–06A</i>			
6			A	$\bar{x}$
7			R/S	$\bar{y}$
8			B	$s_x$
9			R/S	$s_y$
	(optional)		R/S	$s_x'$
	(optional)		R/S	$s_y'$
10			C	$s_{xy}$
	(optional)		R/S	$s_{xy}'$
11			D	$r_{xy}$

## MOMENTS, SKEWNESS AND KURTOSIS (FOR GROUPED OR UNGROUPED DATA)

**MOMENTS, SKEWNESS AND KURTOSIS      STAT 1-07A 1**

$\Sigma+$        $\Sigma-$        $\Sigma+(f_i)$        $\Sigma-(f_i)$



**MOMENTS, SKEWNESS AND KURTOSIS      STAT 1-07A 2**

$\bar{x}$        $m_2$        $m_3$        $m_4$        $\gamma_1, \gamma_2$



This program computes the following statistics for a set of given data  $\{x_1, x_2, \dots, x_n\}$ :

$$1^{\text{st}} \text{ moment} \quad \bar{x} = \frac{1}{n} \sum_{i=1}^n x_i$$

$$2^{\text{nd}} \text{ moment} \quad m_2 = \frac{1}{n} \sum x_i^2 - \bar{x}^2$$

$$3^{\text{rd}} \text{ moment} \quad m_3 = \frac{1}{n} \sum x_i^3 - \frac{3}{n} \bar{x} \sum x_i^2 + 2\bar{x}^3$$

$$4^{\text{th}} \text{ moment} \quad m_4 = \frac{1}{n} \sum x_i^4 - \frac{4}{n} \bar{x} \sum x_i^3 + \frac{6}{n} \bar{x}^2 \sum x_i^2 - 3\bar{x}^4$$

Moment coefficient of skewness

$$\gamma_1 = \frac{m_3}{m_2^{3/2}}$$

Moment coefficient of kurtosis

$$\gamma_2 = \frac{m_4}{m_2^2}$$

This program also provides the option for computing those statistics for grouped data (using similar formulas as for ungrouped data):

data		$y_1$	$y_2$	...	$y_m$	
frequency		$f_1$	$f_2$	...	$f_m$	

**Reference:** Theory and Problems of Statistics, M. R. Spiegel,  
Schaum's Outline, McGraw-Hill, 1961

**Examples:**

1. Ungrouped data

i	1	2	3	4	5	6	7	8	9
$x_i$	2.1	3.5	4.2	6.5	4.1	3.6	5.3	3.7	4.9

$$\bar{x} = 4.21, m_2 = 1.39, m_3 = 0.39, m_4 = 5.49$$

$$\gamma_1 = 0.24, \gamma_2 = 2.84$$


2. Grouped data

j	1	2	3	4	5
$y_j$	3	2	4	6	1
$f_j$	4	5	3	2	1

$$\bar{x} = 3.13, m_2 = 1.98, m_3 = 2.14, m_4 = 11.05$$

$$\gamma_1 = 0.77, \gamma_2 = 2.81$$

LINE	INSTRUCTIONS	DATA	KEYS	DISPLAY
1	Enter program on card 1			
2	Initialize		f REG	
3	For grouped data, go to 12			
4	Perform 4 for $i=1,2,\dots,n$	$x_i$	A	i
	(Correct erroneous data $x_k$ )	$x_k$	B	
5	Enter program on card 2			
6			A	$\bar{x}$
7			B	$m_2$
8			C	$m_3$
9			D	$m_4$
10			E	$\gamma_1$
11			R/S	$\gamma_2$
	(For a new case, go to 1)			
12	Perform 12–13 for $j=1,2,\dots,m$	$y_j$	↑	
13		$f_j$	C	
	(Correct erroneous data $y_h, f_h$ )	$y_h$	↑	
		$f_h$	D	
14	Go to 5			

**RANDOM NUMBER GENERATOR**

This program calculates:

- (1) Uniformly distributed random numbers  $u_i$  in the range

$$0 \leq u_i \leq 1$$

using the following formula:

$$u_i = \text{Fractional part of } [(\pi + u_{i-1})^8]$$

Initial value  $u_0 = 0$  is used.

- (2) Normally distributed random numbers  $n_i$  with mean  $m$  and standard deviation  $\sigma$ . The technique involves transforming uniform random variables to normal variables by the formulas:

$$N_i = (-2 \ln u_i)^{1/2} \cos(2\pi u_{i+1})$$

$$N_{i+1} = (-2 \ln u_i)^{1/2} \sin(2\pi u_{i+1})$$

where  $u_i, u_{i+1}$  are independent uniform random variables,

$$0 < u_i < 1.$$

The  $N_i$  thus generated are normally distributed with mean zero and unity variance.

Numbers  $N_i$  are used to generate a more general set of normally distributed numbers with mean  $m$  and standard deviation  $\sigma$  by

$$n_i = \sigma N_i + m$$

$$n_{i+1} = \sigma N_{i+1} + m$$

**Note:** Two initializing uniform random numbers  $u_a, u_b$  must be specified by the user, such that

$$u_a \neq u_b$$

$$0 < u_a < 1$$

$$0 < u_b < 1$$

**Reference:** Handbook of Mathematical Functions, U.S. Dept. of Commerce, Applied Mathematics Series, 1964

**Examples:**


- The following uniformly distributed pseudo random numbers are generated:  
0.53, 0.52, 0.39, 0.49, 0.97, 0.29, 0.65, 0.30, 0.40, 0.06, 0.14, 0.16, 0.68, 0.22, ...
- If  $m = 2$ ,  $\sigma = 1$ ,  $u_a = 0.23$ ,  $u_b = 0.82$  then the following pseudo normal numbers are obtained:  
2.73, 0.45, 2.52, 1.19, 3.51, 2.75, 1.79, 1.83, 1.00, 0.87, 1.90, 1.62, 1.74, 1.92, 1.24, 2.68, ...

LINE	INSTRUCTIONS	DATA	KEYS	DISPLAY
1	Enter program			
2	For normal numbers, go to 5			
3	Initialize		A	
4	Perform 4 for $i=1,2,3,\dots$		B	$u_i^*$
5	Store $m, \sigma$	$m$	$\uparrow$	
6		$\sigma$	C	
7	Store $u_a, u_b$	$u_a$	$\uparrow$	
8		$u_b$	D	
9	Perform 9 for $i=1,2,3,\dots$		E	$n_i$
	(Machine is set to RAD mode)			
	in subroutine E)			

\*If a different sequence of numbers is desired, choose a starting value  $u_0$  such that  $0 \leq u_0 \leq 1$  and do:

- $u_0$  **STO** **1**
- Skip step 3 and perform step 4.

## ANALYSIS OF VARIANCE (ONE WAY)



The one-way analysis of variance tests the differences between the population means of  $k$  treatment groups. Group  $i$  ( $i = 1, 2, \dots, k$ ) has  $n_i$  observations (treatment group may have equal or unequal number of observations).

$\text{Sum}_i$  = sum of observations in treatment group  $i$

$$= \sum_{j=1}^{n_i} x_{ij}$$

$$\text{Total SS} = \sum_{i=1}^k \sum_{j=1}^{n_i} x_{ij}^2 - \frac{\left( \sum_{i=1}^k \sum_{j=1}^{n_i} x_{ij} \right)^2}{\sum_{i=1}^k n_i}$$

$$\text{Treat SS} = \sum_{i=1}^k \frac{\left( \sum_{j=1}^{n_i} x_{ij} \right)^2}{n_i} - \frac{\left( \sum_{i=1}^k \sum_{j=1}^{n_i} x_{ij} \right)^2}{\sum_{i=1}^k n_i}$$

$$\text{Error SS} = \text{Total SS} - \text{Treat SS}$$

$$\text{df}_1 = \text{Treat df} = k - 1$$

$$\text{df}_2 = \text{Error df} = \sum_{i=1}^k n_i - k$$

$$\text{Treat MS} = \frac{\text{Treat SS}}{\text{Treat df}}$$

$$\text{Error MS} = \frac{\text{Error SS}}{\text{Error df}}$$

$$F = \frac{\text{Treat MS}}{\text{Error MS}} \quad (\text{with } k - 1 \text{ and } \sum_{i=1}^k n_i - k \text{ degrees of freedom})$$

Total SS, Treat SS, Error SS are in registers  $R_1, R_2, R_3$ .

**Note:** Erroneous data of the current treatment group can be corrected by entering the value then pressing **D** key.

**Reference:** Mathematical Statistics, J.E. Freund, Prentice Hall, 1962

**Example:**

		j	1	2	3	4	5	6
		i	10	8	5	12	14	11
		Treatment	6	9	8	13		
			14	13	10	17	16	

$$\text{Sum}_1 = 60.00$$

$$\text{Sum}_2 = 36.00$$

$$\text{Sum}_3 = 70.00$$

$$F = 3.79$$

$$df_1 = 2.00$$

$$df_2 = 12.00$$

LINE	INSTRUCTIONS	DATA	KEYS	DISPLAY
1	Enter program			
2	Initialize		f REG	
3	Perform 3–5 for $i=1,2,\dots,k$			
4	Perform 4 for $j=1,2,\dots,n_i$ (Correct erroneous data $x_{im}$ )	$x_{ij}$ $x_{im}$	A D j R/S	
5			B	$\text{Sum}_i$
6			C	F
7			R/S	$df_1$
8			R/S	$df_2$
	(For a new case, go to 2)			

**NORMAL DISTRIBUTION**

NORMAL DISTRIBUTION

**STAT 1-10A1**


NORMAL DISTRIBUTION

**f(x) Q(x)****STAT 1-10A2**

For a standard normal distribution

$$f(x) = \frac{1}{\sqrt{2\pi}} e^{-\frac{x^2}{2}}$$

$$Q(x) = \frac{1}{\sqrt{2\pi}} \int_x^{\infty} e^{-\frac{t^2}{2}} dt$$

For  $x \geq 0$ , polynomial approximation is used to compute  $Q(x)$ :

$$Q(x) = f(x) (b_1 t + b_2 t^2 + b_3 t^3 + b_4 t^4 + b_5 t^5) + \epsilon(x)$$

where  $|\epsilon(x)| < 7.5 \times 10^{-8}$ 

$$t = \frac{1}{1 + rx}, \quad r = 0.2316419$$

$$b_1 = .31938153, \quad b_2 = -.356563782$$

$$b_3 = 1.781477937, \quad b_4 = -1.821255978$$

$$b_5 = 1.330274429$$

**Note:**  $f(-x) = f(x)$ ,  $Q(-x) = 1 - Q(x)$

**Reference:** Handbook of Mathematical Functions, Abramowitz and Stegun, National Bureau of Standards, 1968

**Examples:**

1.  $f(1.18) = 0.20$   
 $Q(1.18) = 0.12$
2.  $f(2.28) = 0.03$   
 $Q(2.28) = 0.01$

LINE	INSTRUCTIONS	DATA	KEYS	DISPLAY
1	Enter program on card 1			
2			A	
3	Enter program on card 2			
4		x	A	f(x)
5			B	Q(x)
	(For a new x, go to 4)			

**INVERSE NORMAL INTEGRAL**


INIT	$x$	STAT 111A
------	-----	-----------



This program determines the value of  $x$  such that

$$Q = \int_x^{\infty} \frac{e^{-\frac{t^2}{2}}}{\sqrt{2\pi}} dt$$

where  $Q$  is given and  $0 < Q \leq 0.5$ .



The following rational approximation is used:

$$x = t - \frac{c_0 + c_1 t + c_2 t^2}{1 + d_1 t + d_2 t^2 + d_3 t^3} + \epsilon(Q)$$

where  $|\epsilon(Q)| < 4.5 \times 10^{-4}$

$$t = \sqrt{\ln \frac{1}{Q^2}}$$

$$c_0 = 2.515517 \quad d_1 = 1.432788$$

$$c_1 = 0.802853 \quad d_2 = 0.189269$$

$$c_2 = 0.010328 \quad d_3 = 0.001308$$

**Note:** If  $Q > 0.5$ , or  $Q \leq 0$ , flashing zeros will indicate the error.

**Reference:** Handbook of Mathematical Functions, Abramowitz and Stegun, National Bureau of Standards, 1968

**Examples:**

1.  $Q = 0.12$   
 $x = 1.18$

2.  $Q = 0.05$   
 $x = 1.65$

LINE	INSTRUCTIONS	DATA	KEYS	DISPLAY
1	Enter program			
2			A	
3		Q	B	x
	(For a new Q, go to 3)			

**CHI-SQUARE DISTRIBUTION**

CHI-SQUARE DISTRIBUTION


 $\Gamma(\nu/2)$  $f(x)$  $P(x)$ **STAT 1-12A**

This program evaluates the chi-square density

$$f(x) = \frac{1}{2^{\frac{\nu}{2}} \Gamma\left(\frac{\nu}{2}\right)} x^{\frac{\nu}{2}-1} e^{-\frac{x}{2}}$$

where  $x \geq 0$

$\nu$  is the degrees of freedom.



Series approximation is used to evaluate the cumulative distribution

$$P(x) = \int_0^x f(t) dt$$

$$= \left(\frac{x}{2}\right)^{\frac{\nu}{2}} \frac{e^{-\frac{x}{2}}}{\Gamma\left(\frac{\nu+2}{2}\right)} \left[ 1 + \sum_{k=1}^{\infty} \frac{x^k}{(\nu+2)(\nu+4)\dots(\nu+2k)} \right]$$

The program computes successive partial sums of the above series. When two consecutive partial sums are equal, the value is used as the sum of the series.

- Notes:**
1. Program requires  $\nu \leq 141$ . If  $\nu > 141$  and  $\nu$  is even, then display shows all 9's for  $\Gamma(\nu/2)$ ; if  $\nu > 141$  and  $\nu$  is odd, no warnings will be given, but answers are incorrect.
  2. If both  $x$  and  $\nu$  are large,  $f(x)$  may overflow the machine.
  3. If  $\nu$  is even,

$$\Gamma\left(\frac{\nu}{2}\right) = \left(\frac{\nu}{2} - 1\right)!$$

If  $\nu$  is odd,

$$\Gamma\left(\frac{\nu}{2}\right) = \left(\frac{\nu}{2} - 1\right)\left(\frac{\nu}{2} - 2\right) \cdots \left(\frac{1}{2}\right)\Gamma\left(\frac{1}{2}\right)$$


$$4. \quad \Gamma\left(\frac{1}{2}\right) = \sqrt{\pi}$$

**Reference:** Handbook of Mathematical Functions, Abramowitz and Stegun, National Bureau of Standards, 1968

### Examples:

1.  $\nu = 20, \quad \Gamma\left(\frac{\nu}{2}\right) = 362880.00$   
 $f(9.591) = 0.02, \quad P(9.591) = 0.03$   
 $f(15) = 0.06, \quad P(15) = 0.22$
2.  $\nu = 3, \quad \Gamma\left(\frac{\nu}{2}\right) = 0.89$   
 $f(7.82) = 0.02, \quad P(7.82) = 0.95$

LINE	INSTRUCTIONS	DATA	KEYS	DISPLAY
1	Enter program			
2		$\nu$	A	$\Gamma(\nu/2)$
3	Compute $f(x)$ and $P(x)$	x	B	$f(x)$
4			C	$P(x)$
	(For a different $x$ , go to 3.)			
	For a new case, go to 2)			


**t DISTRIBUTION**

This program evaluates the integral for t distribution

$$I(x, \nu) = \int_{-x}^x \frac{\Gamma\left(\frac{\nu+1}{2}\right)\left(1 + \frac{y^2}{\nu}\right)^{-\frac{\nu+1}{2}}}{\sqrt{\pi\nu} \Gamma\left(\frac{\nu}{2}\right)} dy$$

where  $x > 0$ ,

$\nu$  is the degrees of freedom.



Formulas used are:

(1)  $\nu$  even

$$I(x, \nu) = \sin \theta \left\{ 1 + \frac{1}{2} \cos^2 \theta + \frac{1 \cdot 3}{2 \cdot 4} \cos^4 \theta + \dots + \frac{1 \cdot 3 \cdot 5 \dots (\nu-3)}{2 \cdot 4 \cdot 6 \dots (\nu-2)} \cos^{\nu-2} \theta \right\}$$

(2)  $\nu$  odd

$$I(x, \nu) = \begin{cases} \frac{2\theta}{\pi} & \text{if } \nu = 1 \\ \frac{2\theta}{\pi} + \frac{2}{\pi} \cos \theta & \left\{ \sin \theta \left[ 1 + \frac{2}{3} \cos^2 \theta + \dots \right. \right. \\ & \left. \left. + \frac{2 \cdot 4 \dots (\nu - 3)}{1 \cdot 3 \dots (\nu - 2)} \cos^{\nu-3} \theta \right] \right\} & \text{if } \nu > 1 \end{cases}$$

where  $\theta = \tan^{-1}\left(\frac{x}{\sqrt{\nu}}\right)$

**Reference:** Handbook of Mathematical Functions, Abramowitz and Stegun, National Bureau of Standards, 1968

**Example:**

$$I(2.201, 11) = 0.95$$

$$I(2.75, 30) = 0.99$$

LINE	INSTRUCTIONS	DATA	KEYS	DISPLAY
1	Enter program			
2		x	↑	
3		$\nu$	A	$I(x, \nu)$
	(Machine now is in RAD mode)			


**F DISTRIBUTION**

F DISTRIBUTION	STAT 114A			
$v_1$	$v_2$	$x$	$v_1$ even	$v_2$ even

This program evaluates the integral of the F distribution

$$Q(x) = \int_x^{\infty} \frac{\Gamma\left(\frac{v_1 + v_2}{2}\right) y^{\frac{v_1}{2}-1} \left(\frac{v_1}{v_2}\right)^{\frac{v_1}{2}}}{\Gamma\left(\frac{v_1}{2}\right) \Gamma\left(\frac{v_2}{2}\right) \left(1 + \frac{v_1}{v_2} y\right)^{\frac{v_1+v_2}{2}}} dy$$

for given values of  $x$  ( $>0$ ), degrees of freedom  $v_1, v_2$ , provided either  $v_1$  or  $v_2$  is even.



The integral is evaluated by means of the following series:

(1)  $v_1$  even

$$Q(x) = t^{\frac{v_2}{2}} \left[ 1 + \frac{v_2}{2} (1-t) + \dots + \frac{v_2(v_2+2)\dots(v_2+v_1-4)}{2 \cdot 4 \dots (v_1-2)} (1-t)^{\frac{v_1-2}{2}} \right]$$

(2)  $\nu_2$  even

$$Q(x) = 1 - (1-t)^{\frac{\nu_1}{2}} \left[ 1 + \frac{\nu_1}{2} t + \dots + \frac{\nu_1(\nu_1+2)\dots(\nu_2+\nu_1-4)}{2\cdot4\dots(\nu_2-2)} t^{\frac{\nu_2-2}{2}} \right]$$

$$\text{where } t = \frac{\nu_2}{\nu_2 + \nu_1} x$$

**Note:** If both  $\nu_1$ ,  $\nu_2$  are even, the two formulas would generate identical answers. Using the smaller of  $\nu_1$ ,  $\nu_2$  could save computation time. For example, if  $\nu_1 = 10$ ,  $\nu_2 = 20$ , then classify the problem as  $\nu_1$  is even and use the **D** key to obtain the answer.

**Examples:**

1.  $\nu_1 = 7$ ,  $\nu_2 = 6$

$$Q(4.21) = 0.05$$

2.  $\nu_1 = 4$ ,  $\nu_2 = 20$

$$Q(2.25) = 0.10$$

LINE	INSTRUCTIONS	DATA	KEYS	DISPLAY
1	Enter program			
2		$\nu_1$	A	
3		$\nu_2$	B	
4		x	C	
5	If $\nu_1$ is even		D	
6	If $\nu_2$ is even		E	
	(For a new case, go to 2)			

**BIVARIATE NORMAL DISTRIBUTION**

BIVARIATE NORMAL DISTRIBUTION

**STAT 1-15A** $\mu_1, \sigma_1$  $\mu_2, \sigma_2$  $\rho$  $f(x,y)$ 

$$f(x,y) = \frac{1}{2\pi \sigma_1 \sigma_2 \sqrt{1 - \rho^2}} e^{-P(x,y)}$$

where

$$P(x,y) = \frac{1}{2(1 - \rho^2)} \left[ \frac{(x - \mu_1)^2}{\sigma_1^2} - 2\rho \frac{(x - \mu_1)(y - \mu_2)}{\sigma_1 \sigma_2} + \frac{(y - \mu_2)^2}{\sigma_2^2} \right]$$

- Notes:**
1.  $\sigma_1 \neq 0, \sigma_2 \neq 0$
  2. Program requires  $\rho^2 < 1$ .

**Reference:** Mathematical Statistics, J.E. Freund, Prentice Hall, 1962

**Example:**

$$\mu_1 = -1, \sigma_1 = 1.5$$

$$\mu_2 = 1, \sigma_2 = 0.5$$

$$\rho = 0.7$$

$$f(1, 2) = 0.04$$

$$f(-1, 1) = 0.30$$

LINE	INSTRUCTIONS	DATA	KEYS	DISPLAY
1	Enter program			
2		$\mu_1$	$\uparrow$	
3		$\sigma_1$	A	
4		$\mu_2$	$\uparrow$	
5		$\sigma_2$	B	
6		$\rho$	C	
7		x	$\uparrow$	
8		y	D	$f(x, y)$
	(For new values of x, y go to 7)			

**LOGARITHMIC NORMAL DISTRIBUTION**

LOGARITHMIC NORMAL DISTRIBUTION				STAT 1-16A
median	mode	mean	var	f(x)

If  $X$  is a random variable whose logarithm is normally distributed with mean  $m$  and variance  $\sigma^2$ , then  $X$  has a logarithmic normal distribution with density function

$$f(x) = \frac{1}{x \sqrt{2\pi\sigma^2}} e^{-\frac{1}{2\sigma^2}(\ln x - m)^2}$$

where  $x > 0$

This program computes  $f(x)$  and the following statistics for given  $m$ ,  $\sigma^2$ :

$$\text{median} = e^m$$

$$\text{mode} = e^{m - \sigma^2}$$

$$\text{mean} = e^m + \frac{\sigma^2}{2}$$

$$\text{variance} = e^{\sigma^2 + 2m} (e^{\sigma^2} - 1)$$

**Note:** Program requires  $\sigma^2 \neq 0$ .

**Reference:** Statistical Theory and Methodology in Science and Engineering, K.A. Brownlee, John Wiley & Sons, 1965

**Example:**

$$m = 1, \sigma^2 = 1$$

$$\text{median} = 2.72$$

$$\text{mode} = 1.00$$

$$\text{mean} = 4.48$$


$$\text{variance} = 34.51$$

$$f(.1) = 0.02$$

$$f(.6) = 0.21$$

$$f(1) = 0.24$$

LINE	INSTRUCTIONS	DATA	KEYS	DISPLAY
1	Enter program			
2		m	↑	
3		$\sigma^2$	A	median
4			B	mode
5			C	mean
6			D	variance
7	Compute $f(x)$	x	E	$f(x)$
	(For a different x, go to 7)			

**WEIBULL DISTRIBUTION**

This program can be used to find:

$$(1) \quad f(x) = ab x^{b-1} \exp(-ax^b)$$

where  $a > 0$ ,  $b > 0$ ,  $x > 0$

$$(2) \quad Q(x) = \int_x^{\infty} ab t^{b-1} \exp(-at^b) dt \\ = \exp(-ax^b)$$

$$(3) \quad x \text{ (for a given } Q, 0 < Q < 1\text{), such that}$$

$$Q = \int_x^{\infty} f(t) dt$$

The following formula is used:

$$x = \left( \frac{\ln Q}{-a} \right)^{\frac{1}{b}}$$


**Reference:** Statistics in Research, Bernard Ostle,  
Iowa State University Press, 1963

**Example:**

$$a = 0.1, b = 0.8$$

1.  $f(3.2) = 0.05$
2.  $Q(3.2) = 0.78$
3. If  $Q = 0.5$ , then  $x = 11.25$

LINE	INSTRUCTIONS	DATA	KEYS	DISPLAY
1	Enter program			
2		a	↑	
3		b	A	
4	Compute $f(x)$ , $Q(x)$	x	B	$f(x)$
5			C	$Q(x)$
6	Find $x$ for a given $Q$	Q	D	x

**BINOMIAL DISTRIBUTION**

This program evaluates the binomial density function for given p and n:

$$f(x) = \binom{n}{x} p^x (1-p)^{n-x}$$

where      n is a positive integer

$0 < p < 1$  and

$x = 0, 1, 2, \dots, n$

The recursive relation

$$f(x+1) = \frac{p(n-x)}{(x+1)(1-p)} f(x)$$

$$(x = 0, 1, 2, \dots, n-1)$$

is used to find the cumulative distribution

$$P(x) = \sum_{k=0}^{x} f(k)$$

The mean m and the variance  $\sigma^2$  are given by

$$m = np$$

$$\sigma^2 = np(1-p)$$

**Reference:** Modern Probability Theory and its Applications,  
E. Parzen, John Wiley & Sons, 1960

**Example:**

$$p = 0.49, \quad n = 6$$

$$m = 2.94, \quad \sigma^2 = 1.50$$

$$f(4) = 0.22$$

$$P(4) = 0.90$$

LINE	INSTRUCTIONS	DATA	KEYS	DISPLAY
1	Enter program			
2		p	↑	
3		n	A	m
4			R/S	$\sigma^2$
5	Compute $f(x)$ and $P(x)$	x	B	$f(x)$
6			C	$P(x)$
	(For a new value of x, go to 5)			

**NEGATIVE BINOMIAL DISTRIBUTION**

NEGATIVE BINOMIAL DISTRIBUTION				STAT 1-19A	
p, r	m	$\sigma^2$	f(x)	P(x)	

This program evaluates the negative binomial density function for given p and r:

$$f(x) = \binom{x + r - 1}{r - 1} p^r (1 - p)^x$$

where      r is a positive integer

0 < p < 1    and

x = 0, 1, 2, ...

The recursive relation

$$f(x + 1) = \frac{(1 - p)(x + r)}{x + 1} f(x)$$

is used to find the cumulative distribution

$$P(x) = \sum_{k=0}^x f(k)$$

The mean m and the variance  $\sigma^2$  are given by

$$m = \frac{r(1 - p)}{p}$$

$$\sigma^2 = \frac{r(1 - p)}{p^2}$$

**Note:** If we interpret p as the probability of success of a given event, then f(x) is the probability that exactly x + r trials will be required to get r successes.

**Reference:** Modern Probability Theory and its Applications, E. Parzen, John Wiley & Sons, 1960

**Example:**

$$p = 0.9, \quad r = 4$$

$$m = 0.44$$

$$\sigma^2 = 0.49$$

$$f(1) = 0.26$$

$$P(1) = 0.92$$

$$f(2) = 0.07$$

$$P(2) = 0.98$$

LINE	INSTRUCTIONS	DATA	KEYS	DISPLAY
1	Enter program			
2	Enter p, r	p	↑	
3		r	A	
4	Compute mean m		B	m
5	Compute variance $\sigma^2$		C	$\sigma^2$
6	Compute $f(x)$ , $P(x)$	x	D	$f(x)$
7			E	$P(x)$
	(For a different x, go to 6)			

**HYPERGEOMETRIC DISTRIBUTION**

HYPERGEOMETRIC DISTRIBUTION

**STAT 1-20A**

a,b

n

f(x)

P(x)

m,σ<sup>2</sup>

This program evaluates the hypergeometric density function for given a, b and n:

$$f(x) = \frac{\binom{a}{x} \binom{b}{n-x}}{\binom{a+b}{n}}$$

where a, b, n are positive integers

$x \leq a, n - x \leq b$  and

$x = 0, 1, 2, \dots, n$

The recursive relation

$$f(x+1) = \frac{(x-a)(x-n)}{(x+1)(b-n+x+1)} f(x)$$

$(x = 0, 1, 2, \dots, n-1)$

is used to find the cumulative distribution

$$P(x) = \sum_{k=0}^x f(k)$$

The mean m and the variance  $\sigma^2$  are given by

$$m = \frac{an}{a+b}$$

$$\sigma^2 = \frac{abn(a+b-n)}{(a+b)^2(a+b-1)}$$

**Notes:** 1.  $f(0) = P(0)$

2. When x is large, due to round-off error, the computed value for P(x) might be slightly greater than one. In that case, let  $P(x) = 1$ .

3. This program requires  $a + b \leq 69$ .

**Reference:** Mathematical Statistics, J.E. Freund, Prentice Hall, 1962

**Example:**

Given  $a = 8$ ,  $b = 12$ ,  $n = 6$ , then

$$f(0) = P(0) = 0.02$$


$$f(3) = 0.32, \quad P(3) = 0.86$$

$$f(5) = 0.02, \quad P(5) = 1.00$$

$$m = 2.40$$

$$\sigma^2 = 1.06$$

LINE	INSTRUCTIONS	DATA	KEYS	DISPLAY
1	Enter program			
2		a	↑	
3		b	A	
4		n	B	f(0)
5	For $x \geq 1$	x	C	f(x)
6			D	P(x)
	(For a new value of x, go to 5.			
	For a new n, go to 4.			
	For different a, b, go to 2)			
7	Compute $m$ , $\sigma^2$		E	m
8			R/S	$\sigma^2$

**POISSON DISTRIBUTION**

Density function

$$f(x) = \frac{\lambda^x e^{-\lambda}}{x!}$$

where  $\lambda > 0$

$x = 0, 1, 2, \dots$

Cumulative distribution

$$P(x) = \sum_{k=0}^x f(k)$$

This program evaluates  $f(x)$  and  $P(x)$  for a given  $\lambda$  using the recursive relation

$$f(x+1) = \frac{\lambda}{x+1} f(x)$$

Note: Mean = variance =  $\lambda$

**Example:**


$$\lambda = 3.2$$

$$f(0) = 0.04$$

$$f(7) = 0.03$$

$$P(7) = 0.98$$

LINE	INSTRUCTIONS	DATA	KEYS	DISPLAY
1	Enter program			
2		$\lambda$	A	$f(0)$
3	Compute $f(x)$ and $P(x)$	x	B	$f(x)$
4			C	$P(x)$
	(For new value of x, go to 3.)			
	For new value of $\lambda$ , go to 2)			

**LINEAR REGRESSION**

This program must be used in conjunction with *Stat 1-05A, Sums for Two Variables*, to fit a straight line

$$y = a_0 + a_1 x$$

to a set of data points  $\{(x_i, y_i), i = 1, 2, \dots, n\}$  by the least squares method.

The program computes:

- regression coefficients  $a_0, a_1$

$$a_1 = \frac{\sum x_i y_i - \frac{\sum x_i \sum y_i}{n}}{\sum x_i^2 - \frac{(\sum x_i)^2}{n}}$$

$$a_0 = \bar{y} - a_1 \bar{x}$$

where

$$\bar{x} = \frac{\sum x_i}{n}$$

$$\bar{y} = \frac{\sum y_i}{n}$$

- coefficient of determination

$$r^2 = \frac{\left[ \sum x_i y_i - \frac{\sum x_i \sum y_i}{n} \right]^2}{\left[ \sum x_i^2 - \frac{(\sum x_i)^2}{n} \right] \left[ \sum y_i^2 - \frac{(\sum y_i)^2}{n} \right]}$$

$r^2$  can be interpreted as the proportion of total variation about the mean  $\bar{y}$  explained by the regression. In other words,  $r^2$  measures the “goodness of fit” of the regression line. Note that  $0 \leq r^2 \leq 1$ , and if  $r^2 = 1$ , we have a perfect fit.

3. estimated value  $\hat{y}$  on the regression line for any given  $x$

$$\hat{y} = a_0 + a_1 x$$

4. standard error of estimate of  $y$  on  $x$

$$\begin{aligned}s_{y \cdot x} &= \sqrt{\frac{\sum(y_i - \hat{y}_i)^2}{n - 2}} \\&= \sqrt{\frac{\sum y_i^2 - a_0 \sum y_i - a_1 \sum x_i y_i}{n - 2}}\end{aligned}$$

5. standard error of the regression coefficient  $a_0$

$$s_0 = s_{y \cdot x} \sqrt{\frac{\sum x_i^2}{n \left[ \sum x_i^2 - \frac{(\sum x_i)^2}{n} \right]}}$$

6. standard error of the regression coefficient  $a_1$

$$s_1 = \frac{s_{y \cdot x}}{\sqrt{\sum x_i^2 - \frac{(\sum x_i)^2}{n}}}$$

**Note:**  $n$  is a positive integer and  $n \neq 1$  or  $2$ .

### References:

Applied Regression Analysis, Draper and Smith, John Wiley & Sons, 1966

Statistics in Research, B. Ostle, Iowa State University Press, 1963

**Example:**

$x_i$	26	30	44	50	62	68	74
$y_i$	92	85	78	81	54	51	40

1.  $a_0 = 121.04$

$a_1 = -1.03$

Regression line is  $y = 121.04 - 1.03x$ 

2.  $r^2 = 0.92$


3. For  $x = 80$ ,  $\hat{y} = 38.27$

4.  $s_{y \cdot x} = 6.34$

5.  $s_0 = 7.47$

$s_1 = 0.14$

LINE	INSTRUCTIONS	DATA	KEYS	DISPLAY
1	Enter program Stat 1–05A			
2	Initialize		f REG	
3	Perform 3–4 for $i = 1, 2, \dots, n$	$x_i$	$\uparrow$	
4		$y_i$	A	i
	(Correct erroneous data $x_k, y_k$ )	$x_k$	$\uparrow$	
		$y_k$	B	
5	Enter program Stat 1–22A			
6			A	$a_0$
7			R/S	$a_1$
8			B	$r^2$
9		x	C	$\hat{y}$
	(For a new x, go to 9)			
10			D	$s_{y \cdot x}$
11			E	$s_0$
12			R/S	$s_1$

**EXPONENTIAL CURVE FIT**

This program computes the least squares fit of n pairs of data points  $\{(x_i, y_i), i = 1, 2, \dots, n\}$ , where  $y_i > 0$ , for an exponential function of the form

$$y = a e^{bx} \quad (a > 0)$$

The equation is linearized into

$$\ln y = \ln a + bx$$

The following statistics are computed:

1. Coefficients a, b

$$b = \frac{\sum x_i \ln y_i - \frac{1}{n} (\sum x_i) (\sum \ln y_i)}{\sum x_i^2 - \frac{1}{n} (\sum x_i)^2}$$

$$a = \exp \left[ \frac{\sum \ln y_i}{n} - b \frac{\sum x_i}{n} \right]$$

2. Coefficient of determination

$$r^2 = \frac{\left[ \sum x_i \ln y_i - \frac{1}{n} \sum x_i \sum \ln y_i \right]^2}{\left[ \sum x_i^2 - \frac{(\sum x_i)^2}{n} \right] \left[ \sum (\ln y_i)^2 - \frac{(\sum \ln y_i)^2}{n} \right]}$$

3. Estimated value  $\hat{y}$  for a given x

$$\hat{y} = a e^{bx}$$

**Note:** n is a positive integer and n ≠ 1.

**Reference:** Statistical Theory and Methodology in Science and Engineering, K.A. Brownlee, John Wiley & Sons, 1965

**Example:**

$x_i$	.72	1.31	1.95	2.58	3.14
$y_i$	2.16	1.61	1.16	.85	0.5

1.  $a = 3.45, b = -0.58$

$y = 3.45 e^{-0.58x}$

2.  $r^2 = 0.98$

3. For  $x = 1.5, \hat{y} = 1.44$

LINE	INSTRUCTIONS	DATA	KEYS	DISPLAY
1	Enter program			
2	Initialize		f REG	
3	Perform 3–4 for $i=1, 2, \dots, n$	$x_i$	↑	
4		$y_i$	A	i
5			B	a
6			R/S	b
7			C	$r^2$
8	Compute estimated value $\hat{y}$	x	D	$\hat{y}$
	(For a new x, go to 8)			

**POWER CURVE FIT**

Σ+	POWER CURVE FIT	STAT 1-24A
a,b	$r^2$	$\hat{y}$

This program fits a power curve

$$y = ax^b \quad (a > 0)$$

to a set of data points

$$\{(x_i, y_i), i = 1, 2, \dots, n\}$$

where  $x_i > 0, y_i > 0$ .

By writing this equation as

$$\ln y = b \ln x + \ln a$$

the problem can be solved as a linear regression problem.

Output statistics are:

1. Regression coefficients

$$b = \frac{\sum (\ln x_i)(\ln y_i) - \frac{(\sum \ln x_i)(\sum \ln y_i)}{n}}{\sum (\ln x_i)^2 - \frac{(\sum \ln x_i)^2}{n}}$$

$$a = \exp \left[ \frac{\sum \ln y_i}{n} - b \frac{\sum \ln x_i}{n} \right]$$

2. Coefficient of determination

$$r^2 = \frac{\left[ \sum (\ln x_i)(\ln y_i) - \frac{(\sum \ln x_i)(\sum \ln y_i)}{n} \right]^2}{\left[ \sum (\ln x_i)^2 - \frac{(\sum \ln x_i)^2}{n} \right] \left[ \sum (\ln y_i)^2 - \frac{(\sum \ln y_i)^2}{n} \right]}$$

3. Estimated value  $\hat{y}$  for given  $x$

$$\hat{y} = ax^b$$

**Note:**  $n$  is a positive integer and  $n \neq 1$ .

**Reference:** Statistical Theory and Methodology in Science and Engineering, K.A. Brownlee, John Wiley & Sons, 1965

**Example:**

$x_i$	10	12	15	17	20	22	25	27	30	32	35
$y_i$	.95	1.05	1.25	1.41	1.73	2.00	2.53	2.98	3.85	4.59	6.02

1.  $a = 0.03, b = 1.46$


$$y = 0.03x^{1.46}$$

2.  $r^2 = 0.94$

3. For  $x = 18, \hat{y} = 1.76$

$$x = 23, \hat{y} = 2.52$$

LINE	INSTRUCTIONS	DATA	KEYS	DISPLAY
1	Enter program			
2	Initialize		f REG	
3	Perform 3-4 for $i=1, 2, \dots, n$	$x_i$	$\uparrow$	
4		$y_i$	A	i
5			B	a
6			R/S	b
7			C	$r^2$
8	Compute estimated value $\hat{y}$	x	D	$\hat{y}$
	(For a new x, go to 8)			

**LOGARITHMIC CURVE FIT**

This program fits a logarithmic curve

$$y = a + b \ln x$$

to a set of data points

$$\{(x_i, y_i), i = 1, 2, \dots, n\}$$

where

$$x_i > 0.$$

Program computes:

1. Regression coefficients

$$b = \frac{\sum y_i \ln x_i - \frac{1}{n} \sum \ln x_i \sum y_i}{\sum (\ln x_i)^2 - \frac{1}{n} (\sum \ln x_i)^2}$$

$$a = \frac{1}{n} (\sum y_i - b \sum \ln x_i)$$

2. Coefficient of determination

$$r^2 = \frac{\left[ \sum y_i \ln x_i - \frac{1}{n} \sum \ln x_i \sum y_i \right]^2}{\left[ \sum (\ln x_i)^2 - \frac{1}{n} (\sum \ln x_i)^2 \right] \left[ \sum y_i^2 - \frac{1}{n} (\sum y_i)^2 \right]}$$

3. Estimated value  $\hat{y}$  for given  $x$

$$\hat{y} = a + b \ln x$$


**Note:**  $n$  is a positive integer and  $n \neq 1$ .

**Example:**

$x_i$	3	4	6	10	12
$y_i$	1.5	9.3	23.4	45.8	60.1

1.  $a = -47.02, b = 41.39$   
 $y = -47.02 + 41.39 \ln x$
2.  $r^2 = 0.98$
3. For  $x = 8, \hat{y} = 39.06$   
For  $x = 14.5, \hat{y} = 63.67$

LINE	INSTRUCTIONS	DATA	KEYS	DISPLAY
1	Enter program			
2	Initialize		f REG	
3	Perform 3–4 for $i=1, 2, \dots, n$	$x_i$	↑	
4		$y_i$	A	i
5			B	a
6			R/S	b
7			C	$r^2$
8	Compute estimated value $\hat{y}$	x	D	$\hat{y}$
	(For a new x, go to 8)			

**LEAST SQUARES REGRESSION OF  $y = cx^a + dx^b$** 

This program determines the coefficients c, d of the equation

$$y = cx^a + dx^b$$

for a set of data points

$$\{(x_i, y_i), i = 1, 2, \dots, n\}$$

where a, b are any given real numbers.

$$d = \frac{(\sum x_i^{2a})(\sum x_i^b y_i) - (\sum x_i^a y_i)(\sum x_i^{a+b})}{(\sum x_i^{2b})(\sum x_i^{2a}) - (\sum x_i^{a+b})^2}$$

$$c = \frac{\sum x_i^a y_i - d \sum x_i^{a+b}}{\sum x_i^{2a}}$$

where  $x_i > 0$  for  $i = 1, 2, \dots, n$ .

**Note:** n is a positive integer and  $n \neq 1$ .

**Example:**

$$a = 0.5, \quad b = 3$$

$x_i$	1	4	9	16
$y_i$	9	-44	-699	-4056

$$c = 10.00, \quad d = -1.00$$

$$\text{Regression line is } y = 10x^{1/2} - x^3$$

$$\text{For } x = 6, \quad \hat{y} = -191.51$$

LINE	INSTRUCTIONS	DATA	KEYS	DISPLAY
1	Enter program			
2	Initialize		f REG	
3		a	↑	
4		b	A	
5	Perform 5–6 for i=1, 2, ..., n	$x_i$	↑	
6		$y_i$	B	
7			C	c
8			R/S	d
9	Compute estimated value $\hat{y}$ on the line	x	D	$\hat{y}$
	(For a new x, go to 9)			

**MULTIPLE LINEAR REGRESSION**

MULTIPLE LINEAR REGRESSION

**STAT 1-27A1** $\Sigma+$        $\Sigma-$ 

MULTIPLE LINEAR REGRESSION

**STAT 1-27A2** $a_0$        $a_1$        $a_2$        $\hat{z}$ 

For a set of data points  $\{(x_i, y_i, z_i), i = 1, 2, \dots, n\}$  this program fits a linear equation of the form

$$z = a_0 + a_1 x + a_2 y$$

by the least squares method.

Regression coefficients  $a_0, a_1, a_2$  can be found by solving the normal equations:

$$\begin{cases} \sum z_i = a_0 n + a_1 \sum x_i + a_2 \sum y_i \\ \sum x_i z_i = a_0 \sum x_i + a_1 \sum x_i^2 + a_2 \sum x_i y_i \quad i = 1, 2, \dots, n \\ \sum y_i z_i = a_0 \sum y_i + a_1 \sum x_i y_i + a_2 \sum y_i^2 \end{cases}$$

$$a_2 = \frac{A - B}{[n \sum x_i^2 - (\sum x_i)^2] [n \sum y_i^2 - (\sum y_i)^2] - [n \sum x_i y_i - (\sum x_i)(\sum y_i)]^2}$$

$$\text{where } A = [n \sum x_i^2 - (\sum x_i)^2] [n \sum y_i z_i - (\sum y_i)(\sum z_i)]$$

$$B = [n \sum x_i y_i - (\sum x_i)(\sum y_i)] [n \sum x_i z_i - (\sum x_i)(\sum z_i)]$$

$$a_1 = \frac{[n \sum x_i z_i - (\sum x_i)(\sum z_i)] - a_2 [n \sum x_i y_i - (\sum x_i)(\sum y_i)]}{n \sum x_i^2 - (\sum x_i)^2}$$

$$a_0 = \frac{1}{n} (\sum z_i - a_2 \sum y_i - a_1 \sum x_i)$$

**Notes:** 1.  $\sum x_i y_i, \sum x_i z_i, \sum y_i z_i, \sum y_i^2, n, \sum x_i^2, \sum x_i, \sum y_i, \sum z_i$  are in storage registers  $R_1$  through  $R_9$  before program on card 2 is executed. Recall and record these sums if desired when instructions indicate to do so.

2. Erroneous data  $x_k, y_k, z_k$  can be removed by the following keystrokes:  $x_k \uparrow y_k \uparrow z_k \blacksquare B$

3.  $n$  is a positive integer and  $n \neq 1$ .

**Reference:** Introduction to the Theory of Statistics, Mood and Graybill, McGraw-Hill, 1963

**Example:**

i	1	2	3	4	$\Sigma x_i y_i = 17.57$	$\Sigma x_i = 6.55$
$x_i$	1.5	0.45	1.8	2.8	$\Sigma x_i z_i = 38.65$	$\Sigma y_i = 9.10$
$y_i$	0.7	2.3	1.6	4.5	$\Sigma y_i z_i = 59.53$	$\Sigma z_i = 19.60$
$z_i$	2.1	4.0	4.1	9.4	$\Sigma y_i^2 = 28.59$	$a_0 = -0.10$
					$n = 4.00$	$a_1 = 0.79$
					$\Sigma x_i^2 = 13.53$	$a_2 = 1.63$

Regression line is  

$$z = -0.10 + 0.79x + 1.63y$$

For  $x = 2$ ,  $y = 3$ ,  $\hat{z} = 6.37$

LINE	INSTRUCTIONS	DATA	KEYS	DISPLAY
1	Enter program on card 1			
2	Initialize		f REG	
3	Perform 3–5 for $i=1, 2, \dots, n$	$x_i$	$\uparrow$	
4		$y_i$	$\uparrow$	
5		$z_i$	A	i
	(Correct erroneous data)	$x_k$	$\uparrow$	
	$x_k, y_k, z_k$ )	$y_k$	$\uparrow$	
		$z_k$	B	
6	Recall and record sums		RCL 1	$\Sigma x_i y_i$
7			RCL 2	$\Sigma x_i z_i$
8			RCL 3	$\Sigma y_i z_i$
9			RCL 4	$\Sigma y_i^2$
10			RCL 5	n
11			RCL 6	$\Sigma x_i^2$
12			RCL 7	$\Sigma x_i$
13			RCL 8	$\Sigma y_i$
14			RCL 9	$\Sigma z_i$
15	Enter program on card 2			
16			A	$a_0$
17			B	$a_1$
18			C	$a_2$
19	Obtain estimated value $\hat{z}$ on the	x	$\uparrow$	
20	line (for new values, go to 19)	y	D	$\hat{z}$

**PARABOLIC CURVE FIT**

For a set of data points  $\{(x_i, y_i), i = 1, 2, \dots, n\}$  this program fits a parabola

$$y = a_0 + a_1 x + a_2 x^2$$

This program must be used in conjunction with *Stat 1-27A, Multiple Linear Regression*, to compute:

1. Regression coefficients

$$a_2 = \frac{A - B}{[n \sum x_i^2 - (\sum x_i)^2] [n \sum x_i^4 - (\sum x_i^2)^2] - [n \sum x_i^3 - (\sum x_i)(\sum x_i^2)]^2}$$

where

$$A = [n \sum x_i^2 - (\sum x_i)^2] [n \sum x_i^2 y_i - (\sum x_i^2)(\sum y_i)]$$

$$B = [n \sum x_i^3 - (\sum x_i)(\sum x_i^2)] [n \sum x_i y_i - (\sum x_i)(\sum y_i)]$$

$$a_1 = \frac{[n \sum x_i y_i - (\sum x_i)(\sum y_i)] - a_2 [n \sum x_i^3 - (\sum x_i)(\sum x_i^2)]}{n \sum x_i^2 - (\sum x_i)^2}$$

$$a_0 = \frac{1}{n} (\sum y_i - a_2 \sum x_i^2 - a_1 \sum x_i)$$

2. Estimated value  $\hat{y}$  for given  $x$

$$\hat{y} = a_0 + a_1 x + a_2 x^2$$

**Note:**  $n$  is a positive integer and  $n \neq 1$ .


**Reference:** Introduction to the Theory of Statistics, Mood and Graybill, McGraw Hill, 1963

**Example:**

$x_i$	0	1	1.5	3	5
$y_i$	2.1	2	-5	-24.5	-80

1.  $\Sigma x_i^3 = 156.38$ ,  $\Sigma x_i y_i = -479.00$ ,  $\Sigma x_i^2 y_i = -2229.75$   
 $\Sigma x_i^4 = 712.06$ ,  $n = 5.00$ ,  $\Sigma x_i^2 = 37.25$   
 $\Sigma x_i = 10.50$ ,  $\Sigma y_i = -105.40$
2.  $a_0 = 2.28$ ,  $a_1 = 1.85$ ,  $a_2 = -3.66$   
 $y = 2.28 + 1.85x - 3.66x^2$
3. For  $x = 4$ ,  $\hat{y} = -48.83$

LINE	INSTRUCTIONS	DATA	KEYS	DISPLAY
1	Enter program Stat 1–28A			
2	Initialize		f REG	
3	Perform 3–4 for $i=1, 2, \dots, n$	$x_i$	$\uparrow$	
4		$y_i$	A	i
	(Correct erroneous data $x_k, y_k$ )	$x_k$	$\uparrow$	
		$y_k$	B	
5	Recall and record sums		RCL 1	$\Sigma x_i^3$
6			RCL 2	$\Sigma x_i y_i$
7			RCL 3	$\Sigma x_i^2 y_i$
8			RCL 4	$\Sigma x_i^4$
9			RCL 5	n
10			RCL 6	$\Sigma x_i^2$
11			RCL 7	$\Sigma x_i$
12			RCL 9	$\Sigma y_i$
13	Enter program Stat 1–27A2			
14			A	$a_0$
15			B	$a_1$
16			C	$a_2$
17	Enter program Stat 1–28A			
18	Compute estimated value $\hat{y}$	x	C	$\hat{y}$
	(For a new x, go to 18)			

**PAIRED t STATISTIC**

Given a set of paired observations from two normal populations with means  $\mu_1, \mu_2$  (unknown)

$x_i$	$x_1$	$x_2$	...	$x_n$
$y_i$	$y_1$	$y_2$	...	$y_n$

let

$$D_i = x_i - y_i$$

$$\bar{D} = \frac{1}{n} \sum_{i=1}^n D_i$$

$$s_D = \sqrt{\frac{\sum D_i^2 - \frac{1}{n} (\sum D_i)^2}{n-1}}$$

$$s_{\bar{D}} = \frac{s_D}{\sqrt{n}}$$

The test statistic

$$t = \frac{\bar{D}}{s_{\bar{D}}}$$

which has  $n - 1$  degrees of freedom (df) can be used to test the null hypothesis

$$H_0: \mu_1 = \mu_2$$

**Reference:** Statistics in Research, B. Ostle, Iowa State University Press, 1963

**Example:**

$x_i$	14	17.5	17	17.5	15.4
$y_i$	17	20.7	21.6	20.9	17.2

$$\bar{D} = -3.20$$

$$s_D = 1.00$$

$$t = -7.16$$

$$df = 4.00$$

LINE	INSTRUCTIONS	DATA	KEYS	DISPLAY
1	Enter program			
2	Initialize		A	
3	Perform 3–4 for $i=1, 2, \dots, n$	$x_i$	$\uparrow$	
4		$y_i$	B	i
	(Correct erroneous data $x_k, y_k$ )	$x_k$	$\uparrow$	
		$y_k$	E	
5			C	$\bar{D}$
6			R/S	$s_D$
7			D	t
8			R/S	df

**t STATISTIC FOR TWO MEANS**

t STATISTIC FOR TWO MEANS			STAT 1-30A	
INIT	$\Sigma+$	D	t, df	$\Sigma-$

Suppose  $\{x_1, x_2, \dots, x_{n_1}\}$  and  $\{y_1, y_2, \dots, y_{n_2}\}$  are independent random samples from two normal populations having means  $\mu_1, \mu_2$  (unknown) and the same unknown variance  $\sigma^2$ .

We want to test the null hypothesis

$$H_0: \mu_1 - \mu_2 = D$$

Define

$$\bar{x} = \frac{1}{n_1} \sum_{i=1}^{n_1} x_i$$

$$\bar{y} = \frac{1}{n_2} \sum_{i=1}^{n_2} y_i$$

$$t = \frac{\bar{x} - \bar{y} - D}{\sqrt{\frac{1}{n_1} + \frac{1}{n_2}} \sqrt{\frac{\sum x_i^2 - n_1 \bar{x}^2 + \sum y_i^2 - n_2 \bar{y}^2}{n_1 + n_2 - 2}}}$$

We can use this t statistic which has the t distribution with  $n_1 + n_2 - 2$  degrees of freedom (df) to test the null hypothesis  $H_0$ .

**Note:**  $n_2, \Sigma y_i, \Sigma y_i^2, n_1, \Sigma x_i, \Sigma x_i^2$  are in registers R<sub>1</sub> through R<sub>6</sub>.

**Reference:** Statistical Theory and Methodology in Science and Engineering, K. A. Brownlee, John Wiley & Sons, 1965

**Example:**

x: 79, 84, 108, 114, 120, 103, 122, 120

y: 91, 103, 90, 113, 108, 87, 100, 80, 99, 54

$$n_1 = 8$$

$$n_2 = 10$$

If D = 0 (i.e.,  $H_0: \mu_1 = \mu_2$ )

then  $t = 1.73$ ,  $df = 16.00$

LINE	INSTRUCTIONS	DATA	KEYS	DISPLAY
1	Enter program			
2	Initialize		A	
3	Perform 3 for i=1, 2, ..., n <sub>1</sub>	x <sub>i</sub>	B	i
	(Correct erroneous data x <sub>k</sub> )	x <sub>k</sub>	E	
4		D	C R/S	
5	Perform 5 for j=1, 2, ..., n <sub>2</sub>	y <sub>j</sub>	B	j
	(Correct erroneous data y <sub>h</sub> )	y <sub>h</sub>	E	
6			D	t
7			R/S	df
	(For a different value of D)	D	C	
			D	t
			R/S	df

**CHI-SQUARE EVALUATION**

O <sub>i</sub> , E <sub>i</sub>	Σ-	O <sub>i</sub>	Σ-(O <sub>i</sub> )	χ <sup>2</sup>	
---------------------------------	----	----------------	---------------------	----------------	---

This program calculates the value of the  $\chi^2$  statistic for the goodness of fit test by the equation

$$\chi^2 = \sum_{i=1}^n \frac{(O_i - E_i)^2}{E_i}$$

where  $O_i$  = observed frequency

$E_i$  = expected frequency

If the expected values are equal

$$\left( E = E_i = \frac{\Sigma O_i}{n} \text{ for all } i \right)$$

then

$$\chi^2 = \frac{n \Sigma O_i^2}{\Sigma O_i} - \Sigma O_i$$

**Note:** In order to apply the goodness of fit test to a set of given data, combining some classes may be necessary to make sure that each expected frequency is not too small (say, not less than 5).

**Reference:** Mathematical Statistics, J. E. Freund, Prentice Hall, 1962

**Examples:**

1.	O <sub>i</sub>		8	50	47	56	5	14
	E <sub>i</sub>		9.6	46.75	51.85	54.4	8.25	9.15

$$\chi^2 = 4.84$$

2. The following table shows the observed frequencies in tossing a die 120 times.  $\chi^2$  can be used to test if the die is fair.

**Note:** Assume that the expected frequencies are equal.

number	1	2	3	4	5	6
frequency $O_i$	25	17	15	23	24	16

$$\chi^2 = 5.00$$

$$E = 20.00$$

LINE	INSTRUCTIONS	DATA	KEYS	DISPLAY
1	Enter program			
2	Initialize		RTN R/S	
3	For equal expected values,			
	go to 7.			
4	Perform 4–5 for $i = 1, 2, \dots, n$	$O_i$	$\uparrow$	
5		$E_i$	A	$i$
	(Correct erroneous data $O_k, E_k$ )	$O_k$	$\uparrow$	
		$E_k$	B	
6			E	$\chi^2$
	(For a new case, go to 2)			
7	Perform 7 for $i = 1, 2, \dots, n$	$O_i$	C	$i$
	(Correct erroneous data $O_h$ )	$O_h$	D	
8			f SF 1	
9			E	$\chi^2$
10			R/S	E
	(For a new case, go to 2)			

**2 x k CONTINGENCY TABLE**

2xk CONTINGENCY TABLE					STAT 1-32A
INIT	$a_i, b_i$	$\chi^2$	df	C	

Contingency tables can be used to test the null hypothesis that two variables are independent.

	1	2	3	...	k	Totals
A	$a_1$	$a_2$	$a_3$	...	$a_k$	$N_A$
B	$b_1$	$b_2$	$b_3$	...	$b_k$	$N_B$
Totals	$N_1$	$N_2$	$N_3$	...	$N_k$	$N$

Test statistic

$$\chi^2 = \frac{N}{N_A} \sum_{i=1}^k \frac{a_i^2}{N_i} + \frac{N}{N_B} \sum_{i=1}^k \frac{b_i^2}{N_i} - N$$

Degrees of freedom  $df = k - 1$

Pearson's coefficient of contingency C measures the degree of association between the two variables

$$C = \sqrt{\frac{\chi^2}{N + \chi^2}}$$

**Reference:** Statistics in Research, B. Ostle, Iowa State University Press, 1963

**Example:**

	1	2	3
A	2	5	4
B	3	8	7

$$\chi^2 = 0.02$$

$$df = 2.00$$

$$C = 0.03$$

LINE	INSTRUCTIONS	DATA	KEYS	DISPLAY
1	Enter program			
2	Initialize		A	
3	Perform 3–4 for i=1, 2, ..., k	a <sub>i</sub>	↑	
4		b <sub>i</sub>	B	i
5			C	$\chi^2$
6			D	df
7			E	C

**BARTLETT'S CHI-SQUARE STATISTIC**

BARTLETT'S CHI-SQUARE STATISTIC			STAT 1-33A
INIT	$\Sigma^+$	$\chi^2$	$\Sigma^-$

$$\chi^2 = \frac{f \ln s^2 - \sum_{i=1}^k f_i \ln s_i^2}{1 + \frac{1}{3(k-1)} \left[ \left( \sum_{i=1}^k \frac{1}{f_i} \right) - \frac{1}{f} \right]}$$

where  $s_i^2$  = sample variance of the  $i^{\text{th}}$  sample

$f_i$  = degrees of freedom associated with  $s_i^2$

$i = 1, 2, \dots, k$

$k$  = number of samples

$$s^2 = \frac{\sum_{i=1}^k f_i s_i^2}{f}$$

$$f = \sum_{i=1}^k f_i$$

This  $\chi^2$  has a chi-square distribution (approximately) with  $k - 1$  degrees of freedom which can be used to test the null hypothesis that  $s_1^2, s_2^2, \dots, s_k^2$  are all estimates of the same population variance  $\sigma^2$ ; i.e.

$H_0$ : Each of  $s_1^2, s_2^2, \dots, s_k^2$  is an estimate of  $\sigma^2$ .

**Note:** Erroneous data can be corrected by using the **D** key.

**Reference:** Statistical Theory with Engineering Applications,  
A. Hald, John Wiley and Sons, 1960

**Example:**

i	1	2	3	4	5	6
$s_i^2$	5.5	5.1	5.2	4.7	4.8	4.3
$f_i$	10	20	17	18	8	15

$$\chi^2 = 0.25$$

$$df = 5.00$$

LINE	INSTRUCTIONS	DATA	KEYS	DISPLAY
1	Enter program			
2	Initialize		A	
3	Perform 3–4 for $i=1, 2, \dots, k$	$s_i^2$	↑	
4		$f_i$	B	i
	(Correct erroneous data $s_m^2, f_m$ )	$s_m^2$	↑	
		$f_m$	D	
5			C	$\chi^2$
6			R/S	df

**SPEARMAN'S RANK CORRELATION COEFFICIENT**

Spearman's rank correlation coefficient is defined by

$$r_s = 1 - \frac{6 \sum_{i=1}^n D_i^2}{n(n^2 - 1)}$$

where  $n$  = number of paired observations  $(x_i, y_i)$

$$D_i = \text{rank}(x_i) - \text{rank}(y_i) = R_i - S_i$$

If the X and Y random variables from which these n pairs of observations are derived are independent, then  $r_s$  has zero mean and a variance

$$\frac{1}{n - 1}$$

A test for the null hypothesis

$$H_0: X, Y \text{ are independent}$$

is using

$$z = r_s \sqrt{n - 1}$$

which is approximately a standardized normal variable (for large  $n$ , say  $n \geq 10$ ).

If the null hypothesis of independence is not rejected, we can infer that the population correlation coefficient  $\rho(x, y) = 0$ , but dependence between the variables does not necessarily imply that  $\rho(x, y) \neq 0$ .

**Note:**  $-1 \leq r_s \leq 1$

$r_s = 1$  indicates complete agreement in order of the ranks and  $r_s = -1$  indicates complete agreement in the opposite order of the ranks.

**Reference:** Nonparametric Statistical Inference, J. D. Gibbons, McGraw Hill, 1971


**Example:**

Student	$x_i$ Math Grade	$y_i$ Stat Grade	$R_i$ Rank of $x_i$	$S_i$ Rank of $y_i$
1	82	81	6	7
2	67	75	14	11
3	91	85	3	4
4	98	90	1	2
5	74	80	11	8
6	52	60	15	15
7	86	94	4	1
8	95	78	2	9
9	79	83	9	6
10	78	76	10	10
11	84	84	5	5
12	80	69	8	13
13	69	72	13	12
14	81	88	7	3
15	73	61	12	14

$$r_s = 0.76$$

$$z = 2.85$$

LINE	INSTRUCTIONS	DATA	KEYS	DISPLAY
1	Enter program			
2	Initialize		A	
3	Perform 3–4 for $i = 1, 2, \dots, n$	$R_i$	↑	
4		$S_i$	B	i
	(Correct erroneous data $R_k, S_k$ )	$R_k$	↑	
		$S_k$	E	
5			C	$r_s$
6			D	z

**MANN-WHITNEY STATISTIC**

This program computes the Mann-Whitney test statistic on two independent samples of equal or unequal sizes. This test is designed for testing the null hypothesis of no difference between two populations.

Mann-Whitney test statistic is defined as

$$U = n_1 n_2 + \frac{n_1 (n_1 + 1)}{2} - \sum_{i=1}^{n_1} R_i$$

where  $n_1$  and  $n_2$  are the sizes of the two samples. Arrange all values from both samples jointly (as if they were one sample) in an increasing order of magnitude, let  $R_i$  ( $i = 1, 2, \dots, n_1$ ) be the ranks assigned to the values of the first sample (it is immaterial which sample is referred to as the “first”).

When  $n_1$  and  $n_2$  are small, the Mann-Whitney test bases on the exact distribution of  $U$  and specially constructed tables. When  $n_1$  and  $n_2$  are both large (say, greater than 8) then

$$z = \frac{U - \frac{n_1 n_2}{2}}{\sqrt{n_1 n_2 (n_1 + n_2 + 1)/12}}$$

is approximately a random variable having the standard normal distribution.

**Reference:** Mathematical Statistics, J.E. Freund, Prentice Hall, 1962

**Table for small samples:**

Handbook of Statistical Tables, D.B. Owen, Addison-Wesley, 1962

**Example:**

Sample 1	14.9	11.3	13.2	16.6	17	14.1	15.4	13	16.9
Rank $R_i$	7	1	4	12	14	5	10	3	13
Sample 2	15.2	19.8	14.7	18.3	16.2	21.2	18.9	12.2	15.3
Rank	8	18	6	15	11	19	16	2	9

$$n_1 = 9, \quad n_2 = 10$$

$$U = 66.00$$

$$z = 1.71$$

LINE	INSTRUCTIONS	DATA	KEYS	DISPLAY
1	Enter program			
2		$n_2$	A	
3	Perform 3 for $i=1, 2, \dots, n_1$	$R_i$	B	i
	(Correct erroneous data $R_k$ )	$R_k$	E	
4	Compute U		C	U
5	Compute z		D	z

**KENDALL'S COEFFICIENT OF CONCORDANCE**

KENDALL'S COEFFICIENT OF CONCORDANCE			STAT 1-36A		
$\Sigma^+$	$\Sigma\Sigma^+$	W	$\chi^2, df$	$\Sigma^-$	

Suppose  $n$  individuals are ranked from 1 to  $n$  according to some specified characteristic by  $k$  observers, the coefficient of concordance  $W$  measures the agreement between observers (or concordance between rankings).

$$W = \frac{12 \sum_{i=1}^n \left( \sum_{j=1}^k R_{ij} \right)^2}{k^2 n(n^2 - 1)} - \frac{3(n + 1)}{n - 1}$$

Where  $R_{ij}$  is the rank assigned to the  $i^{\text{th}}$  individual by the  $j^{\text{th}}$  observer.

$W$  varies from 0 (no community of preference) to 1 (perfect agreement). The null hypothesis that the observers have no community of preference may be tested using special tables, or if  $n > 7$ , by computing

$$\chi^2 = k(n - 1)W$$

which has approximately the chi-square distribution with  $n - 1$  degrees of freedom (df).

**Reference:** Nonparametric Statistical Inference, J. D. Gibbons, McGraw-Hill, 1971

**Table for small samples:**

Rank Correlation Methods, M.G. Kendall, Hafner Publishing Co., 1962

**Example:**

Table for  $R_{ij}$  ( $n = 10, k = 3$ )

i \ j	1	2	3
1	6	7	3
2	1	4	2
3	9	3	5
4	2	6	1
5	10	8	9
6	3	2	6
7	5	9	8
8	4	1	4
9	8	10	10
10	7	5	7

$$W = 0.69$$

$$\chi^2 = 18.64$$

$$df = 9.00$$

LINE	INSTRUCTIONS	DATA	KEYS	DISPLAY
1	Enter program			
2	Initialize		RTN R/S	
3	Perform 3–5 for $i=1, 2, \dots, n$			
4	Perform 4 for $j=1, 2, \dots, k$	$R_{ij}$	A	j
	(Correct erroneous data $R_{im}$ )	$R_{im}$	E	
5			B	i
6	Compute W		C	W
7	Compute $\chi^2$ and df		D R/S	$\chi^2$ df
8				
	(For a new case, go to 2)			

**BISERIAL CORRELATION COEFFICIENT**

BISERIAL CORRELATION COEFFICIENT			STAT 1-37A
INIT	$x_i=1$	$x_i=0$	$r_b$

The biserial correlation coefficient  $r_b$  is used where one variable Y is quantitatively measured while the other continuous variable X is artificially dichotomized (that is, artificially defined by two groups). It measures the degree of linear association between X and Y.

$$r_b = \frac{n (\Sigma' y_i) - n_1 \sum y_i}{na \sqrt{n \sum y_i^2 - (\sum y_i)^2}}$$

Suppose X takes the value 0 or 1.

Define  $n_1$  = number of x's such that  $x = 1$


$n$  = total number of data points

$\Sigma' y_i$  = sum of the y's for which  $x = 1$

$\sum y_i$  = sum of all y's

$a$  = ordinate of the standard normal curve at point  $z$   
cutting off a tail of that distribution with area

$$\text{equal to } p = \frac{n_1}{n}.$$



- Notes:**
- $p = \frac{n_1}{n}$  must be less than or equal to 0.5, if not, interchange the roles of 0 and 1 for the X variable.
  - z and a can be found by using *Stat 1-10A, Normal Distribution*, and *Stat 1-11A, Inverse Normal Integral*.

3. Among the necessary assumptions for a meaningful interpretation of  $r_b$  are:

- (a) Y is normally distributed
- (b) the true distribution of X should be of normal form.

**Reference:** Statistics in Research, B. Ostle, Iowa State University Press, 1963

**Example:**

$x_i$	0	1	1	0	1	0	0	0	1
$y_i$	3.1	2.8	5.6	0.3	2.5	2.4	4.8	2.9	7.7

$$n_1 = 4$$

$$n = 9$$

$$z = 0.14$$

$$a = 0.40$$

$$r_b = 0.60$$

LINE	INSTRUCTIONS	DATA	KEYS	DISPLAY
1	Enter program Stat 1-11A			
2			A	
3		$n_1$	$\uparrow$	
4		$n$	$\div$	B
5			STO	1
6	Enter program Stat 1-10A1			
7			A	
8	Enter program Stat 1-10A2			
9			RCL	1
10			A	
11	Enter program Stat 1-37A			a
12			A	
13	Perform 14 or 15 for $i=1, \dots, n$			
14	If $x_i = 1$	$y_i$	B	
15	If $x_i = 0$	$y_i$	C	
16			D	
				$r_b$



**PROGRAM LISTINGS**

1. Mean, Standard Deviation, Standard Error . . . . .	84
2. Mean, Standard Deviation, Standard Error (Grouped Data) . . . . .	85
3. Permutation and Combination . . . . .	86
4. Arithmetic, Geometric, Harmonic and Generalized Means . . . . .	87
5. Sums for Two Variables . . . . .	88
6. Basic Statistics (Two Variables) . . . . .	89
7. Moments, Skewness and Kurtosis (For Grouped or Ungrouped Data) Card 1 . . . . .	90
Card 2 . . . . .	91
8. Random Number Generator . . . . .	92
9. Analysis of Variance (One Way) . . . . .	93
10. Normal Distribution Card 1 . . . . .	94
Card 2 . . . . .	95
11. Inverse Normal Integral . . . . .	96
12. Chi-Square Distribution . . . . .	97
13. t Distribution . . . . .	98
14. F Distribution . . . . .	99
15. Bivariate Normal Distribution . . . . .	100
16. Logarithmic Normal Distribution . . . . .	101
17. Weibull Distribution . . . . .	102
18. Binomial Distribution . . . . .	103
19. Negative Binomial Distribution . . . . .	104
20. Hypergeometric Distribution . . . . .	105
21. Poisson Distribution . . . . .	106
22. Linear Regression . . . . .	107
23. Exponential Curve Fit . . . . .	108
24. Power Curve Fit . . . . .	109
25. Logarithmic Curve Fit . . . . .	110
26. Least Squares Regression of $y = cx^a + dx^b$ . . . . .	111
27. Multiple Linear Regression Card 1 . . . . .	112
Card 2 . . . . .	113
28. Parabolic Curve Fit . . . . .	114
29. Paired t Statistic . . . . .	115
30. t Statistic for Two Means . . . . .	116
31. Chi-Square Evaluation . . . . .	117
32. $2 \times k$ Contingency Table . . . . .	118
33. Bartlett's Chi-Square Statistic . . . . .	119
34. Spearman's Rank Correlation Coefficient . . . . .	120
35. Mann-Whitney Statistic . . . . .	121
36. Kendall's Coefficient of Concordance . . . . .	122
37. Biserial Correlation Coefficient . . . . .	123

## MEAN, STANDARD DEVIATION, STANDARD ERROR

CODE	KEYS	CODE	KEYS	CODE	KEYS
00	0	71	x	02	2
33 01	STO 1	51	—	32	f <sup>-1</sup>
33 02	STO 2	34 01	RCL 1	09	$\sqrt{x}$
33 03	STO 3	81	$\div$	33	STO
84	R/S	31	f	51	—
23	LBL	09	$\sqrt{x}$	03	3
11	A	34 01	RCL 1	34 01	RCL 1
33	STO	34 01	RCL 1	01	1
61	+	01	1	51	—
02	2	51	—	33 01	STO 1
32	f <sup>-1</sup>	81	$\div$	24	RTN
09	$\sqrt{x}$	31	f	35 01	g NOP
33	STO	09	$\sqrt{x}$	35 01	g NOP
61	+	71	x	35 01	g NOP
03	3	24	RTN	35 01	g NOP
34 01	RCL 1	35 07	g x $\leftrightarrow$ y	35 01	g NOP
01	1	84	R/S	35 01	g NOP
61	+	23	LBL	35 01	g NOP
33 01	STO 1	14	D	35 01	g NOP
24	RTN	13	C	35 01	g NOP
23	LBL	34 01	RCL 1	35 01	g NOP
12	B	31	f	35 01	g NOP
34 02	RCL 2	09	$\sqrt{x}$	35 01	g NOP
34 01	RCL 1	81	$\div$	35 01	g NOP
81	$\div$	35 07	g x $\leftrightarrow$ y	35 01	g NOP
24	RTN	35 00	g LST X	35 01	g NOP
23	LBL	81	$\div$	35 01	g NOP
13	C	35 07	g x $\leftrightarrow$ y	35 01	g NOP
34 03	RCL 3	84	R/S	35 01	g NOP
34 02	RCL 2	35 07	g x $\leftrightarrow$ y	35 01	g NOP
34 01	RCL 1	24	RTN	35 01	g NOP
81	$\div$	23	LBL		
32	f <sup>-1</sup>	15	E		
09	$\sqrt{x}$	33	STO		
34 01	RCL 1	51	—		

<b>R<sub>1</sub></b>	n	<b>R<sub>4</sub></b>	<b>R<sub>7</sub></b>
<b>R<sub>2</sub></b>	$\Sigma x_i$	<b>R<sub>5</sub></b>	<b>R<sub>8</sub></b>
<b>R<sub>3</sub></b>	$\Sigma x_i^2$	<b>R<sub>6</sub></b>	<b>R<sub>9</sub></b>

**MEAN, STANDARD DEVIATION,  
STANDARD ERROR (GROUPED DATA)**

CODE	KEYS	CODE	KEYS	CODE	KEYS
00	0	34 02	RCL 2	35 07	g $x \leftrightarrow y$
33 01	STO 1	34 01	RCL 1	24	RTN
33 02	STO 2	81	$\div$	23	LBL
33 03	STO 3	32	$f^{-1}$	15	E
33 04	STO 4	09	$\sqrt{x}$	42	CHS
84	R/S	34 01	RCL 1	11	A
23	LBL	71	x	34 04	RCL 4
11	A	51	$-$	02	2
33	STO	34 01	RCL 1	51	$-$
61	+	81	$\div$	33 04	STO 4
01	1	31	f	24	RTN
35 07	g $x \leftrightarrow y$	09	$\sqrt{x}$	35 01	g NOP
71	x	34 01	RCL 1	35 01	g NOP
33	STO	34 01	RCL 1	35 01	g NOP
61	+	01	1	35 01	g NOP
02	2	51	$-$	35 01	g NOP
35 00	g LST X	81	$\div$	35 01	g NOP
71	x	31	f	35 01	g NOP
33	STO	09	$\sqrt{x}$	35 01	g NOP
61	+	71	x	35 01	g NOP
03	3	24	RTN	35 01	g NOP
01	1	35 07	g $x \leftrightarrow y$	35 01	g NOP
34 04	RCL 4	84	R/S	35 01	g NOP
61	+	23	LBL	35 01	g NOP
33 04	STO 4	14	D	35 01	g NOP
24	RTN	13	C	35 01	g NOP
23	LBL	34 01	RCL 1	35 01	g NOP
12	B	31	f	35 01	g NOP
34 02	RCL 2	09	$\sqrt{x}$	35 01	g NOP
34 01	RCL 1	81	$\div$	35 01	g NOP
81	$\div$	35 07	g $x \leftrightarrow y$	35 01	g NOP
24	RTN	35 00	g LST X	35 01	g NOP
23	LBL	81	$\div$		
13	C	35 07	g $x \leftrightarrow y$		
34 03	RCL 3	84	R/S		

<b>R<sub>1</sub></b>	$\sum f_i$	<b>R<sub>4</sub></b>	n	<b>R<sub>7</sub></b>
<b>R<sub>2</sub></b>	$\sum f_i x_i$	<b>R<sub>5</sub></b>		<b>R<sub>8</sub></b>
<b>R<sub>3</sub></b>	$\sum f_i x_i^2$	<b>R<sub>6</sub></b>		<b>R<sub>9</sub></b>

## PERMUTATION AND COMBINATION

CODE	KEYS	CODE	KEYS	CODE	KEYS
23	LBL	23	LBL	35 07	g $x \leftrightarrow y$
11	A	03	3	61	+
35 24	g $x > y$	41	$\uparrow$	35 00	g LST X
22	GTO	01	1	81	$\div$
02	2	24	RTN	34 06	RCL 6
41	$\uparrow$	23	LBL	71	x
00	0	12	B	33 06	STO 6
35 23	g $x = y$	35 24	g $x > y$	22	GTO
22	GTO	22	GTO	00	0
03	3	02	2	23	LBL
44	CLX	51	$-$	04	4
01	1	35 00	g LST X	35 08	g R↓
35 23	g $x = y$	35 22	g $x \leq y$	35 08	g R↑
22	GTO	33 06	STO 6	24	RTN
04	4	35 07	g $x \leftrightarrow y$	35 01	g NOP
51	$-$	33 07	STO 7	35 01	g NOP
33 08	STO 8	01	1	35 01	g NOP
35 08	g R↓	33 08	STO 8	35 01	g NOP
33 07	STO 7	61	+	35 01	g NOP
23	LBL	33 06	STO 6	35 01	g NOP
01	1	44	CLX	35 01	g NOP
34 07	RCL 7	35 23	g $x = y$	35 01	g NOP
01	1	01	1	35 01	g NOP
51	$-$	24	RTN	35 01	g NOP
33 07	STO 7	23	LBL	35 01	g NOP
71	x	00	0	35 01	g NOP
35	g	35 08	g R↓	35 01	g NOP
83	DSZ	01	1	35 01	g NOP
22	GTO	34 08	RCL 8	35 01	g NOP
01	1	61	$+$	35 01	g NOP
24	RTN	33 08	STO 8	35 01	g NOP
23	LBL	35 24	g $x > y$	35 01	g NOP
02	2	34 06	RCL 6	35 01	g NOP
00	0	24	RTN	35 01	g NOP
81	$\div$	34 07	RCL 7	35 01	g NOP

<b>R<sub>1</sub></b>	<b>R<sub>4</sub></b>	<b>R<sub>7</sub></b>	Used
<b>R<sub>2</sub></b>	<b>R<sub>5</sub></b>	<b>R<sub>8</sub></b>	Used
<b>R<sub>3</sub></b>	<b>R<sub>6</sub></b>	Used	<b>R<sub>9</sub></b>

**ARITHMETIC, GEOMETRIC, HARMONIC AND  
GENERALIZED MEANS**

CODE	KEYS	CODE	KEYS	CODE	KEYS
01	1	33 05	STO 5	35 01	g NOP
33 02	STO 2	24	RTN	35 01	g NOP
44	CLX	23	LBL	35 01	g NOP
33 01	STO 1	12	B	35 01	g NOP
33 03	STO 3	34 01	RCL 1	35 01	g NOP
33 04	STO 4	34 05	RCL 5	35 01	g NOP
33 05	STO 5	81	÷	35 01	g NOP
84	R/S	24	RTN	35 01	g NOP
33	STO	23	LBL	35 01	g NOP
09	9	13	C	35 01	g NOP
84	R/S	34 02	RCL 2	35 01	g NOP
23	LBL	34 05	RCL 5	35 01	g NOP
11	A	35	g	35 01	g NOP
33	STO	04	${}^1/\chi$	35 01	g NOP
61	+	35	g	35 01	g NOP
01	1	05	$y^x$	35 01	g NOP
33	STO	24	RTN	35 01	g NOP
71	x	23	LBL	35 01	g NOP
02	2	14	D	35 01	g NOP
35	g	34 05	RCL 5	35 01	g NOP
04	${}^1/\chi$	34 03	RCL 3	35 01	g NOP
33	STO	81	÷	35 01	g NOP
61	+	24	RTN	35 01	g NOP
03	3	23	LBL	35 01	g NOP
35 00	g LST X	15	E	35 01	g NOP
34	RCL	34 04	RCL 4	35 01	g NOP
09	9	34 05	RCL 5	35 01	g NOP
35	g	81	÷	35 01	g NOP
05	$y^x$	34	RCL	35 01	g NOP
33	STO	09	9	35 01	g NOP
61	+	35	g	35 01	g NOP
04	4	04	${}^1/\chi$		
01	1	35	g		
34 05	RCL 5	05	$y^x$		
61	+	24	RTN		

<b>R<sub>1</sub></b>	$\Sigma a$	<b>R<sub>4</sub></b>	$\Sigma a^t$	<b>R<sub>7</sub></b>
<b>R<sub>2</sub></b>	$\Pi a$	<b>R<sub>5</sub></b>	n	<b>R<sub>8</sub></b>
<b>R<sub>3</sub></b>	$\Sigma {}^1/a$	<b>R<sub>6</sub></b>		<b>R<sub>9</sub></b> t

## SUMS FOR TWO VARIABLES

CODE	KEYS	CODE	KEYS	CODE	KEYS
23	LBL	51	—	34 04	RCL 4
11	A	04	4	84	R/S
33 07	STO 7	32	f <sup>-1</sup>	34 05	RCL 5
33	STO	09	$\sqrt{x}$	84	R/S
61	+	33	STO	34 06	RCL 6
04	4	51	—	84	R/S
32	f <sup>-1</sup>	05	5	35 01	g NOP
09	$\sqrt{x}$	35 07	g x $\leftrightarrow$ y	35 01	g NOP
33	STO	33	STO	35 01	g NOP
61	+	51	—	35 01	g NOP
05	5	02	2	35 01	g NOP
35 07	g x $\leftrightarrow$ y	32	f <sup>-1</sup>	35 01	g NOP
33	STO	09	$\sqrt{x}$	35 01	g NOP
61	+	33	STO	35 01	g NOP
02	2	51	—	35 01	g NOP
32	f <sup>-1</sup>	03	3	35 01	g NOP
09	$\sqrt{x}$	35 00	g LST X	35 01	g NOP
33	STO	34 07	RCL 7	35 01	g NOP
61	+	71	x	35 01	g NOP
03	3	33	STO	35 01	g NOP
35 00	g LST X	51	—	35 01	g NOP
34 07	RCL 7	06	6	35 01	g NOP
71	x	34 01	RCL 1	35 01	g NOP
33	STO	01	1	35 01	g NOP
61	+	51	—	35 01	g NOP
06	6	33 01	STO 1	35 01	g NOP
34 01	RCL 1	24	RTN	35 01	g NOP
01	1	23	LBL	35 01	g NOP
61	+	13	C	35 01	g NOP
33 01	STO 1	34 01	RCL 1	35 01	g NOP
24	RTN	84	R/S		
23	LBL	34 02	RCL 2		
12	B	84	R/S		
33 07	STO 7	34 03	RCL 3		
33	STO	84	R/S		

R <sub>1</sub>	n	R <sub>4</sub>	$\Sigma y_i$	R <sub>7</sub>	Used
R <sub>2</sub>	$\Sigma x_i$	R <sub>5</sub>	$\Sigma y_i^2$	R <sub>8</sub>	
R <sub>3</sub>	$\Sigma x_i^2$	R <sub>6</sub>	$\Sigma x_i y_i$	R <sub>9</sub>	

## BASIC STATISTICS (TWO VARIABLES)

CODE	KEYS	CODE	KEYS	CODE	KEYS
23	LBL	84	R/S	32	f <sup>-1</sup>
11	A	35 07	g x $\leftrightarrow$ y	09	$\sqrt{x}$
34 02	RCL 2	24	RTN	34 01	RCL 1
34 01	RCL 1	23	LBL	71	x
81	$\div$	13	C	51	—
84	R/S	34 06	RCL 6	34 01	RCL 1
34 04	RCL 4	34 02	RCL 2	01	1
34 01	RCL 1	34 04	RCL 4	51	—
81	$\div$	71	x	81	$\div$
24	RTN	34 01	RCL 1	31	f
23	LBL	81	$\div$	09	$\sqrt{x}$
12	B	51	—	24	RTN
34 03	RCL 3	34 01	RCL 1	35 01	g NOP
34 02	RCL 2	01	1	35 01	g NOP
15	E	51	—	35 01	g NOP
33 07	STO 7	81	$\div$	35 01	g NOP
84	R/S	24	RTN	35 01	g NOP
34 05	RCL 5	34	RCL	35 01	g NOP
34 04	RCL 4	09	9	35 01	g NOP
15	E	32	f <sup>-1</sup>	35 01	g NOP
33 08	STO 8	09	$\sqrt{x}$	35 01	g NOP
84	R/S	71	x	35 01	g NOP
34 01	RCL 1	84	R/S	35 01	g NOP
01	1	23	LBL	35 01	g NOP
51	—	14	D	35 01	g NOP
34 01	RCL 1	13	C	35 01	g NOP
81	$\div$	34 07	RCL 7	35 01	g NOP
31	f	34 08	RCL 8	35 01	g NOP
09	$\sqrt{x}$	71	x	35 01	g NOP
33	STO	81	$\div$	35 01	g NOP
09	9	24	RTN		
71	x	23	LBL		
34 07	RCL 7	15	E		
35 00	g LST X	34 01	RCL 1		
71	x	81	$\div$		

<b>R<sub>1</sub></b>	n	<b>R<sub>4</sub></b>	$\Sigma y_i$	<b>R<sub>7</sub></b>	$s_x$
<b>R<sub>2</sub></b>	$\Sigma x_i$	<b>R<sub>5</sub></b>	$\Sigma y_i^2$	<b>R<sub>8</sub></b>	$s_y$
<b>R<sub>3</sub></b>	$\Sigma x_i^2$	<b>R<sub>6</sub></b>	$\Sigma x_i y_i$	<b>R<sub>9</sub></b>	$[(n-1)/n]^{1/2}$

**MOMENTS, SKEWNESS AND KURTOSIS  
(FOR GROUPED OR UNGROUPED DATA) (CARD 1)**

CODE	KEYS	CODE	KEYS	CODE	KEYS
33	STO	33	STO	33	STO
61	+	51	—	61	+
02	2	04	4	05	5
32	f <sup>-1</sup>	35 00	g LST X	24	RTN
09	√x	71	x	23	LBL
33	STO	33	STO	14	D
61	+	51	—	33	STO
03	3	05	5	51	—
35 00	g LST X	34 01	RCL 1	01	1
71	x	01	1	35 07	g x↔y
33	STO	51	—	71	x
61	+	33 01	STO 1	33	STO
04	4	24	RTN	51	—
35 00	g LST X	23	LBL	02	2
71	x	13	C	35 00	g LST X
33	STO	33	STO	71	x
61	+	61	+	33	STO
05	5	01	1	51	—
01	1	35 07	g x↔y	03	3
34 01	RCL 1	71	x	35 00	g LST X
61	+	33	STO	71	x
33 01	STO 1	61	+	33	STO
84	R/S	02	2	51	—
23	LBL	35 00	g LST X	04	4
12	B	71	x	35 00	g LST X
33	STO	33	STO	71	x
51	—	61	+	33	STO
02	2	03	3	51	—
32	f <sup>-1</sup>	35 00	g LST X	05	5
09	√x	71	x	24	RTN
33	STO	33	STO		
51	—	61	+		
03	3	04	4		
35 00	g LST X	35 00	g LST X		
71	x	71	x		

<b>R<sub>1</sub></b>	n or $\sum f_j$	<b>R<sub>4</sub></b>	$\sum x_i^3$ or $\sum f_j y_j^3$	<b>R<sub>7</sub></b>
<b>R<sub>2</sub></b>	$\sum x_i$ or $\sum f_j y_j$	<b>R<sub>5</sub></b>	$\sum x_i^4$ or $\sum f_j y_j^4$	<b>R<sub>8</sub></b>
<b>R<sub>3</sub></b>	$\sum x_i^2$ or $\sum f_j y_j^2$	<b>R<sub>6</sub></b>		<b>R<sub>9</sub></b>

**MOMENTS, SKEWNESS AND KURTOSIS  
(FOR GROUPED OR UNGROUPED DATA) (CARD 2)**

CODE	KEYS	CODE	KEYS	CODE	KEYS
23	LBL	61	+	83	.
11	A	33	STO	05	5
34 02	RCL 2	09	9	35	g
34 01	RCL 1	24	RTN	05	$y^x$
81	$\div$	23	LBL	81	$\div$
33 06	STO 6	14	D	84	R/S
24	RTN	34 05	RCL 5	34 06	RCL 6
23	LBL	34 06	RCL 6	34 07	RCL 7
12	B	34 04	RCL 4	32	$f^{-1}$
34 03	RCL 3	71	x	09	$\sqrt{x}$
34 01	RCL 1	04	4	81	$\div$
81	$\div$	71	x	24	RTN
34 06	RCL 6	51	—	35 01	g NOP
32	$f^{-1}$	34 08	RCL 8	35 01	g NOP
09	$\sqrt{x}$	34 03	RCL 3	35 01	g NOP
33 08	STO 8	71	x	35 01	g NOP
51	—	06	6	35 01	g NOP
33 07	STO 7	71	x	35 01	g NOP
24	RTN	61	+	35 01	g NOP
23	LBL	34 01	RCL 1	35 01	g NOP
13	C	81	$\div$	35 01	g NOP
34 04	RCL 4	34 08	RCL 8	35 01	g NOP
34 03	RCL 3	32	$f^{-1}$	35 01	g NOP
34 06	RCL 6	09	$\sqrt{x}$	35 01	g NOP
71	x	03	3	35 01	g NOP
03	3	71	x	35 01	g NOP
71	x	51	—	35 01	g NOP
51	—	33 06	STO 6	35 01	g NOP
34 01	RCL 1	24	RTN	35 01	g NOP
81	$\div$	23	LBL	35 01	g NOP
34 06	RCL 6	15	E	35 01	g NOP
34 08	RCL 8	34	RCL	35 01	g NOP
71	x	09	9	35 01	g NOP
02	2	34 07	RCL 7	35 01	g NOP
71	x	01	1	35 01	g NOP

<b>R<sub>1</sub></b>	n or $\sum f_j$	<b>R<sub>4</sub></b>	$\sum x_i^3$ or $\sum f_j y_j^3$	<b>R<sub>7</sub></b>	$m_2$
<b>R<sub>2</sub></b>	$\sum x_i$ or $\sum f_j y_j$	<b>R<sub>5</sub></b>	$\sum x_i^4$ or $\sum f_j y_j^4$	<b>R<sub>8</sub></b>	$\bar{x}^2$
<b>R<sub>3</sub></b>	$\sum x_i^2$ or $\sum f_j y_j^2$	<b>R<sub>6</sub></b>	$\bar{x}, m_4$	<b>R<sub>9</sub></b>	$m_3$

## RANDOM NUMBER GENERATOR

CODE	KEYS	CODE	KEYS	CODE	KEYS
23	LBL	31	f	34 01	RCL 1
11	A	07	LN	33 06	STO 6
00	0	02	2	34 02	RCL 2
33 01	STO 1	71	x	33 01	STO 1
24	RTN	42	CHS	12	B
23	LBL	31	f	33 02	STO 2
12	B	09	$\sqrt{x}$	34 06	RCL 6
35	g	33 05	STO 5	33 01	STO 1
02	$\pi$	34 01	RCL 1	12	B
34 01	RCL 1	35	g	34 05	RCL 5
61	+	02	$\pi$	84	R/S
08	8	71	x	35 01	g NOP
35	g	02	2	35 01	g NOP
05	$y^x$	71	x	35 01	g NOP
32	$f^{-1}$	33 06	STO 6	35 01	g NOP
83	INT	31	f	35 01	g NOP
33 01	STO 1	05	COS	35 01	g NOP
24	RTN	71	x	35 01	g NOP
23	LBL	34 04	RCL 4	35 01	g NOP
13	C	71	x	35 01	g NOP
33 04	STO 4	34 03	RCL 3	35 01	g NOP
35 07	$g \leftrightarrow y$	61	+	35 01	g NOP
33 03	STO 3	84	R/S	35 01	g NOP
24	RTN	23	LBL	35 01	g NOP
23	LBL	15	E	35 01	g NOP
14	D	34 06	RCL 6	35 01	g NOP
33 01	STO 1	31	f	35 01	g NOP
35 07	$g \leftrightarrow y$	04	SIN	35 01	g NOP
33 02	STO 2	34 05	RCL 5	35 01	g NOP
24	RTN	71	x	35 01	g NOP
23	LBL	34 04	RCL 4		
15	E	71	x		
35	g	34 03	RCL 3		
42	RAD	61	+		
34 02	RCL 2	33 05	STO 5		

<b>R<sub>1</sub></b>	Used	<b>R<sub>4</sub></b>	$\sigma$	<b>R<sub>7</sub></b>	
<b>R<sub>2</sub></b>	Used	<b>R<sub>5</sub></b>	Used	<b>R<sub>8</sub></b>	
<b>R<sub>3</sub></b>	m	<b>R<sub>6</sub></b>	Used	<b>R<sub>9</sub></b>	Used

## ANALYSIS OF VARIANCE (ONE WAY)

CODE	KEYS	CODE	KEYS	CODE	KEYS
23	LBL	33	STO	34 05	RCL 5
11	A	61	+	34 04	RCL 4
33	STO	05	5	51	—
61	+	00	0	33	STO
01	1	33 01	STO 1	09	9
32	f <sup>-1</sup>	33 02	STO 2	81	÷
09	√x	34 08	RCL 8	81	÷
33	STO	84	R/S	84	R/S
61	+	23	LBL	34 08	RCL 8
06	6	13	C	84	R/S
01	1	34 06	RCL 6	34	RCL
34 02	RCL 2	34 07	RCL 7	09	9
61	+	32	f <sup>-1</sup>	84	R/S
33 02	STO 2	09	√x	23	LBL
84	R/S	34 05	RCL 5	14	D
23	LBL	81	÷	33	STO
12	B	51	—	51	—
01	1	33 01	STO 1	01	1
33	STO	34 03	RCL 3	32	f <sup>-1</sup>
61	+	34 07	RCL 7	09	√x
04	4	32	f <sup>-1</sup>	33	STO
34 01	RCL 1	09	√x	51	—
32	f <sup>-1</sup>	34 05	RCL 5	06	6
09	√x	81	÷	34 02	RCL 2
34 02	RCL 2	51	—	01	1
81	÷	33 02	STO 2	51	—
33	STO	51	—	33 02	STO 2
61	+	33 03	STO 3	84	R/S
03	3	35 00	g LST X	35 01	g NOP
34 01	RCL 1	34 04	RCL 4	35 01	g NOP
33 08	STO 8	01	1		
33	STO	51	—		
61	+	33 08	STO 8		
07	7	81	÷		
34 02	RCL 2	35 07	g x $\rightleftharpoons$ y		

<b>R<sub>1</sub></b>	Used	<b>R<sub>4</sub></b>	Used	<b>R<sub>7</sub></b>	$\Sigma \Sigma x_{ij}$
<b>R<sub>2</sub></b>	Used	<b>R<sub>5</sub></b>	$\Sigma n_i$	<b>R<sub>8</sub></b>	$df_1$
<b>R<sub>3</sub></b>	Used	<b>R<sub>6</sub></b>	$\Sigma \Sigma x_{ij}^2$	<b>R<sub>9</sub></b>	$df_2$

## NORMAL DISTRIBUTION (CARD 1)

CODE	KEYS	CODE	KEYS	CODE	KEYS
23	LBL	33 05	STO 5	24	RTN
11	A	01	1	35 01	g NOP
83	.	83	.	35 01	g NOP
02	2	07	7	35 01	g NOP
03	3	08	8	35 01	g NOP
01	1	01	1	35 01	g NOP
06	6	04	4	35 01	g NOP
04	4	07	7	35 01	g NOP
01	1	07	7	35 01	g NOP
09	9	09	9	35 01	g NOP
33 03	STO 3	03	3	35 01	g NOP
01	1	07	7	35 01	g NOP
83	.	33 06	STO 6	35 01	g NOP
03	3	83	.	35 01	g NOP
03	3	03	3	35 01	g NOP
00	0	05	5	35 01	g NOP
02	2	06	6	35 01	g NOP
07	7	05	5	35 01	g NOP
04	4	06	6	35 01	g NOP
04	4	03	3	35 01	g NOP
02	2	07	7	35 01	g NOP
09	9	08	8	35 01	g NOP
33 04	STO 4	02	2	35 01	g NOP
01	1	42	CHS	35 01	g NOP
83	.	33 07	STO 7	35 01	g NOP
08	8	83	.	35 01	g NOP
02	2	03	3	35 01	g NOP
01	1	01	1	35 01	g NOP
02	2	09	9	35 01	g NOP
05	5	03	3	35 01	g NOP
05	5	08	8	35 01	g NOP
09	9	01	1		
07	7	05	5		
08	8	03	3		
42	CHS	33 08	STO 8		

<b>R<sub>1</sub></b>	<b>R<sub>4</sub></b>	b <sub>5</sub>	<b>R<sub>7</sub></b>	b <sub>2</sub>
<b>R<sub>2</sub></b>	<b>R<sub>5</sub></b>	b <sub>4</sub>	<b>R<sub>8</sub></b>	b <sub>1</sub>
<b>R<sub>3</sub></b>	r	b <sub>3</sub>	<b>R<sub>9</sub></b>	

## NORMAL DISTRIBUTION (CARD 2)

CODE	KEYS	CODE	KEYS	CODE	KEYS
23	LBL	41	↑	35 01	g NOP
11	A	41	↑	35 01	g NOP
33 01	STO 1	41	↑	35 01	g NOP
41	↑	34 04	RCL 4	35 01	g NOP
71	x	71	x	35 01	g NOP
02	2	34 05	RCL 5	35 01	g NOP
81	÷	61	+	35 01	g NOP
42	CHS	71	x	35 01	g NOP
32	f <sup>-1</sup>	34 06	RCL 6	35 01	g NOP
07	LN	61	+	35 01	g NOP
35	g	71	x	35 01	g NOP
02	π	34 07	RCL 7	35 01	g NOP
02	2	61	+	35 01	g NOP
71	x	71	x	35 01	g NOP
31	f	34 08	RCL 8	35 01	g NOP
09	√x	61	+	35 01	g NOP
81	÷	71	x	35 01	g NOP
33 02	STO 2	34 02	RCL 2	35 01	g NOP
24	RTN	71	x	35 01	g NOP
23	LBL	24	RTN	35 01	g NOP
12	B	23	LBL	35 01	g NOP
34 01	RCL 1	01	1	35 01	g NOP
00	0	34 01	RCL 1	35 01	g NOP
35 24	g x>y	42	CHS	35 01	g NOP
22	GTO	33 01	STO 1	35 01	g NOP
01	1	13	C	35 01	g NOP
23	LBL	01	1	35 01	g NOP
13	C	35 07	g x↔y	35 01	g NOP
01	1	51	—	35 01	g NOP
34 01	RCL 1	24	RTN	35 01	g NOP
34 03	RCL 3	35 01	g NOP	35 01	g NOP
71	x	35 01	g NOP	35 01	g NOP
61	+	35 01	g NOP	35 01	g NOP
35	g	35 01	g NOP	35 01	g NOP
04	¹/x	35 01	g NOP	35 01	g NOP

<b>R<sub>1</sub></b>	x or -x	<b>R<sub>4</sub></b>	b <sub>5</sub>	<b>R<sub>7</sub></b>	b <sub>2</sub>
<b>R<sub>2</sub></b>	f(x)	<b>R<sub>5</sub></b>	b <sub>4</sub>	<b>R<sub>8</sub></b>	b <sub>1</sub>
<b>R<sub>3</sub></b>	r	<b>R<sub>6</sub></b>	b <sub>3</sub>	<b>R<sub>9</sub></b>	Used

## INVERSE NORMAL INTEGRAL

CODE	KEYS	CODE	KEYS	CODE	KEYS
02	2	01	1	07	LN
83	.	08	8	31	f
05	5	09	9	09	$\sqrt{x}$
01	1	02	2	33 07	STO 7
05	5	06	6	34 03	RCL 3
05	5	09	9	71	x
01	1	33 05	STO 5	34 02	RCL 2
07	7	83	.	61	+
33 01	STO 1	00	0	34 07	RCL 7
83	.	00	0	71	x
08	8	01	1	34 01	RCL 1
00	0	03	3	61	+
02	2	00	0	34 07	RCL 7
08	8	08	8	34 06	RCL 6
05	5	33 06	STO 6	71	x
03	3	24	RTN	34 05	RCL 5
33 02	STO 2	23	LBL	61	+
83	.	12	B	34 07	RCL 7
00	0	41	$\uparrow$	71	x
01	1	00	0	34 04	RCL 4
00	0	35 24	$g\ x > y$	61	+
03	3	00	0	34 07	RCL 7
02	2	81	$\div$	71	x
08	8	35 08	$g\ R \downarrow$	01	1
33 03	STO 3	83	.	61	+
01	1	05	5	81	$\div$
83	.	35 07	$g\ x \leftrightarrow y$	34 07	RCL 7
04	4	35 24	$g\ x > y$	35 07	$g\ x \leftrightarrow y$
03	3	00	0	51	—
02	2	81	$\div$	24	RTN
07	7	41	$\uparrow$		
08	8	71	x		
08	8	35	$g$		
33 04	STO 4	04	${}^1/x$		
83	.	31	f		

$R_1$	$c_0$	$R_4$	$d_1$	$R_7$	$t$
$R_2$	$c_1$	$R_5$	$d_2$	$R_8$	
$R_3$	$c_2$	$R_6$	$d_3$	$R_9$	Used

## CHI-SQUARE DISTRIBUTION

CODE	KEYS	CODE	KEYS	CODE	KEYS
01	1	35	g	34	01
33 03	STO 3	02	$\pi$	81	$\div$
35 07	$g \chi^2 y$	31	f	33	STO
02	2	09	$\sqrt{x}$	71	x
81	$\div$	34 03	RCL 3	05	5
33 01	STO 1	71	x	02	2
31	f	33 03	STO 3	34 01	RCL 1
83	INT	84	R/S	71	x
35 00	$g \text{ LST } X$	23	LBL	33 06	STO 6
35 21	$g x \neq y$	12	B	01	1
22	GTO	33 02	STO 2	33 04	STO 4
01	1	34 01	RCL 1	23	LBL
01	1	01	1	03	3
51	—	51	—	34 02	RCL 2
35	g	35	g	34 06	RCL 6
03	$n!$	05	$y^x$	02	2
33 03	STO 3	34 02	RCL 2	61	+
84	R/S	02	2	33 06	STO 6
23	LBL	81	$\div$	81	$\div$
01	1	42	CHS	34 04	RCL 4
83	•	32	$f^{-1}$	71	x
05	5	07	LN	33 04	STO 4
35 23	$g x = y$	71	x	61	+
22	GTO	02	2	35 21	$g x \neq y$
02	2	34 01	RCL 1	22	GTO
35 07	$g x \leftarrow y$	35	g	03	3
01	1	05	$y^x$	34 05	RCL 5
51	—	81	$\div$	71	x
33	STO	34 03	RCL 3	84	R/S
71	x	81	$\div$	35 01	g NOP
03	3	33 05	STO 5		
22	GTO	84	R/S		
01	1	23	LBL		
23	LBL	13	C		
02	2	34 02	RCL 2		

<b>R<sub>1</sub></b>	$\nu/2$	<b>R<sub>4</sub></b>	Used	<b>R<sub>7</sub></b>	
<b>R<sub>2</sub></b>	x	<b>R<sub>5</sub></b>	Used	<b>R<sub>8</sub></b>	
<b>R<sub>3</sub></b>	$1, \Gamma(\nu/2)$	<b>R<sub>6</sub></b>	Used	<b>R<sub>9</sub></b>	Used

## t DISTRIBUTION

CODE	KEYS
23	LBL
11	A
33 01	STO 1
35	g
42	RAD
31	f
09	$\sqrt{x}$
81	$\div$
32	$f^{-1}$
06	TAN
33 02	STO 2
34 01	RCL 1
02	2
81	$\div$
31	f
83	INT
35 00	g LST X
35 21	g $x \neq y$
22	GTO
02	2
00	0
33 05	STO 5
23	LBL
12	B
34 02	RCL 2
31	f
05	COS
32	$f^{-1}$
09	$\sqrt{x}$
33 03	STO 3
34 02	RCL 2
31	f
04	SIN
33 04	STO 4
34 01	RCL 1

CODE	KEYS
02	2
35 23	g x=y
34 04	RCL 4
24	RTN
81	$\div$
01	1
51	$-$
33 08	STO 8
01	1
33 06	STO 6
23	LBL
01	1
34 03	RCL 3
71	x
34 05	RCL 5
01	1
61	+
71	x
35 00	g LST X
01	1
61	+
33 05	STO 5
81	$\div$
33	STO
61	+
06	6
35	g
83	DSZ
22	GTO
01	1
34 06	RCL 6
34 04	RCL 4
71	x
24	RTN
23	LBL

CODE	KEYS
02	2
34 02	RCL 2
02	2
71	x
35	g
02	$\pi$
81	$\div$
33 07	STO 7
34 01	RCL 1
01	1
33 05	STO 5
33	STO
51	$-$
01	1
35 23	g x=y
34 07	RCL 7
24	RTN
12	B
34 02	RCL 2
31	f
05	COS
71	x
02	2
71	x
35	g
02	$\pi$
81	$\div$
34 07	RCL 7
61	+
24	RTN

<b>R<sub>1</sub></b>	$\nu$ or $\nu - 1$	<b>R<sub>4</sub></b>	$\sin \theta$	<b>R<sub>7</sub></b>	$2\theta/\pi$
<b>R<sub>2</sub></b>	$\theta$	<b>R<sub>5</sub></b>	Used	<b>R<sub>8</sub></b>	Used
<b>R<sub>3</sub></b>	$\cos^2 \theta$	<b>R<sub>6</sub></b>	Used	<b>R<sub>9</sub></b>	Used

## F DISTRIBUTION

CODE	KEYS	CODE	KEYS	CODE	KEYS
33 01	STO 1	35 23	g x=y	33	STO
24	RTN	34 04	RCL 4	61	+
23	LBL	24	RTN	05	5
12	B	01	1	35	g
33 02	STO 2	33 05	STO 5	83	DSZ
24	RTN	34 03	RCL 3	22	GTO
23	LBL	51	—	03	3
13	C	33 03	STO 3	23	LBL
41	↑	34 02	RCL 2	02	2
34 01	RCL 1	02	2	34 05	RCL 5
71	x	81	÷	34 04	RCL 4
34 02	RCL 2	71	x	71	x
61	+	33	STO	24	RTN
34 02	RCL 2	61	+	23	LBL
35 07	g x↔y	05	5	15	E
81	÷	35	g	34 01	RCL 1
33 03	STO 3	83	DSZ	34 02	RCL 2
24	RTN	22	GTO	33 01	STO 1
23	LBL	03	3	35 07	g x↔y
14	D	22	GTO	33 02	STO 2
34 03	RCL 3	02	2	01	1
34 02	RCL 2	23	LBL	34 03	RCL 3
02	2	03	3	51	—
33 07	STO 7	34 02	RCL 2	33 03	STO 3
81	÷	02	2	14	D
35	g	61	+	01	1
05	y <sup>x</sup>	33 02	STO 2	35 07	g x↔y
33 04	STO 4	34 07	RCL 7	51	—
34 01	RCL 1	02	2	84	R/S
02	2	61	+	35 01	g NOP
51	—	33 07	STO 7		
02	2	81	÷		
81	÷	34 03	RCL 3		
33 08	STO 8	71	x		
00	0	71	x		

<b>R<sub>1</sub></b>	$\nu_1$ or $\nu_2$	<b>R<sub>4</sub></b>	$t^{\nu_2/2}$ or $t^{\nu_1/2}$	<b>R<sub>7</sub></b>	Used
<b>R<sub>2</sub></b>	$\nu_2$ or $\nu_1$	<b>R<sub>5</sub></b>	Used	<b>R<sub>8</sub></b>	Used
<b>R<sub>3</sub></b>	$t, 1 - t$	<b>R<sub>6</sub></b>		<b>R<sub>9</sub></b>	Used

## BIVARIATE NORMAL DISTRIBUTION

CODE	KEYS	CODE	KEYS	CODE	KEYS
23	LBL	34 02	RCL 2	35 01	g NOP
11	A	34 04	RCL 4	35 01	g NOP
33 01	STO 1	71	x	35 01	g NOP
35 07	$g x \leftrightarrow y$	02	2	35 01	g NOP
33 06	STO 6	71	x	35 01	g NOP
24	RTN	34 05	RCL 5	35 01	g NOP
23	LBL	71	x	35 01	g NOP
12	B	51	-	35 01	g NOP
33 03	STO 3	01	1	35 01	g NOP
35 07	$g x \leftrightarrow y$	34 05	RCL 5	35 01	g NOP
33 07	STO 7	32	$f^{-1}$	35 01	g NOP
24	RTN	09	$\sqrt{x}$	35 01	g NOP
23	LBL	51	-	35 01	g NOP
13	C	33 08	STO 8	35 01	g NOP
33 05	STO 5	02	2	35 01	g NOP
24	RTN	71	x	35 01	g NOP
23	LBL	81	$\div$	35 01	g NOP
14	D	42	CHS	35 01	g NOP
35 07	$g x \leftrightarrow y$	32	$f^{-1}$	35 01	g NOP
34 06	RCL 6	07	LN	35 01	g NOP
51	-	34 08	RCL 8	35 01	g NOP
34 01	RCL 1	31	f	35 01	g NOP
81	$\div$	09	$\sqrt{x}$	35 01	g NOP
33 02	STO 2	34 01	RCL 1	35 01	g NOP
32	$f^{-1}$	71	x	35 01	g NOP
09	$\sqrt{x}$	34 03	RCL 3	35 01	g NOP
35 07	$g x \leftrightarrow y$	71	x	35 01	g NOP
34 07	RCL 7	02	2	35 01	g NOP
51	-	71	x	35 01	g NOP
34 03	RCL 3	35	g	35 01	g NOP
81	$\div$	02	$\pi$	35 01	g NOP
33 04	STO 4	71	x	35 01	g NOP
32	$f^{-1}$	81	$\div$	35 01	g NOP
09	$\sqrt{x}$	24	RTN	35 01	g NOP
61	+	35 01	g NOP	35 01	g NOP

$R_1$	$\sigma_1$	$R_4$	$(y - \mu_2)/\sigma_2$	$R_7$	$\mu_2$
$R_2$	$(x - \mu_1)/\sigma_1$	$R_5$	$\rho$	$R_8$	$1 - \rho^2$
$R_3$	$\sigma_2$	$R_6$	$\mu_1$	$R_9$	

## LOGARITHMIC NORMAL DISTRIBUTION

CODE	KEYS	CODE	KEYS	CODE	KEYS
23	LBL	71	x	35 01	g NOP
11	A	24	RTN	35 01	g NOP
33 02	STO 2	23	LBL	35 01	g NOP
35 07	$g \sqrt{x}$	15	E	35 01	g NOP
33 01	STO 1	33 03	STO 3	35 01	g NOP
32	$f^{-1}$	31	f	35 01	g NOP
07	LN	07	LN	35 01	g NOP
24	RTN	34 01	RCL 1	35 01	g NOP
23	LBL	51	—	35 01	g NOP
12	B	32	$f^{-1}$	35 01	g NOP
34 01	RCL 1	09	$\sqrt{x}$	35 01	g NOP
34 02	RCL 2	34 02	RCL 2	35 01	g NOP
51	—	81	$\div$	35 01	g NOP
32	$f^{-1}$	02	2	35 01	g NOP
07	LN	81	$\div$	35 01	g NOP
24	RTN	42	CHS	35 01	g NOP
23	LBL	32	$f^{-1}$	35 01	g NOP
13	C	07	LN	35 01	g NOP
34 01	RCL 1	35	g	35 01	g NOP
34 02	RCL 2	02	$\pi$	35 01	g NOP
02	2	02	2	35 01	g NOP
81	$\div$	71	x	35 01	g NOP
61	+	34 02	RCL 2	35 01	g NOP
32	$f^{-1}$	71	x	35 01	g NOP
07	LN	31	f	35 01	g NOP
24	RTN	09	$\sqrt{x}$	35 01	g NOP
23	LBL	81	$\div$	35 01	g NOP
14	D	34 03	RCL 3	35 01	g NOP
32	$f^{-1}$	81	$\div$	35 01	g NOP
09	$\sqrt{x}$	24	RTN	35 01	g NOP
34 02	RCL 2	35 01	g NOP		
32	$f^{-1}$	35 01	g NOP		
07	LN	35 01	g NOP		
01	1	35 01	g NOP		
51	—	35 01	g NOP		

<b>R<sub>1</sub></b>	m	<b>R<sub>4</sub></b>	<b>R<sub>7</sub></b>
<b>R<sub>2</sub></b>	$\sigma^2$	<b>R<sub>5</sub></b>	<b>R<sub>8</sub></b>
<b>R<sub>3</sub></b>	x	<b>R<sub>6</sub></b>	<b>R<sub>9</sub></b>

## WEIBULL DISTRIBUTION

CODE	KEYS	CODE	KEYS	CODE	KEYS
23	LBL	35 24	g x>y	35 01	g NOP
11	A	00	0	35 01	g NOP
33 02	STO 2	81	÷	35 01	g NOP
35 07	g x↔y	31	f	35 01	g NOP
33 01	STO 1	07	LN	35 01	g NOP
24	RTN	34 01	RCL 1	35 01	g NOP
23	LBL	81	÷	35 01	g NOP
12	B	42	CHS	35 01	g NOP
33 03	STO 3	34 02	RCL 2	35 01	g NOP
34 02	RCL 2	35	g	35 01	g NOP
35	g	04	¹/x	35 01	g NOP
05	y <sup>x</sup>	35	g	35 01	g NOP
34 01	RCL 1	05	y <sup>x</sup>	35 01	g NOP
71	x	24	RTN	35 01	g NOP
42	CHS	35 01	g NOP	35 01	g NOP
32	f <sup>-1</sup>	35 01	g NOP	35 01	g NOP
07	LN	35 01	g NOP	35 01	g NOP
33 04	STO 4	35 01	g NOP	35 01	g NOP
35 00	g LST X	35 01	g NOP	35 01	g NOP
42	CHS	35 01	g NOP	35 01	g NOP
34 03	RCL 3	35 01	g NOP	35 01	g NOP
81	÷	35 01	g NOP	35 01	g NOP
71	x	35 01	g NOP	35 01	g NOP
34 02	RCL 2	35 01	g NOP	35 01	g NOP
71	x	35 01	g NOP	35 01	g NOP
24	RTN	35 01	g NOP	35 01	g NOP
23	LBL	35 01	g NOP	35 01	g NOP
13	C	35 01	g NOP	35 01	g NOP
34 04	RCL 4	35 01	g NOP	35 01	g NOP
24	RTN	35 01	g NOP	35 01	g NOP
23	LBL	35 01	g NOP	35 01	g NOP
14	D	35 01	g NOP	35 01	g NOP
41	↑	35 01	g NOP	35 01	g NOP
01	1	35 01	g NOP	35 01	g NOP
35 07	g x↔y	35 01	g NOP	35 01	g NOP

<b>R<sub>1</sub></b>	a	<b>R<sub>4</sub></b>	Q(x)	<b>R<sub>7</sub></b>	
<b>R<sub>2</sub></b>	b	<b>R<sub>5</sub></b>		<b>R<sub>8</sub></b>	
<b>R<sub>3</sub></b>	x	<b>R<sub>6</sub></b>		<b>R<sub>9</sub></b>	Used

## BINOMIAL DISTRIBUTION

CODE	KEYS	CODE	KEYS	CODE	KEYS
23	LBL	00	0	61	+
11	A	33 07	STO 7	05	5
33 01	STO 1	35 23	g x=y	34 07	RCL 7
35 07	g x↔y	34 03	RCL 3	01	1
33 02	STO 2	24	RTN	61	+
33 04	STO 4	44	CLX	33 07	STO 7
01	1	34 01	RCL 1	34 06	RCL 6
51	—	35 07	g x↔y	35 23	g x=y
42	CHS	35 24	g x>y	34 04	RCL 4
34 01	RCL 1	00	0	24	RTN
35	g	81	÷	22	GTO
05	y <sup>x</sup>	32	f <sup>-1</sup>	01	1
33 03	STO 3	83	INT	23	LBL
00	0	00	0	13	C
34 02	RCL 2	35 21	g x≠y	34 06	RCL 6
35 22	g x≤y	00	0	00	0
00	0	81	÷	35 23	g x=y
81	÷	34 03	RCL 3	34 03	RCL 3
01	1	33 04	STO 4	24	RTN
34 02	RCL 2	33 05	STO 5	01	1
51	—	23	LBL	34 05	RCL 5
81	÷	01	1	35 24	g x>y
33 02	STO 2	34 01	RCL 1	35 07	g x↔y
34 01	RCL 1	34 07	RCL 7	35 01	g NOP
34 04	RCL 4	51	—	24	RTN
71	x	34 07	RCL 7	35 01	g NOP
24	RTN	01	1	35 01	g NOP
01	1	61	+	35 01	g NOP
34 04	RCL 4	81	÷	35 01	g NOP
51	—	34 02	RCL 2	35 01	g NOP
71	x	71	x		
84	R/S	34 04	RCL 4		
23	LBL	71	x		
12	B	33 04	STO 4		
33 06	STO 6	33	STO		

<b>R<sub>1</sub></b>	n	<b>R<sub>4</sub></b>	Used	<b>R<sub>7</sub></b>	Counter
<b>R<sub>2</sub></b>	p, p/(1 – p)	<b>R<sub>5</sub></b>	Used	<b>R<sub>8</sub></b>	
<b>R<sub>3</sub></b>	f(0)	<b>R<sub>6</sub></b>	x	<b>R<sub>9</sub></b>	Used

## NEGATIVE BINOMIAL DISTRIBUTION

CODE	KEYS	CODE	KEYS	CODE	KEYS
23	LBL	35 23	g x=y	34 04	RCL 4
11	A	34 03	RCL 3	24	RTN
33 01	STO 1	24	RTN	22	GTO
35 07	g x $\leftrightarrow$ y	35 07	g x $\leftrightarrow$ y	01	1
33 02	STO 2	32	f $^{-1}$	23	LBL
35 07	g x $\leftrightarrow$ y	83	INT	15	E
35	g	35 21	g x $\neq$ y	34 06	RCL 6
05	y $^x$	00	0	00	0
33 03	STO 3	81	$\div$	35 23	g x=y
01	1	34 03	RCL 3	34 03	RCL 3
34 02	RCL 2	33 04	STO 4	24	RTN
35 23	g x=y	33 05	STO 5	01	1
00	0	23	LBL	34 05	RCL 5
81	$\div$	01	1	35 24	g x>y
24	RTN	01	1	35 07	g x $\leftrightarrow$ y
23	LBL	34 02	RCL 2	35 01	g NOP
12	B	51	—	24	RTN
34 01	RCL 1	34 07	RCL 7	35 01	g NOP
34 02	RCL 2	34 01	RCL 1	35 01	g NOP
81	$\div$	61	+	35 01	g NOP
01	1	71	x	35 01	g NOP
34 02	RCL 2	34 07	RCL 7	35 01	g NOP
51	—	01	1	35 01	g NOP
71	x	61	+	35 01	g NOP
24	RTN	33 07	STO 7	35 01	g NOP
23	LBL	81	$\div$	35 01	g NOP
13	C	34 04	RCL 4	35 01	g NOP
34 02	RCL 2	71	x	35 01	g NOP
81	$\div$	33 04	STO 4	35 01	g NOP
24	RTN	33	STO	35 01	g NOP
23	LBL	61	+	35 01	g NOP
14	D	05	5		
33 06	STO 6	34 07	RCL 7		
00	0	34 06	RCL 6		
33 07	STO 7	35 23	g x=y		

<b>R<sub>1</sub></b>	r	<b>R<sub>4</sub></b>	Used	<b>R<sub>7</sub></b>	Counter
<b>R<sub>2</sub></b>	p	<b>R<sub>5</sub></b>	Used	<b>R<sub>8</sub></b>	
<b>R<sub>3</sub></b>	f(0)	<b>R<sub>6</sub></b>	x	<b>R<sub>9</sub></b>	Used

## HYPERGEOMETRIC DISTRIBUTION

CODE	KEYS	CODE	KEYS	CODE	KEYS
33 02	STO 2	33 06	STO 6	22	GTO
35 07	g x $\geq$ y	00	0	01	1
33 01	STO 1	33 08	STO 8	23	LBL
84	R/S	23	LBL	14	D
23	LBL	01	1	34 06	RCL 6
12	B	34 01	RCL 1	24	RTN
33 03	STO 3	51	—	23	LBL
34 02	RCL 2	34 08	RCL 8	15	E
35	g	34 03	RCL 3	34 01	RCL 1
03	n!	51	—	34 03	RCL 3
35 00	g LST X	71	x	71	x
34 03	RCL 3	34 08	RCL 8	34 01	RCL 1
51	—	01	1	34 02	RCL 2
35	g	61	+	61	+
03	n!	81	$\div$	33 08	STO 8
81	$\div$	35 00	g LST X	81	$\div$
34 01	RCL 1	34 02	RCL 2	84	R/S
34 02	RCL 2	34 03	RCL 3	34 02	RCL 2
61	+	51	—	71	x
35	g	61	+	34 08	RCL 8
03	n!	81	$\div$	81	$\div$
35 00	g LST X	34 05	RCL 5	34 08	RCL 8
34 03	RCL 3	71	x	34 03	RCL 3
51	—	33 05	STO 5	51	—
35	g	33	STO	71	x
03	n!	61	+	34 08	RCL 8
81	$\div$	06	6	01	1
81	$\div$	34 07	RCL 7	51	—
33 04	STO 4	01	1	81	$\div$
24	RTN	34 08	RCL 8	24	RTN
23	LBL	61	+		
13	C	33 08	STO 8		
33 07	STO 7	35 23	g x=y		
34 04	RCL 4	34 05	RCL 5		
33 05	STO 5	24	RTN		

<b>R<sub>1</sub></b>	a	<b>R<sub>4</sub></b>	f(0)	<b>R<sub>7</sub></b>	x
<b>R<sub>2</sub></b>	b	<b>R<sub>5</sub></b>	Used	<b>R<sub>8</sub></b>	Counter, a + b
<b>R<sub>3</sub></b>	n	<b>R<sub>6</sub></b>	Used	<b>R<sub>9</sub></b>	Used

## POISSON DISTRIBUTION

CODE	KEYS	CODE	KEYS	CODE	KEYS
23	LBL	01	1	35 01	g NOP
11	A	34 01	RCL 1	35 01	g NOP
41	↑	34 06	RCL 6	35 01	g NOP
00	0	01	1	35 01	g NOP
35 07	g $x \leftrightarrow y$	61	+	35 01	g NOP
35 22	g $x \leq y$	81	÷	35 01	g NOP
00	0	34 03	RCL 3	35 01	g NOP
81	÷	71	x	35 01	g NOP
33 01	STO 1	33 03	STO 3	35 01	g NOP
42	CHS	33	STO	35 01	g NOP
32	$f^{-1}$	61	+	35 01	g NOP
07	LN	04	4	35 01	g NOP
33 02	STO 2	34 06	RCL 6	35 01	g NOP
24	RTN	01	1	35 01	g NOP
23	LBL	61	+	35 01	g NOP
12	B	33 06	STO 6	35 01	g NOP
33 05	STO 5	34 05	RCL 5	35 01	g NOP
00	0	35 23	g $x = y$	35 01	g NOP
33 06	STO 6	34 03	RCL 3	35 01	g NOP
35 24	g $x > y$	24	RTN	35 01	g NOP
00	0	22	GTO	35 01	g NOP
81	÷	01	1	35 01	g NOP
35 23	g $x = y$	23	LBL	35 01	g NOP
34 02	RCL 2	13	C	35 01	g NOP
24	RTN	34 05	RCL 5	35 01	g NOP
35 07	g $x \leftrightarrow y$	00	0	35 01	g NOP
32	$f^{-1}$	35 23	g $x = y$	35 01	g NOP
83	INT	34 02	RCL 2	35 01	g NOP
35 21	g $x \neq y$	24	RTN	35 01	g NOP
00	0	01	1	35 01	g NOP
81	÷	34 04	RCL 4	35 01	g NOP
34 02	RCL 2	35 24	g $x > y$	35 01	g NOP
33 03	STO 3	35 07	g $x \geq y$	35 01	g NOP
33 04	STO 4	35 01	g NOP		
23	LBL	24	RTN		

<b>R<sub>1</sub></b>	$\lambda$	<b>R<sub>4</sub></b>	Used	<b>R<sub>7</sub></b>	
<b>R<sub>2</sub></b>	$f(0)$	<b>R<sub>5</sub></b>	x	<b>R<sub>8</sub></b>	
<b>R<sub>3</sub></b>	Used	<b>R<sub>6</sub></b>	Counter	<b>R<sub>9</sub></b>	Used

## LINEAR REGRESSION

CODE	KEYS	CODE	KEYS	CODE	KEYS
23	LBL	09	9	09	$\sqrt{x}$
11	A	71	x	24	RTN
34 06	RCL 6	34 05	RCL 5	23	LBL
34 02	RCL 2	34 04	RCL 4	15	E
34 04	RCL 4	32	$f^{-1}$	34 03	RCL 3
71	x	09	$\sqrt{x}$	34 02	RCL 2
34 01	RCL 1	34 01	RCL 1	32	$f^{-1}$
81	$\div$	81	$\div$	09	$\sqrt{x}$
51	—	51	—	34 01	RCL 1
33	STO	81	$\div$	81	$\div$
09	9	24	RTN	51	—
34 03	RCL 3	23	LBL	31	f
34 02	RCL 2	13	C	09	$\sqrt{x}$
32	$f^{-1}$	41	↑	81	$\div$
09	$\sqrt{x}$	34 07	RCL 7	34 03	RCL 3
34 01	RCL 1	71	x	34 01	RCL 1
81	$\div$	34 08	RCL 8	81	$\div$
51	—	61	+	31	f
81	$\div$	24	RTN	09	$\sqrt{x}$
33 07	STO 7	23	LBL	35 07	$g \ x \rightarrow y$
34 04	RCL 4	14	D	71	x
34 07	RCL 7	34 05	RCL 5	84	R/S
34 02	RCL 2	34 08	RCL 8	35 00	$g \ LST \ X$
71	x	34 04	RCL 4	24	RTN
51	—	71	x	35 01	$g \ NOP$
34 01	RCL 1	51	—	35 01	$g \ NOP$
81	$\div$	34 06	RCL 6	35 01	$g \ NOP$
33 08	STO 8	34 07	RCL 7	35 01	$g \ NOP$
84	R/S	71	x	35 01	$g \ NOP$
34 07	RCL 7	51	—	35 01	$g \ NOP$
24	RTN	34 01	RCL 1	35 01	$g \ NOP$
23	LBL	02	2		
12	B	51	—		
34 07	RCL 7	81	$\div$		
34	RCL	31	f		

<b>R<sub>1</sub></b>	n	<b>R<sub>4</sub></b>	$\Sigma y_i$	<b>R<sub>7</sub></b>	$a_1$
<b>R<sub>2</sub></b>	$\Sigma x_i$	<b>R<sub>5</sub></b>	$\Sigma y_i^2$	<b>R<sub>8</sub></b>	$a_0$
<b>R<sub>3</sub></b>	$\Sigma x_i^2$	<b>R<sub>6</sub></b>	$\Sigma x_i y_i$	<b>R<sub>9</sub></b>	Used

## EXPONENTIAL CURVE FIT

CODE	KEYS	CODE	KEYS	CODE	KEYS
23	LBL	34 06	RCL 6	09	9
11	A	34 02	RCL 2	71	x
31	f	34 04	RCL 4	34 05	RCL 5
07	LN	71	x	34 04	RCL 4
33 07	STO 7	34 01	RCL 1	32	$f^{-1}$
33	STO	81	$\div$	09	$\sqrt{x}$
61	+	51	—	34 01	RCL 1
04	4	33	STO	81	$\div$
32	$f^{-1}$	09	9	51	—
09	$\sqrt{x}$	34 03	RCL 3	81	$\div$
33	STO	34 02	RCL 2	24	RTN
61	+	32	$f^{-1}$	23	LBL
05	5	09	$\sqrt{x}$	14	D
35 07	$g x \leftrightarrow y$	34 01	RCL 1	41	$\uparrow$
33	STO	81	$\div$	34 07	RCL 7
61	+	51	—	71	x
02	2	81	$\div$	32	$f^{-1}$
32	$f^{-1}$	33 07	STO 7	07	LN
09	$\sqrt{x}$	34 04	RCL 4	34 08	RCL 8
33	STO	34 07	RCL 7	71	x
61	+	34 02	RCL 2	24	RTN
03	3	71	x	35 01	$g NOP$
35 00	$g LST X$	51	—	35 01	$g NOP$
34 07	RCL 7	34 01	RCL 1	35 01	$g NOP$
71	x	81	$\div$	35 01	$g NOP$
33	STO	32	$f^{-1}$	35 01	$g NOP$
61	+	07	LN	35 01	$g NOP$
06	6	33 08	STO 8	35 01	$g NOP$
34 01	RCL 1	84	R/S	35 01	$g NOP$
01	1	34 07	RCL 7	35 01	$g NOP$
61	+	24	RTN	35 01	$g NOP$
33 01	STO 1	23	LBL		
24	RTN	13	C		
23	LBL	34 07	RCL 7		
12	B	34	RCL		

<b>R<sub>1</sub></b>	n	<b>R<sub>4</sub></b>	$\Sigma \ln y_i$	<b>R<sub>7</sub></b>	$\ln y_i, b$
<b>R<sub>2</sub></b>	$\Sigma x_i$	<b>R<sub>5</sub></b>	$\Sigma (\ln y_i)^2$	<b>R<sub>8</sub></b>	a
<b>R<sub>3</sub></b>	$\Sigma x_i^2$	<b>R<sub>6</sub></b>	$\Sigma x_i \ln y_i$	<b>R<sub>9</sub></b>	Used

## POWER CURVE FIT

CODE	KEYS	CODE	KEYS	CODE	KEYS
23	LBL	23	LBL	34 07	RCL 7
11	A	12	B	34	RCL
31	f	34 06	RCL 6	09	9
07	LN	34 02	RCL 2	71	x
33 07	STO 7	34 04	RCL 4	34 05	RCL 5
33	STO	71	x	34 04	RCL 4
61	+	34 01	RCL 1	32	$f^{-1}$
04	4	81	$\div$	09	$\sqrt{x}$
32	$f^{-1}$	51	-	34 01	RCL 1
09	$\sqrt{x}$	33	STO	81	$\div$
33	STO	09	9	51	-
61	+	34 03	RCL 3	81	$\div$
05	5	34 02	RCL 2	24	RTN
35 07	$g \times \leftrightarrow y$	32	$f^{-1}$	23	LBL
31	f	09	$\sqrt{x}$	14	D
07	LN	34 01	RCL 1	41	$\uparrow$
33	STO	81	$\div$	34 07	RCL 7
61	+	51	-	35	g
02	2	81	$\div$	05	$y^x$
32	$f^{-1}$	33 07	STO 7	34 08	RCL 8
09	$\sqrt{x}$	34 04	RCL 4	71	x
33	STO	34 07	RCL 7	24	RTN
61	+	34 02	RCL 2	35 01	g NOP
03	3	71	x	35 01	g NOP
35 00	$g LST X$	51	-	35 01	g NOP
34 07	RCL 7	34 01	RCL 1	35 01	g NOP
71	x	81	$\div$	35 01	g NOP
33	STO	32	$f^{-1}$	35 01	g NOP
61	+	07	LN	35 01	g NOP
06	6	33 08	STO 8	35 01	g NOP
34 01	RCL 1	84	R/S	35 01	g NOP
01	1	34 07	RCL 7	35 01	g NOP
61	+	24	RTN	35 01	g NOP
33 01	STO 1	23	LBL	35 01	g NOP
24	RTN	13	C		

<b>R<sub>1</sub></b>	n	<b>R<sub>4</sub></b>	$\sum \ln y_i$	<b>R<sub>7</sub></b>	$\ln y_i, b$
<b>R<sub>2</sub></b>	$\sum \ln x_i$	<b>R<sub>5</sub></b>	$\sum (\ln y_i)^2$	<b>R<sub>8</sub></b>	a
<b>R<sub>3</sub></b>	$\sum (\ln x_i)^2$	<b>R<sub>6</sub></b>	$\sum (\ln x_i) (\ln y_i)$	<b>R<sub>9</sub></b>	Used

## LOGARITHMIC CURVE FIT

CODE	KEYS	CODE	KEYS	CODE	KEYS
23	LBL	34 06	RCL 6	34 05	RCL 5
11	A	34 02	RCL 2	34 04	RCL 4
33 07	STO 7	34 04	RCL 4	32	$f^{-1}$
33	STO	71	x	09	$\sqrt{x}$
61	+	34 01	RCL 1	34 01	RCL 1
04	4	81	$\div$	81	$\div$
32	$f^{-1}$	51	-	51	-
09	$\sqrt{x}$	33	STO	81	$\div$
33	STO	09	9	24	RTN
61	+	34 03	RCL 3	23	LBL
05	5	34 02	RCL 2	14	D
35 07	$g \ x \leftrightarrow y$	32	$f^{-1}$	31	f
31	f	09	$\sqrt{x}$	07	LN
07	LN	34 01	RCL 1	34 07	RCL 7
33	STO	81	$\div$	71	x
61	+	51	-	34 08	RCL 8
02	2	81	$\div$	61	+
32	$f^{-1}$	33 07	STO 7	24	RTN
09	$\sqrt{x}$	34 04	RCL 4	35 01	g NOP
33	STO	34 07	RCL 7	35 01	g NOP
61	+	34 02	RCL 2	35 01	g NOP
03	3	71	x	35 01	g NOP
35 00	$g \ LST \ X$	51	-	35 01	g NOP
34 07	RCL 7	34 01	RCL 1	35 01	g NOP
71	x	81	$\div$	35 01	g NOP
33	STO	33 08	STO 8	35 01	g NOP
61	+	84	R/S	35 01	g NOP
06	6	34 07	RCL 7	35 01	g NOP
34 01	RCL 1	24	RTN	35 01	g NOP
01	1	23	LBL	35 01	g NOP
61	+	13	C	35 01	g NOP
33 01	STO 1	34 07	RCL 7		
24	RTN	34	RCL		
23	LBL	09	9		
12	B	71	x		

<b>R<sub>1</sub></b>	n	<b>R<sub>4</sub></b>	$\Sigma y_i$	<b>R<sub>7</sub></b>	$y_i, b$
<b>R<sub>2</sub></b>	$\Sigma \ln x_i$	<b>R<sub>5</sub></b>	$\Sigma y_i^2$	<b>R<sub>8</sub></b>	a
<b>R<sub>3</sub></b>	$\Sigma (\ln x_i)^2$	<b>R<sub>6</sub></b>	$\Sigma y_i \ln x_i$	<b>R<sub>9</sub></b>	Used

LEAST SQUARES REGRESSION OF  $y = cx^a + dx^b$ 

CODE	KEYS	CODE	KEYS	CODE	KEYS
33 02	STO 2	61	+	71	x
35 07	$g x \leftrightarrow y$	05	5	34 06	RCL 6
33 01	STO 1	35 08	$g R \downarrow$	35 07	$g x \leftrightarrow y$
24	RTN	32	$f^{-1}$	51	-
23	LBL	09	$\sqrt{x}$	34 08	RCL 8
12	B	33	STO	81	$\div$
35 07	$g x \leftrightarrow y$	61	+	33	STO
33 03	STO 3	08	8	09	9
34 01	RCL 1	34	RCL	24	RTN
35	g	09	9	34 03	RCL 3
05	$y^x$	32	$f^{-1}$	84	R/S
41	$\uparrow$	09	$\sqrt{x}$	23	LBL
41	$\uparrow$	33	STO	14	D
35 09	$g R \uparrow$	61	+	41	$\uparrow$
71	x	07	7	41	$\uparrow$
33	STO	24	RTN	34 01	RCL 1
61	+	23	LBL	35	g
06	6	13	C	05	$y^x$
44	CLX	34 08	RCL 8	34	RCL
35 00	$g LST X$	34 04	RCL 4	09	9
34 03	RCL 3	71	x	71	x
34 02	RCL 2	34 06	RCL 6	35 07	$g x \leftrightarrow y$
35	g	34 05	RCL 5	34 02	RCL 2
05	$y^x$	71	x	35	g
33	STO	51	-	05	$y^x$
09	9	34 07	RCL 7	34 03	RCL 3
71	x	34 08	RCL 8	71	x
33	STO	71	x	61	+
61	+	34 05	RCL 5	24	RTN
04	4	32	$f^{-1}$	35 01	g NOP
44	CLX	09	$\sqrt{x}$		
34	RCL	51	-		
09	9	81	$\div$		
71	x	33 03	STO 3		
33	STO	34 05	RCL 5		

$R_1$	a	$R_4$	$\sum y_i x_i^b$	$R_7$	$\sum x_i^{2b}$
$R_2$	b	$R_5$	$\sum x_i^{a+b}$	$R_8$	$\sum x_i^{2a}$
$R_3$	$x_i, d$	$R_6$	$\sum x_i^a y_i$	$R_9$	$x_i^b, c$

## MULTIPLE LINEAR REGRESSION (CARD 1)

CODE	KEYS	CODE	KEYS	CODE	KEYS
23	LBL	71	x	51	—
11	A	33	STO	04	4
41	↑	61	+	44	CLX
35 08	g R↓	01	1	35 00	g LST X
33	STO	44	CLX	35 07	g x↔y
61	+	35 00	g LST X	33	STO
09	9	71	x	51	—
35 07	g x↔y	33	STO	07	7
71	x	61	+	32	f⁻¹
33	STO	02	2	09	√x
61	+	01	1	33	STO
03	3	34 05	RCL 5	51	—
44	CLX	61	+	06	6
35 00	g LST X	33 05	STO 5	44	CLX
33	STO	24	RTN	35 00	g LST X
61	+	23	LBL	71	x
08	8	12	B	33	STO
32	f⁻¹	41	↑	51	—
09	√x	35 08	g R↓	01	1
33	STO	33	STO	44	CLX
61	+	51	—	35 00	g LST X
04	4	09	9	71	x
44	CLX	35 07	g x↔y	33	STO
35 00	g LST X	71	x	51	—
35 07	g x↔y	33	STO	02	2
33	STO	51	—	34 05	RCL 5
61	+	03	3	01	1
07	7	44	CLX	51	—
32	f⁻¹	35 00	g LST X	33 05	STO 5
09	√x	33	STO	24	RTN
33	STO	51	—		
61	+	08	8		
06	6	32	f⁻¹		
44	CLX	09	√x		
35 00	g LST X	33	STO		

<b>R<sub>1</sub></b>	$\sum x_i y_i$	<b>R<sub>4</sub></b>	$\sum y_i^2$	<b>R<sub>7</sub></b>	$\sum x_i$
<b>R<sub>2</sub></b>	$\sum x_i z_i$	<b>R<sub>5</sub></b>	n	<b>R<sub>8</sub></b>	$\sum y_i$
<b>R<sub>3</sub></b>	$\sum y_i z_i$	<b>R<sub>6</sub></b>	$\sum x_i^2$	<b>R<sub>9</sub></b>	$\sum z_i$

## MULTIPLE LINEAR REGRESSION (CARD 2)

CODE	KEYS	CODE	KEYS	CODE	KEYS
23	LBL	51	—	34 02	RCL 2
11	A	33 02	STO 2	34 07	RCL 7
34 05	RCL 5	71	x	71	x
34 06	RCL 6	34 03	RCL 3	51	—
71	x	35 07	g $x \leftrightarrow y$	34 05	RCL 5
34 07	RCL 7	51	—	81	÷
32	$f^{-1}$	34 06	RCL 6	33 01	STO 1
09	$\sqrt{x}$	34 05	RCL 5	84	R/S
51	—	34 04	RCL 4	23	LBL
33 06	STO 6	71	x	12	B
34 05	RCL 5	34 08	RCL 8	34 02	RCL 2
34 03	RCL 3	32	$f^{-1}$	84	R/S
71	x	09	$\sqrt{x}$	23	LBL
34 08	RCL 8	51	—	13	C
34	RCL	71	x	34 03	RCL 3
09	9	34 01	RCL 1	84	R/S
71	x	32	$f^{-1}$	23	LBL
51	—	09	$\sqrt{x}$	14	D
71	x	51	—	41	↑
33 03	STO 3	81	÷	34 03	RCL 3
34 05	RCL 5	33 03	STO 3	71	x
34 01	RCL 1	34 02	RCL 2	35 07	g $x \leftrightarrow y$
71	x	34 01	RCL 1	34 02	RCL 2
34 07	RCL 7	34 03	RCL 3	71	x
34 08	RCL 8	71	x	61	+
71	x	51	—	34 01	RCL 1
51	—	34 06	RCL 6	61	+
33 01	STO 1	81	÷	24	RTN
34 05	RCL 5	33 02	STO 2	35 01	g NOP
34 02	RCL 2	34	RCL	35 01	g NOP
71	x	09	9		
34 07	RCL 7	34 03	RCL 3		
34	RCL	34 08	RCL 8		
09	9	71	x		
71	x	51	—		

<b>R<sub>1</sub></b>	Used	<b>R<sub>4</sub></b>	$\sum y_i^2$	<b>R<sub>7</sub></b>	$\sum x_i$
<b>R<sub>2</sub></b>	Used	<b>R<sub>5</sub></b>	n	<b>R<sub>8</sub></b>	$\sum y_i$
<b>R<sub>3</sub></b>	Used	<b>R<sub>6</sub></b>	$\sum x_i^2, n \sum x_i^2 - (\sum x_i)^2$	<b>R<sub>9</sub></b>	$\sum z_i$

## PARABOLIC CURVE FIT

CODE	KEYS	CODE	KEYS	CODE	KEYS
23	LBL	61	+	71	x
11	A	04	4	33	STO
33	STO	34 05	RCL 5	51	—
61	+	01	1	01	1
09	9	61	+	35 00	g LST X
35 07	g x↔y	33 05	STO 5	71	x
33	STO	24	RTN	33	STO
61	+	23	LBL	51	—
07	7	12	B	04	4
71	x	33	STO	34 05	RCL 5
33	STO	51	—	01	1
61	+	09	9	51	—
02	2	35 07	g x↔y	33 05	STO 5
35 00	g LST X	33	STO	24	RTN
71	x	51	—	23	LBL
33	STO	07	7	13	C
61	+	71	x	33 04	STO 4
03	3	33	STO	34 03	RCL 3
35 00	g LST X	51	—	71	x
32	f <sup>-1</sup>	02	2	34 02	RCL 2
09	√x	35 00	g LST X	61	+
33	STO	71	x	34 04	RCL 4
61	+	33	STO	71	x
06	6	51	—	34 01	RCL 1
33	STO	03	3	61	+
61	+	35 00	g LST X	24	RTN
08	8	32	f <sup>-1</sup>	35 01	g NOP
35 00	g LST X	09	√x	35 01	g NOP
71	x	33	STO	35 01	g NOP
33	STO	51	—	35 01	g NOP
61	+	06	6		
01	1	33	STO		
35 00	g LST X	51	—		
71	x	08	8		
33	STO	35 00	g LST X		

<b>R<sub>1</sub></b>	$\sum x_i^3, a_0$	<b>R<sub>4</sub></b>	$\sum x_i^4, x$	<b>R<sub>7</sub></b>	$\sum x_i$
<b>R<sub>2</sub></b>	$\sum x_i y_i, a_1$	<b>R<sub>5</sub></b>	n	<b>R<sub>8</sub></b>	$\sum x_i^2$
<b>R<sub>3</sub></b>	$\sum x_i^2 y_i, a_2$	<b>R<sub>6</sub></b>	$\sum x_i^2$	<b>R<sub>9</sub></b>	$\sum y_i$

## PAIRED t STATISTIC

CODE	KEYS	CODE	KEYS	CODE	KEYS
23	LBL	51	—	24	RTN
11	A	34 01	RCL 1	35 01	g NOP
00	0	01	1	35 01	g NOP
33 01	STO 1	51	—	35 01	g NOP
33 02	STO 2	81	÷	35 01	g NOP
33 03	STO 3	31	f	35 01	g NOP
24	RTN	09	$\sqrt{x}$	35 01	g NOP
23	LBL	24	RTN	35 01	g NOP
12	B	23	LBL	35 01	g NOP
51	—	14	D	35 01	g NOP
33	STO	34 01	RCL 1	35 01	g NOP
61	+	31	f	35 01	g NOP
02	2	09	$\sqrt{x}$	35 01	g NOP
32	$f^{-1}$	81	÷	35 01	g NOP
09	$\sqrt{x}$	81	÷	35 01	g NOP
33	STO	84	R/S	35 01	g NOP
61	+	34 01	RCL 1	35 01	g NOP
03	3	01	1	35 01	g NOP
34 01	RCL 1	51	—	35 01	g NOP
01	1	24	RTN	35 01	g NOP
61	+	23	LBL	35 01	g NOP
33 01	STO 1	15	E	35 01	g NOP
24	RTN	51	—	35 01	g NOP
23	LBL	33	STO	35 01	g NOP
13	C	51	—	35 01	g NOP
34 02	RCL 2	02	2	35 01	g NOP
34 01	RCL 1	32	$f^{-1}$	35 01	g NOP
81	÷	09	$\sqrt{x}$	35 01	g NOP
84	R/S	33	STO	35 01	g NOP
34 03	RCL 3	51	—	35 01	g NOP
34 02	RCL 2	03	3	35 01	g NOP
32	$f^{-1}$	34 01	RCL 1	35 01	g NOP
09	$\sqrt{x}$	01	1	35 01	g NOP
34 01	RCL 1	51	—	35 01	g NOP
81	÷	33 01	STO 1		

<b>R<sub>1</sub></b>	n	<b>R<sub>4</sub></b>	<b>R<sub>7</sub></b>
<b>R<sub>2</sub></b>	$\Sigma D_i$	<b>R<sub>5</sub></b>	<b>R<sub>8</sub></b>
<b>R<sub>3</sub></b>	$\Sigma D_i^2$	<b>R<sub>6</sub></b>	<b>R<sub>9</sub></b>

## t STATISTIC FOR TWO MEANS

CODE	KEYS	CODE	KEYS	CODE	KEYS
23	LBL	14	D	34 05	RCL 5
11	A	34 06	RCL 6	34 04	RCL 4
00	0	34 05	RCL 5	81	÷
33 01	STO 1	32	f <sup>-1</sup>	34 02	RCL 2
33 02	STO 2	09	√x	34 01	RCL 1
33 03	STO 3	34 04	RCL 4	81	÷
24	RTN	81	÷	51	—
23	LBL	51	—	34 07	RCL 7
12	B	34 03	RCL 3	51	—
33	STO	61	+	35 07	g x↔y
61	+	34 02	RCL 2	81	÷
02	2	32	f <sup>-1</sup>	84	R/S
32	f <sup>-1</sup>	09	√x	34 08	RCL 8
09	√x	34 01	RCL 1	24	RTN
33	STO	81	÷	23	LBL
61	+	51	—	15	E
03	3	34 01	RCL 1	33	STO
34 01	RCL 1	34 04	RCL 4	51	—
01	1	61	+	02	2
61	+	02	2	32	f <sup>-1</sup>
33 01	STO 1	51	—	09	√x
24	RTN	33 08	STO 8	33	STO
23	LBL	81	÷	51	—
13	C	31	f	03	3
33 07	STO 7	09	√x	34 01	RCL 1
84	R/S	01	1	01	1
34 01	RCL 1	34 01	RCL 1	51	—
33 04	STO 4	81	÷	33 01	STO 1
34 02	RCL 2	01	1	24	RTN
33 05	STO 5	34 04	RCL 4	35 01	g NOP
34 03	RCL 3	81	÷		
33 06	STO 6	61	+		
11	A	31	f		
24	RTN	09	√x		
23	LBL	71	x		

<b>R<sub>1</sub></b>	$n_1, n_2$	<b>R<sub>4</sub></b>	$n_1$	<b>R<sub>7</sub></b>	D
<b>R<sub>2</sub></b>	$\Sigma x_i, \Sigma y_i$	<b>R<sub>5</sub></b>	$\Sigma x_i$	<b>R<sub>8</sub></b>	$n_1 + n_2 - 2$
<b>R<sub>3</sub></b>	$\Sigma x_i^2, \Sigma y_i^2$	<b>R<sub>6</sub></b>	$\Sigma x_i^2$	<b>R<sub>9</sub></b>	

## CHI-SQUARE EVALUATION

CODE	KEYS	CODE	KEYS	CODE	KEYS
00	0	01	1	15	E
33 01	STO 1	51	—	31	f
33 02	STO 2	33 01	STO 1	61	TF 1
33 03	STO 3	24	RTN	22	GTO
32	f <sup>-1</sup>	23	LBL	01	1
51	SF 1	13	C	34 02	RCL 2
84	R/S	33	STO	24	RTN
23	LBL	61	+	23	LBL
11	A	02	2	01	1
33 03	STO 3	32	f <sup>-1</sup>	34 01	RCL 1
51	—	09	$\sqrt{x}$	34 03	RCL 3
32	f <sup>-1</sup>	33	STO	71	x
09	$\sqrt{x}$	61	+	34 02	RCL 2
34 03	RCL 3	03	3	81	$\div$
81	$\div$	34 01	RCL 1	34 02	RCL 2
33	STO	01	1	51	—
61	+	61	+	84	R/S
02	2	33 01	STO 1	34 02	RCL 2
34 01	RCL 1	24	RTN	34 01	RCL 1
01	1	23	LBL	81	$\div$
61	+	14	D	24	RTN
33 01	STO 1	33	STO	35 01	g NOP
24	RTN	51	—	35 01	g NOP
23	LBL	02	2	35 01	g NOP
12	B	32	f <sup>-1</sup>	35 01	g NOP
33 03	STO 3	09	$\sqrt{x}$	35 01	g NOP
51	—	33	STO	35 01	g NOP
32	f <sup>-1</sup>	51	—	35 01	g NOP
09	$\sqrt{x}$	03	3	35 01	g NOP
34 03	RCL 3	34 01	RCL 1	35 01	g NOP
81	$\div$	01	1		
33	STO	51	—		
51	—	33 01	STO 1		
02	2	24	RTN		
34 01	RCL 1	23	LBL		

<b>R<sub>1</sub></b>	n	<b>R<sub>4</sub></b>	<b>R<sub>7</sub></b>
<b>R<sub>2</sub></b>	Used	<b>R<sub>5</sub></b>	<b>R<sub>8</sub></b>
<b>R<sub>3</sub></b>	Used	<b>R<sub>6</sub></b>	<b>R<sub>9</sub></b>

## 2 x k CONTINGENCY TABLE

CODE	KEYS	CODE	KEYS	CODE	KEYS
23	LBL	05	5	01	1
11	A	35 08	g R↓	51	—
31	f	34 08	RCL 8	24	RTN
43	REG	41	↑	23	LBL
24	RTN	71	x	15	E
23	LBL	35 07	g x↔y	34 07	RCL 7
12	B	81	÷	34 04	RCL 4
33 08	STO 8	33	STO	34 07	RCL 7
35 07	g x↔y	61	+	61	+
33 07	STO 7	06	6	81	÷
41	↑	01	1	31	f
41	↑	34 03	RCL 3	09	√x
71	x	33 03	STO 3	24	RTN
35 08	g R↓	24	RTN	35 01	g NOP
33	STO	23	LBL	35 01	g NOP
61	+	13	C	35 01	g NOP
01	1	34 04	RCL 4	35 01	g NOP
33	STO	34 05	RCL 5	35 01	g NOP
61	+	71	x	35 01	g NOP
04	4	34 01	RCL 1	35 01	g NOP
35 07	g x↔y	81	÷	35 01	g NOP
33	STO	34 04	RCL 4	35 01	g NOP
61	+	34 06	RCL 6	35 01	g NOP
02	2	34 02	RCL 2	35 01	g NOP
33	STO	81	÷	35 01	g NOP
61	+	71	x	35 01	g NOP
04	4	61	+	35 01	g NOP
61	+	34 04	RCL 4	35 01	g NOP
41	↑	51	—	35 01	g NOP
35 08	g R↓	33 07	STO 7	35 01	g NOP
35 07	g x↔y	24	RTN	35 01	g NOP
35 08	g R↓	23	LBL	35 01	g NOP
81	÷	14	D	35 01	g NOP
33	STO	34 03	RCL 3	35 01	g NOP
61	+				

<b>R<sub>1</sub></b>	N <sub>A</sub>	<b>R<sub>4</sub></b>	N	<b>R<sub>7</sub></b>	a <sub>i</sub> , χ <sup>2</sup>
<b>R<sub>2</sub></b>	N <sub>B</sub>	<b>R<sub>5</sub></b>	Σ a <sub>i</sub> <sup>2</sup> /N <sub>i</sub>	<b>R<sub>8</sub></b>	b <sub>i</sub>
<b>R<sub>3</sub></b>	k	<b>R<sub>6</sub></b>	Σ b <sub>i</sub> <sup>2</sup> /N <sub>i</sub>	<b>R<sub>9</sub></b>	0

## BARTLETT'S CHI-SQUARE STATISTIC

CODE	KEYS	CODE	KEYS	CODE	KEYS
23	LBL	33 05	STO 5	51	—
11	A	24	RTN	03	3
31	f	23	LBL	35	g
43	REG	13	C	04	$\frac{1}{x}$
24	RTN	34 08	RCL 8	33	STO
23	LBL	34 03	RCL 3	51	—
12	B	81	÷	04	4
33 01	STO 1	31	f	35 08	g R↓
33	STO	07	LN	41	↑
61	+	34 03	RCL 3	41	↑
03	3	71	x	34 01	RCL 1
35	g	34 07	RCL 7	71	x
04	$\frac{1}{x}$	51	—	33	STO
33	STO	34 04	RCL 4	51	—
61	+	34 03	RCL 3	08	8
04	4	35	g	35 07	g $x \leftrightarrow y$
35 08	g R↓	04	$\frac{1}{x}$	31	f
41	↑	51	—	07	LN
41	↑	34 05	RCL 5	34 01	RCL 1
34 01	RCL 1	01	1	71	x
71	x	51	—	33	STO
33	STO	33 02	STO 2	51	—
61	+	03	3	07	7
08	8	71	x	34 05	RCL 5
35 07	g $x \leftrightarrow y$	81	÷	01	1
31	f	01	1	51	—
07	LN	61	+	33 05	STO 5
34 01	RCL 1	81	÷	24	RTN
71	x	84	R/S	35 01	g NOP
33	STO	34 02	RCL 2	35 01	g NOP
61	+	24	RTN		
07	7	23	LBL		
34 05	RCL 5	14	D		
01	1	33 01	STO 1		
61	+	33	STO		

<b>R<sub>1</sub></b>	$f_i$	<b>R<sub>4</sub></b>	$\sum 1/f_i$	<b>R<sub>7</sub></b>	$\sum f_i \ln s_i^2$
<b>R<sub>2</sub></b>	df	<b>R<sub>5</sub></b>	k	<b>R<sub>8</sub></b>	$\sum f_i s_i^2$
<b>R<sub>3</sub></b>	$\Sigma f_i$	<b>R<sub>6</sub></b>	0	<b>R<sub>9</sub></b>	0

## SPEARMAN'S RANK CORRELATION COEFFICIENT

CODE	KEYS	CODE	KEYS	CODE	KEYS
23	LBL	23	LBL	35 01	g NOP
11	A	14	D	35 01	g NOP
00	0	34 01	RCL 1	35 01	g NOP
33 01	STO 1	01	1	35 01	g NOP
33 02	STO 2	51	—	35 01	g NOP
24	RTN	31	f	35 01	g NOP
23	LBL	09	$\sqrt{x}$	35 01	g NOP
12	B	71	x	35 01	g NOP
51	—	24	RTN	35 01	g NOP
32	$f^{-1}$	23	LBL	35 01	g NOP
09	$\sqrt{x}$	15	E	35 01	g NOP
33	STO	51	—	35 01	g NOP
61	+	32	$f^{-1}$	35 01	g NOP
02	2	09	$\sqrt{x}$	35 01	g NOP
34 01	RCL 1	33	STO	35 01	g NOP
01	1	51	—	35 01	g NOP
61	+	02	2	35 01	g NOP
33 01	STO 1	34 01	RCL 1	35 01	g NOP
24	RTN	01	1	35 01	g NOP
23	LBL	51	—	35 01	g NOP
13	C	33 01	STO 1	35 01	g NOP
01	1	24	RTN	35 01	g NOP
34 02	RCL 2	35 01	g NOP	35 01	g NOP
06	6	35 01	g NOP	35 01	g NOP
71	x	35 01	g NOP	35 01	g NOP
34 01	RCL 1	35 01	g NOP	35 01	g NOP
32	$f^{-1}$	35 01	g NOP	35 01	g NOP
09	$\sqrt{x}$	35 01	g NOP	35 01	g NOP
01	1	35 01	g NOP	35 01	g NOP
51	—	35 01	g NOP	35 01	g NOP
34 01	RCL 1	35 01	g NOP	35 01	g NOP
71	x	35 01	g NOP	35 01	g NOP
81	÷	35 01	g NOP	35 01	g NOP
51	—	35 01	g NOP	35 01	g NOP
24	RTN	35 01	g NOP	35 01	g NOP

<b>R<sub>1</sub></b>	n	<b>R<sub>4</sub></b>	<b>R<sub>7</sub></b>
<b>R<sub>2</sub></b>	$\sum D_i^2$	<b>R<sub>5</sub></b>	<b>R<sub>8</sub></b>
<b>R<sub>3</sub></b>		<b>R<sub>6</sub></b>	<b>R<sub>9</sub></b>

## MANN-WHITNEY STATISTIC

CODE	KEYS	CODE	KEYS	CODE	KEYS
23	LBL	02	2	35 01	g NOP
11	A	81	÷	35 01	g NOP
33 02	STO 2	51	—	35 01	g NOP
00	0	34 01	RCL 1	35 01	g NOP
33 01	STO 1	34 02	RCL 2	35 01	g NOP
33 03	STO 3	61	+	35 01	g NOP
24	RTN	01	1	35 01	g NOP
23	LBL	61	+	35 01	g NOP
12	B	34 01	RCL 1	35 01	g NOP
33	STO	71	x	35 01	g NOP
61	+	34 02	RCL 2	35 01	g NOP
03	3	71	x	35 01	g NOP
34 01	RCL 1	01	1	35 01	g NOP
01	1	02	2	35 01	g NOP
61	+	81	÷	35 01	g NOP
33 01	STO 1	31	f	35 01	g NOP
24	RTN	09	$\sqrt{x}$	35 01	g NOP
23	LBL	81	÷	35 01	g NOP
13	C	24	RTN	35 01	g NOP
34 02	RCL 2	23	LBL	35 01	g NOP
34 01	RCL 1	15	E	35 01	g NOP
01	1	33	STO	35 01	g NOP
61	+	51	—	35 01	g NOP
02	2	03	3	35 01	g NOP
81	÷	34 01	RCL 1	35 01	g NOP
61	+	01	1	35 01	g NOP
71	x	51	—	35 01	g NOP
34 03	RCL 3	33 01	STO 1	35 01	g NOP
51	—	24	RTN	35 01	g NOP
24	RTN	35 01	g NOP	35 01	g NOP
23	LBL	35 01	g NOP	35 01	g NOP
14	D	35 01	g NOP	35 01	g NOP
34 01	RCL 1	35 01	g NOP	35 01	g NOP
34 02	RCL 2	35 01	g NOP	35 01	g NOP
71	x	35 01	g NOP	35 01	g NOP

<b>R<sub>1</sub></b>	n <sub>1</sub>	<b>R<sub>4</sub></b>	<b>R<sub>7</sub></b>
<b>R<sub>2</sub></b>	n <sub>2</sub>	<b>R<sub>5</sub></b>	<b>R<sub>8</sub></b>
<b>R<sub>3</sub></b>	$\Sigma R_i$	<b>R<sub>6</sub></b>	<b>R<sub>9</sub></b>

## KENDALL'S COEFFICIENT OF CONCORDANCE

CODE	KEYS	CODE	KEYS	CODE	KEYS
00	0	23	LBL	51	—
33 01	STO 1	13	C	71	x
33 02	STO 2	34 03	RCL 3	84	R/S
33 03	STO 3	01	1	34 04	RCL 4
33 04	STO 4	02	2	01	1
84	R/S	71	x	51	—
23	LBL	34 05	RCL 5	24	RTN
11	A	32	f <sup>-1</sup>	23	LBL
33	STO	09	√x	15	E
61	+	81	÷	33	STO
02	2	34 04	RCL 4	51	—
34 01	RCL 1	81	÷	02	2
01	1	34 04	RCL 4	34 01	RCL 1
61	+	32	f <sup>-1</sup>	01	1
33 01	STO 1	09	√x	51	—
24	RTN	01	1	33 01	STO 1
23	LBL	51	—	24	RTN
12	B	81	÷	35 01	g NOP
34 01	RCL 1	34 04	RCL 4	35 01	g NOP
33 05	STO 5	01	1	35 01	g NOP
34 02	RCL 2	61	+	35 01	g NOP
32	f <sup>-1</sup>	03	3	35 01	g NOP
09	√x	71	x	35 01	g NOP
33	STO	34 04	RCL 4	35 01	g NOP
61	+	01	1	35 01	g NOP
03	3	51	—	35 01	g NOP
34 04	RCL 4	81	÷	35 01	g NOP
01	1	51	—	35 01	g NOP
61	+	24	RTN	35 01	g NOP
33 04	STO 4	23	LBL	35 01	g NOP
00	0	14	D	35 01	g NOP
33 01	STO 1	34 05	RCL 5		
33 02	STO 2	71	x		
34 04	RCL 4	34 04	RCL 4		
24	RTN	01	1		

<b>R<sub>1</sub></b>	j	<b>R<sub>4</sub></b>	n	<b>R<sub>7</sub></b>
<b>R<sub>2</sub></b>	$\Sigma R_{ij}$	<b>R<sub>5</sub></b>	k	<b>R<sub>8</sub></b>
<b>R<sub>3</sub></b>	$\Sigma(\Sigma R_{ij})^2$	<b>R<sub>6</sub></b>		<b>R<sub>9</sub></b>

## BISERIAL CORRELATION COEFFICIENT

CODE	KEYS	CODE	KEYS	CODE	KEYS
23	LBL	01	1	35 01	g NOP
11	A	61	+	35 01	g NOP
31	f	33 03	STO 3	35 01	g NOP
43	REG	24	RTN	35 01	g NOP
33 01	STO 1	23	LBL	35 01	g NOP
24	RTN	14	D	35 01	g NOP
23	LBL	34 04	RCL 4	35 01	g NOP
12	B	33	STO	35 01	g NOP
33	STO	61	+	35 01	g NOP
61	+	05	5	35 01	g NOP
04	4	34 03	RCL 3	35 01	g NOP
32	$f^{-1}$	71	x	35 01	g NOP
09	$\sqrt{x}$	34 05	RCL 5	35 01	g NOP
33	STO	34 02	RCL 2	35 01	g NOP
61	+	71	x	35 01	g NOP
06	6	51	-	35 01	g NOP
01	1	34 03	RCL 3	35 01	g NOP
33	STO	81	$\div$	35 01	g NOP
61	+	34 01	RCL 1	35 01	g NOP
02	2	81	$\div$	35 01	g NOP
34 03	RCL 3	34 03	RCL 3	35 01	g NOP
61	+	34 06	RCL 6	35 01	g NOP
33 03	STO 3	71	x	35 01	g NOP
24	RTN	34 05	RCL 5	35 01	g NOP
23	LBL	32	$f^{-1}$	35 01	g NOP
13	C	09	$\sqrt{x}$	35 01	g NOP
33	STO	51	-	35 01	g NOP
61	+	31	f	35 01	g NOP
05	5	09	$\sqrt{x}$	35 01	g NOP
32	$f^{-1}$	81	$\div$	35 01	g NOP
09	$\sqrt{x}$	24	RTN	35 01	g NOP
33	STO	35 01	g NOP	35 01	g NOP
61	+	35 01	g NOP	35 01	g NOP
06	6	35 01	g NOP	35 01	g NOP
34 03	RCL 3	35 01	g NOP	35 01	g NOP

<b>R<sub>1</sub></b>	a	<b>R<sub>4</sub></b>	$\Sigma' y_i$	<b>R<sub>7</sub></b>	0
<b>R<sub>2</sub></b>	$n_1$	<b>R<sub>5</sub></b>	$\Sigma y_i$	<b>R<sub>8</sub></b>	0
<b>R<sub>3</sub></b>	n	<b>R<sub>6</sub></b>	$\Sigma y_i^2$	<b>R<sub>9</sub></b>	0















Sales and service from 172 offices in 65 countries.  
**19310 Pruneridge Avenue, Cupertino, California 95014**